

Aetos Minoxidil Topical Solution 5%w/v

To be supplied only on the prescription of a registered medical practitioner

Composition

Each ml contains 50mg of minoxidil (5%w/v) Contains ethanol (51%w/v)

Product Description

Aetos Minoxidil Topical Solution 5%w/v is a cutaneous solution available as a clear, colourless to yellowish homogeneous solution, free from particles in suspension.

Pharmacodynamics

The mechanism by which cutaneous minoxidil and/or its metabolite stimulates hair growth is not yet clear. However, it is believed that minoxidil acts at a hair follicle level, by increasing the blood flow in the scalp due to local vasodilation, stimulating anagen growth, prolonging the anagen phase and stimulating anagen recovery from telogen phase. As a peripheral vasodilator, minoxidil enhances microcirculation to hair follicles. The Vascular Endothelial Growth Factor (VEGF) is stimulated by minoxidil and VEGF is responsible for the increased capillary fenestration, which indicating of high metabolic activity during the anagen phase.

Pharmacokinetics

Minoxidil percutaneous absorption is approximately 0.3% to 4.5% of the total dose applied, after cutaneous application of hydroalcoholic solutions at 2% containing propylene glycol. Minoxidil is not bound to plasma protein. It is distributed into breast milk. Minoxidil is extensively metabolised by the liver. It requires sulfation to become active, but the major metabolites is a glucuronide conjugate. Minoxidil is excreted predominantly in the urine mainly in the form of metabolites. Minoxidil and its metabolites are dialysable although the pharmacological effect is not reversed.

Indication

Aetos Minoxidil Topical Solution 5%w/v is exclusively indicated for the treatment of alopecia androgenetica in males and females.

Aetos Minoxidil Topical Solution 5%w/v stimulates hair growth and stabilizes hair loss in patients with alopecia androgenetic.

Dosage and Administration

Use Aetos Minoxidil Topical Solution 5%w/v only as directed. Do not apply Aetos Minoxidil Topical Solution 5%w/v to any other area of the body.

A total dose of 6 sprays (equivalent to 1 mL) Aetos Minoxidil Topical Solution 5%w/v should be applied twice per day to the scalp, beginning at the center of the affected area. This dose should be used regardless of the size of the affected area. The total daily dose should not exceed 2 mL.

Apply Aetos Minoxidil Topical Solution 5%w/v when the hair and scalp are thoroughly dry. After applying Aetos Minoxidil Topical Solution 5%w/v wash hands thoroughly.

Method of application:

According to the affected area and the application device provided.

A. Spray pump



Fig. 1

1. Remove the outer cap of the bottle.
2. Apply 1 ml of solution, by pressing the spray pump 6 times (Fig. 1).

B. Applicator



Fig. 2



Fig.3

1. Remove the outer cap of the bottle.
2. Remove the top part of the spray head (piece with the orifice), by pulling it up, (Fig. 2) and insert the applicator (Fig. 3 & 4).



Fig. 4



Fig.5

3. Apply 1 ml of solution, by pressing the applicator 6 times (Fig. 5).
- Remark: Kindly wash your hand after each application.

Contraindications

Not to be used by those who are known hypersensitivity to Minoxidil.

Should not be applied to inflamed scalp skin or areas affected by psoriasis, severe sunburn, or severe excoriations, because of the risk of increased absorption.

Warnings and Precautions

Topical Minoxidil is only indicated for the treatment of alopecia androgenetica and should not be used in other type of hair loss for example when there is no family history of hair loss, hair loss is sudden and/or patchy, hair loss is due to childbirth, or the reason for hair loss is unknown. Topical application of minoxidil should be restricted to the scalp.

Patients being treated for hypertension should be monitored if topical minoxidil is used concurrently.

Minoxidil topical solution will cause burning and irritation of the eye. In the event of accidental contact with sensitive surfaces (eye, abraded ski, mucous membranes), the area should be bathed with copious amounts of cool tap water.

Use in Children:

Safely and efficacy of minoxidil topical solution 5% in patients under 18 years of age have not been established.

Use in elderly:

Safely and efficacy of minoxidil topical solution 5% in patients over 65 years of age have not been established.

Interactions with Other Medicaments

Should not be used concomitantly with other medications applied topically on the scalp. Topical drugs eg: corticosteroid, tretinoin, dithranol or petrolatum which may alter the stratum corneum barrier and could increase absorption of minoxidil if apply concurrently.

Although it has not been demonstrated clinically, there exists the theoretical possibility of absorbed minoxidil potentiating orthostatic hypotension caused by peripheral vasodilators.

Pregnancy and Lactation

This product should not be used during pregnancy or lactation.

Pregnancy

There are no adequate and well-controlled studies in pregnant women. Studies in animal have shown a risk to the foetus at exposure levels that are very high compared to those intended for human exposure.

Lactation

Systemically absorbed minoxidil is secreted in human milk. Hence, Minoxidil should not be used by pregnant or nursing mothers

Side Effects:

Like all medicines, this medicine may cause side effects, although not everybody gets them.

Body System (SOC)	Frequency	Adverse Drug Reaction (Preferred Term)
Immune System Disorders	Common	Hypersensitivity reactions (including face oedema, generalised erythema, pruritus generalised, swelling face, and throat tightness)
	Not known	Angioedema (including lip oedema, lip swelling, oedema mouth, oropharyngeal swelling, pharyngeal oedema, swollen tongue and tongue oedema)
Psychiatric Disorders	Not known	Depressed mood
Nervous System Disorders	Very common	Headache
	Uncommon	Dizziness
Eye disorders	Not known	Eye irritation
Cardiac disorders	Common	Chest pain
	Uncommon	Palpitations
	Not known	Heart rate increased
Vascular disorders	Not known	Hypotension
Respiratory, thoracic and mediastinal disorders	Uncommon	Dyspnoea
Gastrointestinal Disorders	Uncommon	Nausea
	Not known	Vomiting
Skin and subcutaneous tissue disorders	Common	Hypertrichosis (unwanted non-scalp hair including facial hair growth in women) Pruritus (including rash pruritic generalised and eye pruritus) Rash (including macular, popular, generalised, vestibular and macular rash) Dermatitis (including contact, allergic, atopic and seborrhoeic dermatitis)
	Rare	Changes in hair
	Not known	Dry skin Skin exfoliation (including exfoliative rash and dermatitis exfoliative) Acne (acneiform rash) Temporary hair loss (see section 4.4) Changes in hair colour
General disorders and administration site conditions	Common	Oedema peripheral
	Not known	Application site reactions (These sometimes involve nearby structures like the ears and face and typically consist of pruritus, irritation, pain, rash, oedema, dry skin, erythema and rash erythematous but can sometimes be more severe and include exfoliation, dermatitis, blistering, bleeding and ulceration)
Investigations	Common	Weight increased

Symptoms and Treatment of Overdose

Increased systemic absorption of minoxidil may potentially occur if higher than recommended doses are applied to larger surface areas of the body or areas other than the scalp.

Accidental ingestion has the potential of producing systemic effects related to the pharmacological action of the drug. Signs and symptoms of minoxidil overdose would primarily be cardiovascular effects associated with sodium and water retention. Tachycardia, hypotension, dizziness and lethargy can also occur.

Treatment of minoxidil overdose should be symptomatic and supportive.

Fluid retention can be managed with appropriate diuretic therapy. Clinically significant tachycardia can be controlled by administration of beta-adrenergic blocking agent.

Effects on Ability to Drive and Use Machine

This product may cause dizziness or hypotension. If affected, patients should not drive or operate machinery.

Storage Conditions: Do not store above 30°C. Keep the bottle tightly closed. Protect from light.

Dosage Forms and Packaging Available

Aetos Minoxidil Topical Solution 5%w/v is available in 100 ml bottle. For further information, please consult your pharmacist/physician.

Name and Address of Manufacturer

MEDINFAR MANUFACTURING, S.A.
Parque Industrial Armando Martins Tavares,
Rua Outeiro Da Armada No 5,
Condeixa-A-Nova,
3150-194 Sebal, Portugal

Name and Address of Product Registration Holder

Aetos Pharma Sdn Bhd (1322949-D)
Add: No. 66B-Tingkat 2, Jalan Cerdas,
Taman Connaught,
56000 Kuala Lumpur, Malaysia

Arial
Title:12.5pt ,Body: 7pt
Date of revision: 06/11/2023

Product

Minoxidil Aetos 100 ml MY

Client

Laboratórios Medinfar

C.I.:

76011156rev01



  mm
190 290

 5 mm

  mm
3 190x36

TXT



1/1

● - Preto U

○ -

○ -

○ -

○ -

Rev.

18/01/2024 prova: 1

24/01/2024 prova: 2

I

I X

Component

IOR 50 gr/m2

Logistics

Production

Regulatory affairs

Quality Control