

LINCOFEED 4.4% POWDER

Description and Composition

LINCOFEED 4.4% POWDER is a buff coloured powder. Each kg contains 44g Lincomycin (as hydrochloride) as its active ingredient and 5g Calcium Propionate as its preservative.

Pharmacodynamics

Lincomycin is active mainly against Gram-positive bacteria. Lincomycin exerts its antibacterial action by inhibiting RNA-dependent protein synthesis by acting on the 50S subunit of the ribosome.

Pharmacokinetics

In chickens (1/sex/time point) given twice daily oral boluses of ^{14}C -lincomycin (simulating a dietary dose of 1 mg/bird/day) for 35 days approximately 75% of the administered dose was radiometrically detected in the excrement 1 to 3 days after dosing while approximately 30% was detectable by microbiological assay. When chickens (2/time point) were fed a diet containing 11 mg ^{14}C -lincomycin/kg for 35 days then 2 times 0.5 mg ^{14}C -lincomycin/day by oral bolus, the 'total' tissue residue concentrations in bile ranged from 5 194 to 10 μg equivalents/kg after 1 hour to 3 day withdrawal periods, respectively.

In pigs, oral doses of 4.4, 11 (the therapeutic dose), and 22 mg lincomycin/kg bw resulted in therapeutic serum concentrations being reached by 0.5 hour after dosing. Peak serum concentrations of 1.8, 3.9 and 5.1 mg/ml were reached within an hour and less than 4% of the plasma lincomycin was protein bound. In studies where pigs were administered lincomycin by intravenous or oral routes the distribution of liver metabolites were determined by thin layer chromatography (TLC) to have been quantitatively the same. An oral bolus or intravenous injection of 10 mg lincomycin/kg bw resulted in plasma $t_{1/2}$ values of 3.4 and 2.0 hours, respectively. In another pig study, the half-life for liver and kidney tissues after oral administration of lincomycin was reported to have been 24 and 29 hours, respectively.

Indication

Effective strain of *Mycoplasma hyopneumoniae*, *Mycoplasma hyorhinis*, *Clostridium perfringens*, and *Brachyspira hyodysenteriae* susceptible to this drug.

Pigs: Swine mycoplasma pneumonia and swine dysentery.

Chickens: Necrotic enteritis.

Recommended Dose

Pigs: 44g to 110g of Lincomycin (equivalent to 1kg to 2.5kg of product) per ton of feed for 5 to 10 days.

Chickens (excluding laying hens): 44g of Lincomycin (equivalent to 1kg of product) per ton of feed for 7 days.

Route of Administration

For oral administration via feed.

Contraindications

Lincosamides are considered contraindicated for use in rabbits, hamsters, guinea pigs, horses, and ruminants because of serious gastrointestinal effects that may occur, including death.

Lincomycin is contraindicated in animal with known hypersensitivity to it or having a preexisting monilial infection.

Lincomycin is generally not recommended for use in neonatal animals because of its effects on gut flora.

Warning and Precautions

Special warnings for each target species:

None known.

Special precautions for use

Special precautions for use in animals:

If any adverse reactions are observed, immediately consult a veterinarian.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- In case of accidental swallowing, see a doctor immediately.
- If this product accidentally gets into the eyes, nose, mouth, etc. of the user, immediately wash with water, gargle, etc. and consult a doctor.

- When mixing with feed, wear a mask and be careful not to inhale dust.
- When working, wear long-sleeved work clothes, work caps, protective glasses, masks, gloves, and other protective equipment, and be careful not to get this in your eyes, nose, mouth, etc.
- After using this product or if it adheres to the skin, wash it thoroughly with soap and gargle thoroughly with water.
- Keep out of reach of children.

Other Precautions:

- Use it according to the prescription and instructions of a veterinarian.
- This drug should be used only for the treatment of indications specified in the indications.
- Strictly adhere to the prescribed dosage and administration of this drug. In addition, even if the administration is within the period specified in the dosage and administration, do not repeat the administration.
- When using this drug, stop administration for the minimum period necessary for treatment.

Interaction with Other Medicaments

Do not mix this drug with other drugs.

Pregnancy and Lactation

Lincomycin crosses the placenta and cord blood concentrations are approximately 25% of those found in maternal serum. Safe use during pregnancy has not been established, but neither has the drug been implicated in causing teratogenic effects. Because lincomycin is distributed into milk, nursing animals of mothers given lincomycin may develop diarrhoea.

Side Effects

It has been reported that temporary loose stools were occasionally observed in pigs after administration of this drug.

Symptoms and Treatment of Overdose

In pigs treated with 2 to 10 times the recommended dose orally for 14 days altered the stool consistency, from loose stool to diarrhoea, without the loss of appetite. If suspected toxic reactions occur due to overdose, the medication should be discontinued and appropriate symptomatic treatment should be initiated if necessary.

When offered with the feed in amounts that were 3 to 5 times higher than the therapeutic ones for broilers in the course of 45 days, it did not cause mortality.

Storage Condition

Store below 30°C.

Shelf life

3 years. To be used within 24 hours after mixing into feed.

Withdrawal Periods

Pigs: 4 days

Chickens (excluding laying hens): 3 days

Packing

10kg & 25kg.

Revised date: 23.3.23

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com