



TRIAM-GUARD 0.15MG/ML TOPICAL SPRAY

Description and Composition

TRIAM-GUARD 0.15MG/ML TOPICAL SPRAY is a colourless clear solution. Each ml contains 0.15mg Triamcinolone Acetonide as its active ingredient and 5mg Phenoxyethanol as its preservative.

Pharmacodynamics

Triamcinolone acetonide is considered an intermediate acting glucocorticoid that is approximately 4 to 10 times more potent than hydrocortisone. Glucocorticoids have effects on virtually every cell type and system in mammals.

Pharmacokinetics

Triamcinolone is a potent anti-inflammatory corticosteroid. Upon absorption it distributes into the extravascular compartment with a monophasic half-life of approximately 35 minutes. In dogs, biliary metabolism and excretion is the primary means of elimination.

Indication

For the control of pruritus associated with allergic dermatitis in dogs.

Recommended Dose

Apply sufficient pump sprays to uniformly and thoroughly wet the affected areas while avoiding run-off of excess product. Avoid getting the spray in dog's eyes. The recommended treatment schedule is twice daily for seven days, then once daily for seven days, then every other day for an additional 14 days (28 days total).

Route of Administration

For external use only.

Contraindication

Systemic use of glucocorticoids is generally considered contraindicated in systemic fungal infections. Animals that have received glucocorticoids systemically, other than with "burst" therapy, should be tapered off the drugs. Should the animal undergo a "stressor" (e.g., surgery, trauma, illness, etc.) during the tapering process or until normal adrenal and pituitary function resume, additional glucocorticoids should be administered.

Warning and Precautions

Special warning for each target species

None known.

Special precautions for use

Special precautions for use in animals

Avoid getting the spray in dog's eyes.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wear gloves when applying the product. Spray in a well ventilated area. If the spray causes irritation to mucous membranes, discontinue use. Keep this and all drugs out of reach of children.

Interaction with Other Medicaments

The following drug interactions have either been reported or are theoretical in humans or animals receiving triamcinolone and may be of significance in veterinary patients: Amphotericin B, analgesics, opiate and/or anesthetics, local (epidural injections), anticholinesterase agents (e.g., pyridostigmine, neostigmine, etc.), aspirin, barbiturates, cyclophosphamide, cyclosporine, diuretics, potassium-depleting (e.g., spironolactone, triamterene), erythromycin, clarithromycin, estrogens, insulin, isoniazid, ketoconazole and other azole antifungals, mitotane, NSAIDS, phenobarbital, rifampin, vaccines, and warfarin.

Pregnancy and Lactation

Glucocorticoids are probably necessary for normal fetal development. They may be required for adequate surfactant production, myelin, retinal, pancreas and mammary development. Excessive dosages early in pregnancy may lead to teratogenic effects. Glucocorticoids unbound to plasma proteins will enter milk. High dosages or prolonged administration to mothers may potentially inhibit the growth of nursing newborns.

Side Effects

In dogs, polydipsia (PD), polyphagia (PP) and polyuria (PU), may all be seen with short-term "burst" therapy as well as with alternate-day maintenance therapy on days when giving the drug. Adverse effects in dogs can include dull, dry haircoat, weight gain, panting, vomiting, diarrhea, elevated liver enzymes, pancreatitis, GI ulceration, lipidemias, activation or worsening of diabetes mellitus, muscle wasting and behavioral changes (depression, lethargy, viciousness). Discontinuation of the drug may be necessary; changing to an alternate steroid may also alleviate the problem. Adverse effects associated with immunosuppressive doses are more common and potentially, more severe.

Symptoms and Treatment of Overdose

Glucocorticoids when given short-term are unlikely to cause harmful effects, even in massive dosages. One incidence of a dog developing acute CNS effects after accidental ingestion of glucocorticoids has been reported.

Storage Condition

Store below 30°C.

Shelf life

3 years. Discard 28 days after first opening.

Packing

50ml, 100ml, 200ml, 500ml.

Manufacturer & Product Registration Holder

Range Pharma Sdn Bhd

No. 1, Jalan TSB 11, Taman Industri Sg. Buloh,
47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia

Tel: 603-61568708 Fax: 603-61568707

Email: support@vetscareline.com

Revised date: 9.6.2026