

Product Name:

LPM Pain Relief Cream

Active ingredients:

Menthol 16%w/w, Methyl Salicylate 17%w/w

Preservative: Chlorocresol 0.2%w/w

Product Description:

White colour, smooth application, and characteristic odor cream.

Indication:

Pain relief for neck and shoulder aches, stressed, tired and stiff muscles.

Direction of use:

Gently massage a sufficient amount of the cream into the affected area 2 to 3 times a day.

Route of Administration: Topical

Pharmacodynamics*Menthol*

Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl Salicylate

Methyl salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins and cause platelet dysfunction. Methyl salicylate is used topically as a counter irritant. Upon application, it is absorbed through the skin and is applied for the relief of the pain in rheumatic conditions and painful muscle or joints.

Pharmacokinetics*Menthol*

After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl Salicylate

May be absorbed through intact skin. The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity. Absorption characteristics of salicylates vary with the dose, formulation and route of administration.

Percutaneous absorption is enhanced by exercise, heat, occlusion or disruption of the integrity of the skin or application to large areas of skin. Both the rate and extent of absorption increase after repeated application: increasing the bioavailability. Methyl salicylate is extensively metabolised to salicylic acid in the dermal and subcutaneous tissues after topical application. At therapeutic levels, the half-life of salicylates is 2-4 hours. As salicylate level increases into the toxic range, the half-life can be greater than 18 hours.

Contraindications:

Do not use in patients with aspirin or salicylate idiosyncrasy. Patient with known hypersensitivity to any of the ingredients in the product.

Warning and Precautions:

This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to larger skin area. Discontinue use if irritation develops. Do not apply to open wound or damaged skin. Avoid excessive use during pregnancy and lactation. Do not apply over large areas of the body or bandage or use with heating pads.

If symptoms persist, consult your doctor.

For external use only. Keep out of reach of children
Jauhkan daripada capaian kanak-kanak.

Effects on Ability to Drive and use Machines:
Not known

Interactions with Other Medicaments:
Warfarin.

Statement on Usage During Pregnancy and Lactation:
Skin irritation, contact dermatitis, rash, itching, redness or swelling, burning or stinging sensation may occur.

May cause hypersensitivity/ allergic reaction in some individuals with sensitive skin.

Symptoms and Treatment of Overdose:
Symptoms of methyl salicylate overdose
Salicylate intoxication can occur after ingestion or topical application of methyl salicylate. Mild chronic salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage. Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.

Symptoms of menthol overdose
Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness and coma; instant collapse in infants after the local application of menthol or camphor to their nostrils. Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self-limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema and status

epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure and death.

Treatment
The stomach should be emptied by gastric lavage or administration of oral activated charcoal. Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalemia and dehydration if present. Any convulsion must be controlled first through supportive care including anticonvulsant therapy.

Storage Conditions:
Store below 30°C.

Shelf Life:
Please refer to the packaging. Do not use beyond the expiry date.

Dosage Form: Cream

Packing: 30g, 50g

MAL XXXXXC

Product Registration Holder:
LPM BIO SCIENCE SBN BHD
Wisma SJC, No 13B, Persiaran KIP 1, Taman Perindustrian KIP, 52200 Kuala Lumpur.

Manufacturer:
SYARIKAT WEN KEN DRUG SDN BHD
24, Jalan Lambak, Taman Johor, 81200 Johor Bahru, Johor.

Revision Date: Oct 2023