

ENROXON 5MG/ML SOLUTION

Description and Composition

ENROXON 5MG/ML SOLUTION is a pale yellow clear solution. Each ml contains 5mg of Enrofloxacin as its active ingredient and 14mg Benzyl Alcohol as its preservative.

Pharmacodynamics

It is bactericidal in action with activity against Gram positive and Gram negative bacteria and mycoplasmas. The mechanism of action of the quinolones is unique among antimicrobials – they act primarily to inhibit bacterial DNA gyrase, an enzyme responsible for controlling the supercoiling of bacterial DNA during replication. Resealing of the double standard helix is inhibited resulting in irreversible degradation of the chromosomal DNA. The fluoroquinolones also possess activity against bacteria in the stationary phase by an alteration of the permeability of the outer membrane phospholipid cell wall.

Pharmacokinetics

The pharmacokinetics of enrofloxacin are such that oral and parenteral administration leads to similar serum levels. Enrofloxacin possesses a high distribution volume. Tissue levels 2-3 times higher than that found in the serum have been demonstrated in laboratory animals and target species. Organs in which high levels can be expected are the lungs, liver, kidney, bone and lymphatic system. Enrofloxacin also distributes into the cerebrospinal fluid, the aqueous humour and the foetus in pregnant animals.

Indication

For the treatment of diseases of the respiratory and alimentary tracts of bacterial or mycoplasmal origin (e.g. pasteurellosis, mycoplasmosis, coli-bacillosis, coli-septicaemia and salmonellosis), and multifactorial diseases such as atrophic rhinitis and enzootic pneumonia, where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice.

Recommended Dose

Dose rate is 1.5mg to 5mg Enrofloxacin per kg body weight daily.

For piglets up to 3kg body weight: 1ml once daily for 3 to 5 days.

For piglets 3kg up to 10kg body weight: 3ml once daily for 3 to 5 days.

1 pump stroke delivers 1ml of product.

Route of Administration

For oral administration.

Contraindications

Should not be used for prophylaxis.

Warning and Precautions

Special warnings for each target species:

None known.

Special precautions for use

Special precautions for use in animals:

Official and local antimicrobial policies should be taken into account when the product is used.

Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials.

Whenever possible, fluoroquinolones should only be used based on susceptibility testing.

Use of the product deviating from the instructions given may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to the potential for cross resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wear impervious gloves when handling the product.

Wash any splashes from skin or eyes immediately with water.

Wash hands and exposed skin after use.

Do not eat, drink or smoke while using the product.

Interaction with Other Medicaments

None known.

Pregnancy and Lactation

Not applicable – this product is only indicated for piglets.

Side Effects

None known.

Symptoms and Treatment of Overdose

Do not exceed the recommended dose. In accidental overdose there is no antidote and treatment should be symptomatic.

Storage Condition

Store below 30°C.

Shelf life

3 years.

After first opening the container, any product remaining after 24 hours should be discarded.

Withdrawal Period

Meat: 10 days

Packing

100ml, 200ml.

Revised date: 13.7.2023

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com