

TOFENAT 4% (w/v) Solution For Injection

For Animal Use Only

1. Name and Strength of Active Substance(s)

Tofenamic acid - 40mg/ml

2. Product Description:

Sterile injectable solution. Clear, very light yellow to yellowish liquid.

3. Pharmacodynamics:

Tolfenamic acid (N-(2-methyl-3-chlorophenyl) anthranilic acid) is a non-steroidal anti-inflammatory drug belonging to the fenamate group. Tolfenamic acid possesses anti-inflammatory, analgesic and antipyretic properties.

Its anti-inflammatory activity is due to the inhibition of cyclooxygenase, thus reducing the synthesis of prostaglandins and thromboxanes, important inflammatory mediators.

4. Pharmacokinetics:

In dogs, tolfenamic acid is readily absorbed by injectable administration. By injection, maximum plasma concentrations of about 4 µg/ml (subcutaneously) and about 3 µg/ml (intramuscularly) are obtained 2 hours after administration at 4 mg/kg.

In cats, absorption is quite rapid. By injection, a peak of 3.9 µg/ml is obtained within 1 hour of administration at 4 mg/kg.

In dogs and cats, over 99% of tolfenamic acid is bound to plasma proteins.

In the dog, only tolfenamic acid and its conjugate with glucuronic acid are found in urine.

The hydroxylated metabolites and their conjugates are mainly excreted by the kidneys. The unchanged tolfenamic acid and its glucuronides are predominantly excreted into the bile. Moreover, tolfenamic acid undergoes an intensive enterohepatic recycling.

Tolfenamic acid is distributed to all organs with high concentrations in plasma, digestive tract, liver, lungs and kidneys. The concentration in the brain however is low.

In dogs with renal insufficiency, the elimination of tolfenamic acid is unchanged and accumulation does not occur.

In cattle and pigs, tolfenamic acid injected intramuscularly at a dose of 2mg/kg is rapidly absorbed from the injection site with mean maximum plasma concentrations of about 1.4 µg/ml in cattle and 2.3 µg/ml in pigs obtained at about 1 hour.

The volume of distribution is about 1.3 l/kg.

Tolfenamic acid is extensively bound to plasma albumin (>97%).

Tolfenamic acid is distributed in all the organs with a high concentration in the plasma, digestive tract, liver, lungs and kidneys.

However, the concentration in the brain is low. Tolfenamic acid and its metabolites do not cross the placenta to any great extent.

Tolfenamic acid distribution involves extracellular fluids where concentrations similar to plasma are achieved both in healthy and inflamed peripheral tissues. It also appears in milk in the active form, mainly associated with the curds.

Tolfenamic acid undergoes extensive enterohepatic recirculation and, as a result prolonged concentrations are found in plasma.

The elimination half-life varies from 3 - 5 hours in pigs to 8 - 15 hours in cattle.

In cattle and pigs, tolfenamic acid is eliminated mainly unchanged in faeces (~30%) and urine (~70%).

5. Enviromental properities:

Not use in aquaculture.

6. Indications:

Cattle: Adjuvant in the treatment of pneumonia, improving general conditions, nasal secretion and as an adjuvant in the treatment of acute mastitis.

Porcine: Adjuvant in the treatment of Mastitis, Metritis and Agalactia syndrome.

Dogs: Treatment of inflammation associated with musculoskeletal disorders and reduction of postoperative pain.

Cats: Adjuvant treatment of upper respiratory tract diseases, in association with antimicrobial therapy, if necessary.

7. Recommended Dosage:

Cats and dogs: The recommended dose is 4mg/ kg bodyweight (1ml/ 10kg bodyweight) given as a single injection and repeated once after 24 to 48 hours if required and depending upon clinical assessment. Alternatively, a single injection of 1ml/ 10kg can be given with the treatment being continued by the oral route, using tablets.

In dogs administer by intramuscular or subcutaneous injection.

For the reduction of post-operative pain, this is best given pre-operatively, at the time of premedication one hour before induction of anaesthesia

In cats, administer by the subcutaneous route only.

Cattle: For inflammation associated with respiratory disease in cattle, the recommended dosage is 2mg/ kg (1ml/ 20kg bodyweight) by intramuscular injection into the neck area. Treatment may be repeated once after 48 hours.

For use in mastitis, the recommended dosage is 4 mg/kg bodyweight (1ml per 10kg bodyweight) as a single IV injection.

When administering intravenously, the product should be injected slowly. At the first signs of intolerance, the injection should be interrupted.

Porcine: the recommended dosage is 2mg/ kg (1ml/ 20kg bodyweight) as a single intramuscular injection.

The maximum injected volume is 20ml per injection site.

The stopper for the 20ml, 50ml and 100ml vial sizes may be safely punctured up to 20 times.

The stopper for the 250ml vial size may be safely punctured up to 50 times. The user should choose the most appropriate vial size according to the target species to be treated.

The use of an insulin-type needle/syringe is advisable particularly in low-weight animals to ensure an accurate dose.

8. Mode of Administration:

Cattle: intramuscular (IM) or intravenous (IV) route.

Porcine: intramuscular (IM) route.

Dogs: intramuscular or subcutaneous routes.

Cats: subcutaneous (SC) route.

9. Contraindication:

Do not use in cases of cardiac disease.

Do not use in cases of impaired hepatic function or acute renal insufficiency.

Do not use in cases of ulceration or digestive bleeding, in case of blood dyscrasia.

Do not inject intramuscularly in cats.

Do not use in known cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in dehydrated, hypovolaemic or hypotonic animals (due to its potential risk of increasing renal toxicity).

Do not administer other steroidal or non - steroidal anti - inflammatory drugs concurrently or within 24 hours of each other.

10. Warning and Precautions:

NSAIDS can cause inhibition of phagocytosis and hence in the treatment of inflammatory conditions associated with bacterial infections appropriate concurrent antimicrobial therapy should be instigated.

Special precautions for use in animals

Use in animals less than 6 weeks of age, or in aged animals, may involve additional risk. If such a use cannot be avoided, animals may require a reduced dosage and careful clinical management is essential. Reduced metabolism and excretion in these animals should be considered.

Concurrent administration of potential nephrotoxic drugs should be avoided.

It is preferable that the product is not administered to cats undergoing general anaesthesia until fully recovered.

Do not exceed the prescribed dosage or duration of treatment. The scale of pain relief after pre-operative administration may be influenced by the severity and duration of the operation.

Animals suffering from a chronic renal insufficiency and requiring an anti-inflammatory treatment may be treated with tolfenamic acid without requiring an adjustment of the dosage. However, the use of this product is contra-indicated in acute cases of renal insufficiency.

In case of undesirable effects (anorexia, vomiting, diarrhoea, presence of blood in faeces) occurring during the treatment, your veterinarian should be contacted for advice and the

possibility of stopping treatment should be considered.

Use aseptic precautions when administering the product.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

The product may cause skin sensitisation. People with known hypersensitivity to non-steroidal anti-inflammatory (NSAID) or to any of the excipients should avoid contact with the veterinary medicinal product.

Administer the veterinary medicinal product with caution to avoid accidental self-injection. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

This product may cause skin and eye irritation. Avoid contact with skin or eyes. In case of accidental contact, wash immediately exposed area with plenty of clean water.

Seek medical attention if irritation persists.

Wash hands after use.

11. Interactions with Other Medicaments:

Do not administer with other non-steroidal anti-inflammatory drugs simultaneously or with an interval of 24 hours between them. Other NSAIDs, diuretics, anticoagulants and substances with high affinity to plasma proteins may compete for binding and produce toxic effects.

Do not administer in conjunction with anticoagulants.

Avoid simultaneous administration of potentially nephrotoxic drugs.

Do not administer in conjunction with glucocorticoids.

12. Statement on usage during pregnancy and lactation:

Cats and dogs: Do not use during pregnancy.

Cattle and Porcine:

Pregnancy: Use only according to the benefit-risk assessment by the responsible veterinarian.

Lactation: The product can be used during lactation.

13. Adverse Effects/ Undesirable Effects

Collapse following rapid intravenous injection in cattle may occur uncommonly.

A temporary increase in thirst and/or diuresis may occur very rarely. In most of the cases, these signs cease spontaneously after treatment.

Diarrhoea and vomiting may occur during treatment very rarely. Where either persists, treatment should be discontinued.

Local injection site reactions may occur very rarely.

The frequency of adverse reactions is defined using the following convention:

- Very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- Common (more than 1 but less than 10 animals in 100 animals treated)
- Uncommon (more than 1 but less than 10 animals in 1000 animals treated)

- Rare (more than 1 but less than 10 animals in 10000 animals treated)
- Very rare (less than 1 animal in 10000 animals treated, including isolated reports).

14. Overdose and Treatment:

At high dosages, neurological disorders have been observed.

In case of overdose, administer symptomatic treatment.

15. Incompatibilities (for injections only): None known.

16. Withdrawal Period:

Bovine: Intramuscular injection: Meat: 12 days. Milk: zero hours.

Intravenous injection: Meat: 4 days. Milk: 24 hours.

Porcine: Meat: 16 days

17. Storage Condition:

Store in a dry place below 30°C. Keep away from direct sunlight.

18. Dosage Forms and packaging available: Solution - 100ml

19. Name & Address of Manufacturer:

SAI GON VET MANUFACTURING AND TRADING JOINT STOCK COMPANY
Anova Corporation Industrial zone, Long Cang commune, Tay Ninh province, Vietnam

20. Marketing Authorisation Holder:

Sunzen Corporation Sdn Bhd (470468-W)

11, Jalan Anggerik Mokara 31/47, Kota Kemuning, 40460 Shah Alam, Selangor, Malaysia.

21. Date of Revision of Package Insert: 22nd January 2026.

Changed the manufacturer name and update address from:
ANOVA PHARMA JOINT STOCK COMPANY
Anova Corporation Industrial Zone, Long Cang Commune, Can
Duoc District, Long An Province, Vietnam- Tel:(+84 272)3726262
.

to:

SAI GON VET MANUFACTURING AND TRADING JOINT
STOCK COMPANY
Anova Corporation Industrial zone, Long Cang commune, Tay
Ninh province, Vietnam- Tel:(+84 272)3726262.