

→

Pharmacode

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only.

PATASYN

(Olopatadine Hydrochloride Ophthalmic Solution USP 0.2% w/v)

QUALITATIVE AND QUANTITATIVE COMPOSITION

Active ingredient: 2.22 mg Olopatadine Hydrochloride USP equivalent to 2 mg Olopatadine

The full list of excipients is listed at the end of this leaflet.

PHARMACEUTICAL FORM

Ophthalmic drops

Clear colourless to pale yellow solution

PHARMACODYNAMICS

ATC Code: S01GX09

Pharmaco­therapeutic group: ophthalmologicals; decongestant and antiallergics; other antiallergics

Olopatadine is a potent selective antiallergic/antihistaminic agent that exerts its effects through multiple distinct mechanisms of actions. It antagonizes histamine (the primary mediator of allergic response in humans) and prevents histamine induced inflammatory cytokine production by human conjunctival epithelial cells. It may act on human conjunctival mast cells to inhibit the release of pro-inflammatory mediators.

In patients with patent nasolacrimal ducts, topical ocular administration of olopatadine Eye Drops, Solution was suggested to reduce the nasal signs and symptoms that frequently accompany seasonal allergic conjunctivitis. It does not produce a clinically significant change in pupil diameter.

PHARMACOKINETICS

Absorption

Olopatadine is absorbed systemically, as are other topically administered medicinal products. However, systemic absorption of topically applied olopatadine is minimal with plasma concentrations ranging from below the assay quantitation limit (<0.5 ng/ml) up to 1.3 ng/ml. These concentrations are 50-to 200-fold lower than those following well tolerated oral doses.

Biotransformation/ Metabolism

The major metabolites of olopatadine following oral administration in humans are N-desmethyl olopatadine (M1) and olopatadine N-oxide (M3). N-desmethyl olopatadine (M1) is almost exclusively demethylated by the cytochrome P-450 isozyme 3A4 (CYP3A4). Olopatadine was not an inhibitor of cytochrome P-450 isozymes and therefore drug-drug interactions due to metabolic interactions were not expected.

Elimination

In humans after topical ocular administration, N-desmethyl metabolite of olopatadine (M1) was not quantifiable (≤0.050 ng/mL) in plasma.

Warnings and Precautions

ocular tissues after topical ocular instillation.

Indication

PATASYN is indicated for the temporary prevention of itching of the eye due to allergic conjunctivitis.

Posology and Method of Administration

Adults

The recommended dose is one drop in each affected eye once a day.

Paediatric Use

Safety and effectiveness in paediatric patients below the age of 3 years have not been established.

Geriatric Use

No overall differences in safety and effectiveness have been observed between elderly and younger patients.

Method of Administration

- For topical ocular use only. Not for injection or oral use.
- To prevent contamination of the dropper tip and solution, care must be taken not to touch the eyelids, surrounding areas, or other surfaces with the dropper tip of the bottle.
- Keep the bottle tightly closed when not in use.
- In case of concomitant therapy with other topical ocular medicines, an interval of 5 minutes should be allowed between successive applications. Eye ointments should be administered last.
- Patients should be advised not to wear a contact lens if their eye is red.
- PATASYN should not be used to treat contact lens related irritation.

Contraindications

Hypersensitivity to the active substance or to any of the excipients.

Warnings and Precautions

PATASYN is an antiallergic/antihistaminic agent and, although administered topically, is absorbed systemically. If signs of serious reactions or hypersensitivity occur, discontinue the use of this treatment.

PATASYN contains benzalkonium chloride, a preservative that has been reported to cause punctate keratopathy and/or toxic ulcerative keratopathy. Close monitoring is required with frequent or prolonged use in dry eye patients, or in conditions where the cornea is compromised.

PATASYN contains benzalkonium chloride which may cause eye irritation and is known to discolour soft contact lenses. Avoid contact with soft contact lenses. Patients must be instructed to remove contact lenses prior to application of PATASYN and wait at least 15 minutes before reinsertion.

Interactions with other Medicaments

No clinically relevant interactions have been described.

Pregnancy and Lactation

Fertility

Studies have not been performed to evaluate the effect of administration of olopatadine on human fertility. Effects in non-clinical fertility studies in male and female animals were observed only at dosages considered sufficiently in excess of the maximum human exposure indicating little relevance to clinical use.

No effects on human fertility are anticipated since systemic exposure to olopatadine is negligible by the topical ocular route (AUC₀₋₆ of 9.7 ng*hr/mL in humans administered 1 drop of 0.77% olopatadine in both eyes once daily for 6.5 days).

Olopatadine can be used by women of childbearing potential.

Pregnancy

There are no or limited amount of data from the use of olopatadine in pregnant women. Effects in non-clinical reproduction and developmental toxicity studies were observed only at exposures considered sufficiently in excess of the maximum human exposure indicating little relevance to clinical use.

No effects during pregnancy are anticipated since systemic exposure to olopatadine is negligible by the topical ocular route (AUC₀₋₆ of 9.7 ng*hr/mL in humans

240 mm

190 mm

USV Private Limited 		
Market/Country : Malaysia	Co-ordinator Name : Sachin	Software : Corel Draw
SAP Code : 3017XXX	Item type : Packinsert	Date: 30/08/2023
Product Name : Patasyn		Version : 2022
Actual Size : 190 mm x 240 mm (L x H) (Font & Back)		Artist Name : Gajanan
Reference SAP Code :-----		
Pharmacode: -----	Font: Swis721 Cn BT	Size: 8 pt
CMYK/Pantone :  Black		
Reason : New Development.		
Path : M:\Govandi\Artwork\Venaa - Final Artworks in CDR\Current 08.2010\Formulation\Export\Malaysia		
NOTE TO THE PRINTER: Processor / Printer shall not make any correction / changes / deviation in the approved artwork. If any discrepancy observed in the artwork, Printer / Processor shall return the artwork to USV.		

Prepared by RA-Artwork	Checked by RA-Artwork	Checked by CQA	Checked by Legal	Checked by Other Dept. (If required)	Approved by RA-Artwork

administered 1 drop of 0.77% olopatadine in both eyes once daily for 6.5 days).
Before prescribing olopatadine to a pregnant woman, a physician should weigh the benefit of administration to the woman to the risk of the foetus.

Breastfeeding

Available pharmacodynamic/toxicological data in animals have shown excretion of olopatadine/metabolites in milk following high dose systemic olopatadine administration. Radioactivity has been identified in the milk of nursing rats at concentrations of 0.33 to 4.28 times that of plasma concentrations (1,184 ng*hr/mL AUC₀₋₂₄ in lactating rat plasma) following a 1 mg/kg oral administration of ¹⁴C-olopatadine.

Based upon the low level of olopatadine present in human plasma following topical ocular administration (AUC₀₋₆ of 9.7 ng*hr/mL in humans administered 1 drop of 0.77% olopatadine in both eyes once daily for 6.5 days), the concentration of olopatadine potentially present in breast milk is expected to be negligible. However, as there is no data available on the concentration of olopatadine/metabolites in human milk following topical ocular administration, a risk to the suckling child cannot be excluded.

Patients should be informed that antihistamines may affect the milk production of a nursing mother. Before prescribing olopatadine to a nursing mother, a physician should weigh the benefit of administration to the mother to the risk of the breastfeeding child.

SIDE EFFECTS

No serious ophthalmic or systemic adverse reactions related to olopatadine were reported. The most frequent treatment-related adverse reaction was eye pain.

Tabulated list of adverse reactions

System Organ Classification	Frequency	Adverse Reactions
Infections and infestations	Uncommon	Rhinitis
Immune system disorders	Not known	Hypersensitivity, swelling face
Nervous system disorders	Common	Headache, dysgeusia
	Uncommon	Dizziness, hypoaesthesia
	Not known	Somnolence
Eye disorders	Common	Eye pain, eye irritation, dry eye, abnormal sensation in eyes
	Uncommon	Corneal erosion, corneal epithelium defect, corneal epithelium disorder, punctate keratitis, keratitis, corneal staining, eye discharge, photophobia, vision blurred, visual acuity reduced, blepharospasm, ocular discomfort, eye pruritus, conjunctival follicles, conjunctival disorder, foreign body sensation in eyes, lacrimation increased, erythema of eyelid, eyelid oedema, eyelid disorder, ocular hyperaemia
	Not known	Corneal oedema, eye oedema, eye swelling, conjunctivitis, mydriasis, visual disturbance, eyelid margin crusting
Respiratory, thoracic and mediastinal disorders	Common	Nasal dryness
	Not known	Dyspnoea, sinusitis
Gastrointestinal disorders	Not known	Nausea, vomiting
Skin and subcutaneous tissue disorders	Uncommon	Dermatitis contact, skin burning sensation, dry skin
	Not known	Dermatitis, erythema
General disorders and administration site conditions	Common	Fatigue
	Not known	Asthenia, malaise

Cases of corneal calcification have been reported very rarely in association with the use of phosphate containing eyedrops in some patients with significantly damaged corneas.

SYMPTOMS AND TREATMENT OF OVERDOSE

No data are available in humans regarding overdose by accidental or deliberate ingestion. In the case of overdose, appropriate monitoring and management of the patient should be implemented.

EFFECTS ON ABILITY TO DRIVE OR USE MACHINE

Olopatadine is a non-sedating anti-histamine. Temporary blurred vision after drop use, or other visual disturbances may affect the ability to drive or use machines. If blurred vision occurs after instillation, the patient must wait until the vision clears before driving or using machinery.

LIST OF EXCIPIENTS

Benzalkonium chloride
Edetate disodium
Povidone K 30
Dibasic sodium phosphate anhydrous
Sodium chloride
Hydrochloric acid
Sodium hydroxide
Water for injection
Nitrogen

STORAGE CONDITION

Store below 30°C. Discard 28 days after opening.

SHELF LIFE

24 months.

PACKAGING AVAILABLE

2.5 mL of solution in white, round LDPE bottle attached with a transparent LDPE nozzle. The bottle is closed with a white HDPE tamper evident screwcap. One bottle per outer box with leaflet.

LEAFLET REVISION DATE

30/08/2023

Product Registration Holder

Synercam (M) Sdn. Bhd.
A-17-7, Block A, Jaya One,
No. 72A, Jalan Profesor Diraja
Ungku Aziz, Seksyen 13, 46200,
Petaling Jaya, Malaysia

Mfd. by:

USV Private Limited

H-13,16,16A,17,18,19,20,21, & E-22, O IDC,
Mahatma Gandhi Udyog Nagar, Dabhel,
Daman - 396210, India.



3017XX

240 mm

190 mm