

59.4mm

420mm

Immune system disorders				anaphylactic reaction, hypersensitivity	
Endocrine disorders			hyperprolactinaemia ^a	inappropriate antidiuretic hormone secretion ^c , glucose urine present	
Metabolism and nutrition disorders		weight increased, increased appetite, weight decreased, decreased appetite	diabetes mellitus ^d , hyperglycaemia, waist circumference increased, anorexia, blood triglycerides increased	water intoxication, diabetic ketoacidosis ^c , hypoglycaemia, polydipsia, blood cholesterol increased	hyperinsulinaemia
Psychiatric disorders	insomnia ^e	mania, agitation, depression, anxiety	sleep disorder, confusional state, libido decreased, anorgasmia, nervousness, nightmare	catatonia, somnambulism, blunted affect ^c	
Nervous system disorders	parkinsonism ^b , akathisia ^b , sedation/somnolence, headache	dystonia ^b , dizziness, dyskinesia ^b , tremor	tardive dyskinesia, convulsion ^o , syncope, psychomotor hyperactivity, dizziness postural, disturbance in attention, dysarthria, dysgeusia, hypoaesthesia, paresthaesia	neuroleptic malignant syndrome, cerebral ischaemia, unresponsive to stimuli ^c , loss of consciousness, depressed level of consciousness ^c , diabetic coma ^c , balance disorder, coordination abnormal, head titubation ^c	
Eye disorders		vision blurred	photophobia, conjunctivitis, dry eye	glaucoma, eye movement disorder ^c , eye rolling ^c , lacrimation increased, ocular hyperaemia	
Ear and labyrinth disorders			vertigo, tinnitus, ear pain		
Cardiac disorders		atrioventricular block, conduction disorder, electrocardiogram QT prolonged, bradycardia, tachycardia	sinus arrhythmia, electrocardiogram abnormal, palpitations	atrial fibrillation, postural orthostatic tachycardia syndrome ^c	
Vascular disorders		orthostatic hypotension, hypertension	hypotension	pulmonary embolism, venous thrombosis, ischaemia, flushing	
Respiratory, thoracic and mediastinal disorders		pharyngolaryngeal pain, cough, nasal congestion	dyspnoea, wheezing, epistaxis	sleep apnoea* syndrome, hyperventilation, pneumonia aspiration, respiratory tract congestion dysphonia	pulmonary congestion
Gastrointestinal disorders		abdominal pain, abdominal discomfort, vomiting, nausea, constipation, diarrhoea, dyspepsia, dry mouth, toothache	swollen tongue, gastroenteritis, dysphagia, flatulence	pancreatitis ^c , intestinal obstruction, ileus, faecal incontinence, faecaloma ^c , cheilitis	
Hepatobiliary disorders		transaminases increased	gamma-glutamyltransaminase increased, hepatic enzyme increased	jaundice	
Skin and subcutaneous tissue disorders		pruritus, rash	urticaria, alopecia, eczema, acne	angioedema, drug eruption ^c , hyperkeratosis, dry skin, erythema, skin discolouration, seborrheic dermatitis, dandruff	
Musculoskeletal and connective tissue disorders		musculoskeletal pain, back pain, arthralgia	blood creatine phosphokinase increased, muscle spasms, joint stiffness, joint swelling, muscular weakness, neck pain	rhabdomyolysis ^c , posture abnormal ^c	
Renal and urinary disorders			urinary incontinence, pollakiuria, urinary retention, dysuria		
Pregnancy, puerperium and perinatal conditions				drug withdrawal syndrome neonatal (see section Pregnancy and Lactation) ^c	
Reproductive system and breast disorders		amenorrhoea	erectile dysfunction, ejaculation disorder, menstrual disorder ^o , galactorrhoea, sexual dysfunction, breast pain, breast discomfort	priapism ^c , menstruation delayed ^c , gynaecomastia, breast engorgement, breast enlargement ^c , breast discharge, vaginal discharge	
General disorders		pyrexia, asthenia,	face oedema,	hypothermia ^c , body	
disorders		fatigue	oedema ^o , chills, body temperature	temperature decreased ^c , drug withdrawal syndrome ^c , induration ^c	
Injury, poisoning and procedural complications			fall		

^a Refer to 'Hyperprolactinaemia' below.
^b Refer to 'Extrapyramidal symptoms' below.
^c Not observed in Paliperidone Extended Release Tablet clinical studies but observed in post-marketing environment with paliperidone.
^d In placebo-controlled pivotal trials, diabetes mellitus was reported in 0.05% in Paliperidone Extended Release Tablet-treated subjects compared to a rate of 0% in placebo group.
Overall incidence from all clinical trials was 0.14% in all Paliperidone Extended Release Tablet-treated subjects.
^e **Insomnia includes:** initial insomnia, middle insomnia; **Convulsion includes:** grand mal convulsion; **Oedema includes:** generalised oedema, oedema peripheral, pitting oedema; **Menstrual disorder includes:** menstruation irregular, oligomenorrhoea.
^a Atypical antipsychotic drugs, such as Paliperidone have been associated with cases of sleep apnoea, with or without concomitant weight gain. In patients who have a history of or are at risk for sleep apnoea, HEALOL Paliperidone Extended Release Tablet should be prescribed with caution.

Undesirable effects noted with risperidone formulations
Paliperidone is the active metabolite of risperidone, therefore, the adverse reaction profiles of these compounds (including both the oral and injectable formulations) are relevant to one another. In addition to the above adverse reactions, the following adverse reactions have been noted with the use of risperidone products and can be expected to occur with HEALOL Paliperidone Extended Release Tablet.
Psychiatric disorders: sleep-related eating disorder
Nervous system disorders: cerebrovascular disorder, Restless legs syndrome
Eye disorders: floppy iris syndrome (intraoperative)
Respiratory, thoracic and mediastinal disorders: rales
Skin and subcutaneous tissue disorders: Stevens-Johnson syndrome/toxic epidermal necrolysis

Description of selected adverse reactions
Extrapyramidal symptoms (EPS)
In schizophrenia clinical trials, there was no difference observed between placebo and the 3 and 6 mg doses of HEALOL Paliperidone Extended Release Tablet. Dose dependence for EPS was seen with the two higher doses of HEALOL Paliperidone Extended Release Tablet (9 mg). In the schizoaffective disorder studies, the incidence of EPS was observed at a higher rate than placebo in all dose groups without a clear relationship to dose.

EPS included a pooled analysis of the following terms: Parkinsonism (includes salivary hypersecretion, musculoskeletal stiffness, parkinsonism, drooling, cogwheel rigidity, bradykinesia, hypokinesia, masked facies, muscle tightness, akinesia, nuchal rigidity, muscle rigidity, parkinsonian gait, and glabellar reflex abnormal, parkinsonian rest tremor), akathisia (includes akathisia, restlessness, hyperkinesia, and restless leg syndrome), dyskinesia (dyskinesia, muscle twitching, choreoathetosis, athetosis, and myoclonus), dystonia (includes dystonia, hypertonia, torticollis, muscle contractions involuntary, muscle contracture, blepharospasm, oculogyration, tongue paralysis, facial spasm, laryngospasm, myotonia,

opisthotonus, oropharyngeal spasm, pleurothotonus, tongue spasm, and trismus), and tremor. It should be noted that a broader spectrum of symptoms are included that do not necessarily have an extrapyramidal origin.

Weight gain
In schizophrenia clinical trials, the proportions of subjects meeting a weight gain criterion of $\geq 7\%$ of body weight were compared, revealing a similar incidence of weight gain for HEALOL Paliperidone Extended Release Tablet 3 mg and 6 mg compared with placebo, and a higher incidence of weight gain for HEALOL Paliperidone Extended Release Tablet 9 mg and 12 mg compared with placebo.

In schizoaffective disorder clinical trials, a higher percentage of HEALOL Paliperidone Extended Release Tablet-treated subjects (5%) had an increase in body weight of $\geq 7\%$ compared with placebo-treated subjects (1%). In the study that examined two dose groups (see section Pharmacodynamics), the increase in body weight of $\geq 7\%$ was 3% in the lower-dose (3-6 mg) group, 7% in the higher-dose (9-12 mg) group, and 1% in the placebo group.
Hyperprolactinaemia
In schizophrenia clinical trials, increases in serum prolactin were observed with HEALOL Paliperidone Extended Release Tablet in 67% of subjects. Adverse reactions that may suggest increase in prolactin levels (e.g., amenorrhoea, galactorrhoea, menstrual disturbances, gynaecomastia) were reported overall in 2% of subjects. Maximum mean increases of serum prolactin concentrations were generally observed on day 15 of treatment, but remained above baseline levels at study endpoint.

Class effects
QT prolongation, ventricular arrhythmias (ventricular fibrillation, ventricular tachycardia), sudden unexplained death, cardiac arrest and Torsade de pointes may occur with antipsychotics. Cases of venous thromboembolism, including cases of pulmonary embolism and cases of deep vein thrombosis have been reported with antipsychotic drugs - Frequency unknown.

Paliperidone is the active metabolite of risperidone. The safety profile of risperidone may be pertinent.

Elderly
In a study conducted in elderly subjects with schizophrenia, the safety profile was similar to that seen in non-elderly subjects. HEALOL Paliperidone Extended Release Tablet has not been studied in elderly patients with dementia. In clinical trials with some other atypical antipsychotics, increased risks of death and cerebrovascular accidents have been reported (see section Warnings & Precautions).

Paediatric population
Summary of the safety profile
In one short-term and two longer-term studies with HEALOL Paliperidone Extended Release Tablet conducted in adolescents 12 years and older with schizophrenia, the overall safety profile was similar to that seen in adults. In the pooled adolescent schizophrenia population (12 years and older, N = 545) exposed to HEALOL Paliperidone Extended Release Tablet, the frequency and type of undesirable effects were similar to those in adults except for the following ADRs that were reported more frequently in adolescents receiving Paliperidone Extended Release Tablet than adults receiving HEALOL Paliperidone Extended Release Tablet (and more frequently than placebo): sedation/somnolence, parkinsonism, weight increase, upper respiratory tract infection, akathisia, and tremor were reported very commonly ($\geq 1/10$) in adolescents; abdominal pain, galactorrhoea, gynaecomastia, acne, dysarthria, gastroenteritis, epistaxis, ear infection, blood triglyceride increased, and vertigo were reported commonly ($\geq 1/100$, $< 1/10$) in adolescents.

Extrapyramidal Symptoms (EPS)
In the short-term, placebo-controlled, fixed-dose adolescent study, the incidence of EPS was higher than placebo for all doses of HEALOL Paliperidone Extended Release Tablet with an increased frequency of EPS at higher doses. Across all adolescent studies, EPS was more common in adolescents than in adults for each HEALOL Paliperidone Extended Release Tablet dose.

Weight gain
In the short-term, placebo-controlled, fixed-dose adolescent study, a higher percentage of HEALOL Paliperidone Extended Release Tablet-treated subjects (6-19% depending on dose) had an increase in body weight of $\geq 7\%$ compared to placebo-treated subjects (2%). There was no clear dose relationship. In the long-term 2-year study, the subjects who were exposed to HEALOL Paliperidone Extended Release Tablet during both the double-blind and open-label studies reported a modest weight gain (4.9 kg).

In adolescents, weight gain should be assessed against that expected with normal growth.

Prolactin
In the up to 2-year, open-label treatment study of HEALOL Paliperidone Extended Release Tablet in adolescents with schizophrenia, incidence of elevated serum prolactin levels occurred in 48% of females and 60% of males. Adverse reactions that may suggest increase in prolactin levels (e.g., amenorrhoea, galactorrhoea, menstrual disturbances, gynaecomastia) were reported overall in 9.3% of subjects.

Reporting of suspected adverse reactions
Reporting suspected adverse reactions after authorisation of the medical product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Postmarketing Data
Eye Disorders
Frequency: Not known – Floppy iris syndrome (intraoperative)
The NPCB would like to alert healthcare professionals on the risk of intraoperative floppy iris syndrome (IFIS) during and after cataract surgery in patients treated with risperidone or paliperidone-containing products. Patients must be counselled to inform their surgeon about current or prior use of these products before the surgery. In patients with such medication history, the cataract surgery should be approached with caution. If IFIS is suspected, modifications to surgical technique may be required.

Not known: frequency cannot be estimated from the available data
Lung congestion
Increased insulin (a hormone that controls blood sugar levels) in your blood.

The following side effects have been seen with the use of another medicine called risperidone that is very similar to paliperidone, so these can also be expected with HEALOL Paliperidone Extended Release Tablet: sleep-related eating disorder, other types of blood vessel problems in the brain, crackly lung sounds, and severe or life-threatening rash with blisters and peeling skin that may start in and around the mouth, nose, eyes and genitals and spread to other areas of the body (Stevens-Johnson syndrome/toxic epidermal necrolysis). Eye problems during cataract surgery may also occur. During cataract surgery, a condition called intraoperative floppy iris syndrome (IFIS) can happen if you take or have taken HEALOL Paliperidone Extended Release Tablet. If you need to have cataract surgery, be sure to tell your eye doctor if you take or have taken this medicine.

Symptoms and Treatment of Overdose
In general, expected signs and symptoms are those resulting from an exaggeration of paliperidone's known pharmacological effects, i.e., drowsiness and sedation, tachycardia and hypotension, QT prolongation, and extrapyramidal symptoms. Torsade de pointes and ventricular fibrillation have been reported in association with overdose. In the case of acute overdosage, the possibility of multiple medicinal product involvement should be considered.

Consideration should be given to the prolonged-release nature of the product when assessing treatment needs and recovery. There is no specific antidote to paliperidone. General supportive measures should be employed. Establish and maintain a clear airway and ensure adequate oxygenation and ventilation. Cardiovascular monitoring should commence immediately and should include continuous electrocardiographic monitoring for possible arrhythmias. Hypotension and circulatory collapse should be treated with appropriate measures such as intravenous fluid and/or sympathomimetic agents.

Administration of activated charcoal together with a laxative should be considered. In case of severe extrapyramidal symptoms, anticholinergic agents should be administered. Close supervision and monitoring should continue until the patient recovers.

Effects on Ability to Drive and Use Machine
Paliperidone can have minor or moderate influence on the ability to drive and use machines due to potential nervous system and visual effects (see section Side Effects). Therefore, patients should be advised not to drive or operate machines until their individual susceptibility to HEALOL Paliperidone Extended Release Tablet is known.

Preclinical Safety Data
Not applicable.

List of Excipients
Inactive ingredients of HEALOL Paliperidone Extended Release Tablet are Polyethylene oxide WSR N-80, Sodium chloride, Povidone K-30, Magnesium stearate, Polyethylene oxide WSR-303, Iron Oxide (Ferric oxidier red), Butylated hydroxytoluene, Opadry CA. The 9 mg tablets also contain color overcoating material Opadry pink. The 6 mg tablets also contain color overcoating material Opadry orange. The 3 mg tablets also contain color overcoating material Opadry white.

Instructions for Use
NA

Storage Condition
Do not store above 30 °C
Store in the original package in order to protect from moisture.

Shelf Life
2 years
See expiry date on the outer pack.

Product Registration Holder
HEALOL PHARMACEUTICALS SDN. BHD.
NO. 74-3, JALAN WANGSA DELIMA 6 KLSC WANGSA MAJU 53300 KUALA LUMPUR WILAYAH PERSEKUTUAN KUALA LUMPUR MALAYSIA

Manufacturer
AUPA Biopharm Co., Ltd.
No.1, Kwang-Fu Rd., Huko, Hsinchu County, Taiwan (R.O.C.), TAIWAN

Therapeutic Code
Pharmacologic group: Psycholeptics, other antipsychotics, ATC code: N05AX13

Date of Revision: 17 January 2024