

65 mm

UNIGEM

Gemcitabine 38 mg/mL concentrate for solution for Infusion

1. Name of the medicinal product

Gemcitabine 38 mg/ml Concentrate for Solution for Infusion

2. Qualitative and quantitative composition

One ml of Gemcitabine Concentrate for Solution for Infusion contains gemcitabine hydrochloride, equivalent to 38 mg gemcitabine.

The quantitative composition of each presentation is provided in the table below:

Presentation	Strength	Quantity of gemcitabine (as hydrochloride)	Volume of Solution
1 g/26.3 ml	38 mg/ml	1 g	26.3 ml

Excipients with known effect: Each ml of concentrate contains up to 0.76mg (0.03 mmol) sodium. For a full list of excipients, see section 6.1.

3. Pharmaceutical form

Gemcitabine 38 mg /mL concentrate for solution for infusion is clear and colorless to light straw colored solution and packed in 30 mL/20 mm flint moulded USP type-I glass vial, stoppered with 20 mm grey bromobutyl omniflex plus coated rubber stopper (ready for sterilization) and sealed with 20 mm flip-off red colour seal.

After dilution, a clear and colorless solution.

4. Clinical particulars

4.1 Therapeutic indications

- Gemcitabine is indicated for the treatment of locally advanced or metastatic non-small cell-lung cancer.
- Gemcitabine is indicated for the treatment of adult patients with locally advanced or metastatic adenocarcinoma of the pancreas. Gemcitabine is indicated for patients with 5-FU refractory pancreatic cancer.
- Gemcitabine is indicated for the treatment of patients suffering from bladder cancer, at the invasive stage.
- Gemcitabine, in combination with paclitaxel, is indicated for the treatment of patients with unresectable, locally recurrent or metastatic breast cancer who have relapsed following adjuvant/neoadjuvant chemotherapy. Prior chemotherapy should have included an anthracycline unless clinically contraindicated.
- Gemcitabine, in combination with carboplatin, is indicated for the treatment of patients with recurrent epithelial ovarian carcinoma, who have relapse > 6 months, following platinum-base therapy.

4.2 Posology and method of administration

Dosage

Adults:

Non-small cell lung cancer

Single agent use

The recommended dose is 1000 mg/m², given by intravenous infusion.

The administration must be repeated once weekly for three weeks, followed by a one-week rest period. This four-week cycle is then repeated. A dose reduction or delay before each administration of the chemotherapy may be applied, based upon the amount of toxicity experienced by the patient.

Combination use:

Gemcitabine in combination with cisplatin can be administered using two dosage regimens one regimen use a three-week schedule, the other uses a four-week schedule.

The three-week schedule is the usual regimen; this three-week cycle uses gemcitabine 1250 mg/m² given by 30 minutes intravenous infusion on days 1 and 8, followed by one-week rest period. This three-week cycle is then repeated. A dosage reduction or delay before each administration of the chemotherapy may be applied, based upon the amount of toxicity experienced by the patient.

The four-week cycle uses Gemcitabine 1000 mg/m² given by 30 minutes intravenous infusion on days 1, 8 and 15, followed by one-week rest period. This four-week cycle is then repeated. A dosage reduction or delay before each administration of the chemotherapy may be applied, based upon the amount of toxicity experienced by the patient.

Pancreatic adenocarcinoma.

The recommended dose is 1000 mg/m², given by 30 minutes intravenous infusion. This should be repeated once weekly for 7 weeks, followed by a week rest Subsequent cycles should consist of injections once weekly for 3 consecutive weeks out of every 4 weeks. A dosage reduction or delay before each administration of the chemotherapy may be applied based upon the amount of toxicity experienced by the patient.

Bladder cancer, at the invasive stage

The recommended dose of Gemcitabine in combination with cisplatin, is 1000 mg/m², given by 30 minutes intravenous infusion on days 1, 8 and 15, followed by one-week rest period for a 28-day cycle. Cisplatin is given at a recommended dose of 70 mg/m² on day 2. This four-week cycle is then repeated. A dosage reduction or delay before each administration of the chemotherapy may be applied based upon the amount of toxicity experienced by the patient. A clinical trial showed more myelosuppression when cisplatin was used in doses of 100 mg/m².

Breast cancer

Paclitaxel (175 mg/m²) administered on Day 1 over approximately 3 hours as an intravenous infusion, followed by Gemcitabine (1250 mg/m²) as a 30-minute intravenous infusion on Days 1 and 8 of each 21-day cycle. Dose reduction with each cycle or within a cycle may be applied based upon the amount of toxicity experienced by the patient.

Ovarian Cancer

Gemcitabine in combination with carboplatin is recommended using Gemcitabine 1000 mg/m² administered on Days 1 and 8 of each 21-day cycle as a 30-minute intravenous infusion. After Gemcitabine, carboplatin should be given on Day 1 to attain a target AUC of 4.0 mg/mL/min. Dosage reduction with each cycle or within a cycle may be applied based upon the amount of toxicity experienced by the patient.

For all indications

Patients receiving Gemcitabine should be monitored prior to each dose for platelet, leukocyte, and granulocyte counts and, if necessary, the dose of Gemcitabine may be either reduced or withheld in the presence of hematologic toxicity, according to the following scale:

Absolute granulocyte count (x 10 ⁹ /l)		Platelet count (x 10 ⁹ /l)	% of total dose
> 1000	And	> 100,000	100
500 - 1,000	or	50,000 - 100,000	75
<500	or	< 50,000	0

Periodic physical examination and checks of renal and hepatic function should be made to detect nonhematologic toxicity. Dosage reduction with each cycle or within a cycle may be applied based upon the amount of toxicity experienced by the patient.

Doses should be withheld until toxicity has resolved in the opinion of the physician.

Elderly patients

Gemcitabine has been well tolerated by patients over 65 years of age. The pharmacokinetic data suggest that the metabolism of the drug is not affected by age.

Patients with hepatic or renal impairment

Gemcitabine should be used with caution in patients with hepatic insufficiency or with impaired renal function as there is insufficient information from clinical studies to allow clear dose recommendation for this patient population.

Mild to moderate renal insufficiency (GFR from 30mL/min to 80mL/min) has no consistent, significant effect on Gemcitabine pharmacokinetics.

Children

Gemcitabine has been studied in limited Phase I and II trials in children in a variety of tumor types. These studies did not provide sufficient data to establish the efficacy and safety of Gemcitabine in children.

Method of Administration

Intravenous route.

Gemcitabine is well tolerated during infusion and is usually easy to administer.

If extravasation, the administration must be stopped immediately.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

Breast-feeding: It is not known whether Gemcitabine is excreted in human milk and adverse effects on the suckling child cannot be excluded. Breast-feeding must be discontinued during Gemcitabine therapy.

4.4 Special warnings and precautions for use

Prolongation of the infusion time and increased dosing frequency have been shown to increase toxicity.

Haematological toxicity

Gemcitabine can suppress bone marrow function as manifested by leucopenia, thrombocytopenia and anaemia.

Patients receiving gemcitabine should be monitored prior to each dose for platelet, leucocyte and granulocyte counts. Suspension or modification of therapy should be considered when drug-induced bone marrow depression is detected.

However, myelosuppression is short lived and usually does not result in dose reduction and rarely in discontinuation.

Peripheral blood counts may continue to deteriorate after gemcitabine administration has been stopped. In patients with impaired bone marrow function, the treatment should be started with caution. As with other cytotoxic treatments, the risk of cumulative bone-marrow suppression must be considered when gemcitabine treatment is given together with other chemotherapy.

Hepatic and renal impairment

Gemcitabine should be used with caution in patients with hepatic impairment or with impaired renal function as there is insufficient information from clinical studies to allow clear dose recommendation for this patient population.

Administration of gemcitabine in patients with concurrent liver metastases or a pre-existing medical history of hepatitis, alcoholism or liver cirrhosis may lead to exacerbation of the underlying hepatic impairment. Laboratory evaluation of renal and hepatic function (including virological tests) should be performed periodically.

Concomitant radiotherapy

Concomitant radiotherapy (given together or ≤7 days apart): Toxicity has been reported.

Live vaccinations

Yellow fever vaccine and other live attenuated vaccines are not recommended in patients treated with gemcitabine.

Posterior reversible encephalopathy syndrome

Reports of posterior reversible encephalopathy syndrome (PRES) with potentially severe consequences have been reported in patients receiving gemcitabine as single agent or in combination with other chemotherapeutic agents. Acute hypertension and seizure activity were reported in most gemcitabine patients experiencing PRES, but other symptoms such as headache, lethargy, confusion and blindness could also be present. Diagnosis is optimally confirmed by magnetic resonance imaging (MRI). PRES was typically reversible with appropriate supportive measures. Gemcitabine should be permanently discontinued and supportive measures implemented, including blood pressure control and anti-seizure therapy, if PRES develops during therapy.

Cardiovascular

Due to the risk of cardiac and/or vascular disorders with gemcitabine, particular caution must be exercised with patients presenting a history of cardiovascular events.

Capillary leak syndrome

Capillary leak syndrome has been reported in patients receiving gemcitabine as single agent or in combination with other chemotherapeutic agents. The condition is usually treatable if recognised early and managed appropriately, but fatal cases have been reported. The

condition involves systemic capillary hyperpermeability during which fluid and proteins from the intravascular space leak into the interstitium. The clinical features include generalised oedema, weight gain, hypoalbuminaemia, severe hypotension, acute renal impairment and pulmonary oedema. Gemcitabine should be discontinued and supportive measures implemented if capillary leak syndrome develops during therapy. Capillary leak syndrome can occur in later cycles and has been associated in the literature with adult respiratory distress syndrome.

Pulmonary

Pulmonary effects, sometimes severe (such as pulmonary oedema, interstitial pneumonitis or adult respiratory distress syndrome (ARDS)) have been reported in association with gemcitabine therapy. The aetiology of these effects is unknown. If such effects develop, consideration should be made to discontinuing gemcitabine therapy. Early use of supportive care measure may help ameliorate the condition.

Renal

Haemolytic uraemic syndrome

Clinical findings consistent with the haemolytic uraemic syndrome (HUS) were rarely reported in patients receiving gemcitabine. HUS is a potentially life-threatening disorder. Gemcitabine should be discontinued at the first signs of any evidence of microangiopathic haemolytic anaemia, such as rapidly falling haemoglobin with concomitant thrombocytopenia, elevation of serum bilirubin, serum creatinine, blood urea nitrogen, or LDH. Renal failure may not be reversible with discontinuation of therapy and dialysis may be required.

Fertility

In fertility studies gemcitabine caused hypospermatogenesis in male mice. Therefore, men being treated with gemcitabine are advised not to father a child during and up to 6 months after treatment and to seek further advice regarding cryoconservation of sperm prior to treatment because of the possibility of infertility due to therapy with gemcitabine.

Excipient information

Gemcitabine 1 g Concentrate for Solution for Infusion.

This medicinal product contains less than 1 mmol sodium (23 mg) per vial, that is to say essentially “sodium-free”.

4.5 Interaction with other medicaments

No specific interaction studies have been performed.

Radiotherapy

Concurrent (given together or ≤7 days apart) - Toxicity associated with this multimodality therapy is dependent on many different factors, including dose of gemcitabine, frequency of gemcitabine administration, dose of radiation, radiotherapy planning technique, the target tissue, and target volume. Pre-clinical and clinical studies have shown that gemcitabine has radiosensitising activity. In a single trial, where gemcitabine at a dose of 1,000 mg/m² was administered concurrently for up to 6 consecutive weeks with therapeutic thoracic radiation to patients with non-small cell lung cancer, significant toxicity in the form of severe, and potentially life threatening mucositis, especially oesophagitis, and pneumonitis was observed, particularly in patients receiving large volumes of radiotherapy [median treatment volumes 4,795 cm3]. Studies done subsequently have suggested that it is feasible to administer gemcitabine at lower doses with concurrent radiotherapy with predictable toxicity, such as a phase II study in non-small cell lung cancer, where thoracic radiation doses of 66 Gy were applied concomitantly with an administration with gemcitabine (600 mg/m², four times) and cisplatin (80 mg/m² twice) during 6 weeks. The optimum regimen for safe administration of gemcitabine with therapeutic doses of radiation has not yet been determined in all tumour types.

Non-concurrent (given >7 days apart) - Analysis of the data does not indicate any enhanced toxicity when gemcitabine is administered more than 7 days before or after radiation, other than radiation recall. Data suggest that gemcitabine can be started after the acute effects of radiation have resolved or at least one week after radiation.

Radiation injury has been reported on targeted tissues (e.g. oesophagitis, colitis, and pneumonitis) in association with both concurrent and non-concurrent use of gemcitabine.

Others

Yellow fever and other live attenuated vaccines are not recommended due to the risk of systemic, possibly fatal, disease, particularly in immunosuppressed patients.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no adequate data from the use of gemcitabine in pregnant women. Studies in animals have shown reproductive toxicity. Based on results from animal studies and the mechanism of action of gemcitabine, this substance should not be used during pregnancy unless clearly necessary. Women should be advised not to become pregnant during treatment with gemcitabine and to warn their attending physician immediately, should this occur after all.

Breast-feeding

It is not known whether gemcitabine is excreted in human milk and adverse effects on the suckling child cannot be excluded. Breast-feeding must be discontinued during gemcitabine therapy.

Fertility

In fertility studies gemcitabine caused hypospermatogenesis in male mice. Therefore, men being treated with gemcitabine are advised not to father a child during and up to 6 months after treatment and to seek further advice regarding cryoconservation of sperm prior to treatment because of the possibility of infertility due to therapy with gemcitabine.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. However, gemcitabine has been reported to cause mild to moderate somnolence, especially in combination with alcohol consumption. Patients should be cautioned against driving or operating machinery until it is established that they do not become somnolent.

4.8 Undesirable effects

The most commonly reported adverse drug reactions associated with Gemcitabine treatment include: nausea with or without vomiting, raised liver transaminases (AST/ALT) and alkaline phosphatase, reported in approximately 60% of patients; proteinuria and haematuria reported in approximately 50% patients; dyspnoea reported in 10-40% of patients (highest incidence

in lung cancer patients); allergic skin rashes occur in approximately 25% of patients and are associated with itching in 10% of patients.

The frequency and severity of the adverse reactions are affected by the dose, infusion rate and intervals between doses. Dose-limiting adverse reactions are reductions in thrombocyte, leucocyte and granulocyte counts.

Clinical trial data

Frequencies are defined as: Very common (≥ 1/10), Common (≥ 1/100 to <1/10), Uncommon (≥ 1/1000 to <1/100), Rare (≥ 1/10,000 to <1/1000), Very Rare (<1/10,000), Not known (cannot be estimated from the available data).

The following table of undesirable effects and frequencies is based on data from clinical trials. Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

SYSTEM ORGAN CLASS	FREQUENCY GROUPING
Infections and infestations	Common • Infections Not known • Sepsis
Blood and lymphatic system disorders	Very common • Leucopenia (Neutropoenia Grade 3 = 19.3 %; Grade 4 =6%) Bone-marrow suppression is usually mild to moderate and mostly affects the granulocyte count. • Thrombocytopenia • Anaemia Common • Febrile neutropoenia Very rare • Thrombocytosis • Thrombotic microangiopathy
Immune system disorders	Very Rare • Anaphylactoid reaction
Metabolism and nutrition disorders	Common • Anorexia
Nervous system disorders	Common • Headache • Insomnia • Somnolence Uncommon • Cerebrovascular accident Very rare • Posterior reversible encephalopathy syndrome.
Cardiac disorders	Uncommon • Arrhythmias, predominantly supraventricular in nature • Heart failure Rare • Myocardial infarct
Vascular disorders	Rare • Clinical signs of peripheral vasculitis and gangrene • Hypotension Very rare • Capillary leak syndrome.
Respiratory, thoracic and mediastinal disorders	Very common • Dyspnoea –usually mild and passes rapidly without treatment Common • Cough • Rhinitis Uncommon • Interstitial pneumonitis. • Bronchospasm – usually mild and transient but may require parenteral treatment Rare • Pulmonary oedema • Adult respiratory distress syndrome. Not known • Pulmonary eosinophilia
Gastrointestinal disorders	Very common • Vomiting • Nausea Common • Diarrhoea • Stomatitis and ulceration of the mouth • Constipation Very rare • Ischaemic colitis
Hepatobiliary disorders	Very common • Elevation of liver transaminases (AST and ALT) and alkaline phosphatase Common • Increased bilirubin Uncommon • Serious hepatotoxicity, including liver failure and death Rare • Increased gamma-glutamyl transferase (GGT)

Size: 515(L) x 330(W) mm
Folding Size: 65(L) x 34(W) mm
Font: Arial Narrow
Min. Font Size: 9pt
Market: Unimed(Malaysia)
Color: Black
R00 (02.07.2024)

Skin and subcutaneous tissue disorders	Very common - Allergic skin rash frequently associated with pruritus Common - Alopecia - Itching - Sweating Rare - Severe skin reactions, including desquamation and bullous skineruptions - Ulceration - Vesicle and sore formation - Scaling Very rare - Toxic epidermal necrolysis - Stevens-Johnson Syndrome Not known - Pseudoocellulitis
	Common - Back pain - BMyalgia
Musculoskeletal and connective tissuedisorders	Common - Haematuria - Mild proteinuria Uncommon - Renal failure - Haemolytic uraemic syndrome
Renal and urinary disorders	Very Common - Haematuria - Mild proteinuria Uncommon - Renal failure - Haemolytic uraemic syndrome
General disorders and administration site conditions	Very common - Influenza-like symptoms - the most common symptoms are fever,headache, chills, myalgia, asthenia and anorexia. Cough, rhinitis,malaise, perspiration and sleeping difficulties have also been reported. - Oedema/peripheral oedema, including facial oedema. Oedema is usually reversible after stopping treatment Common - Fever - Asthenia - Chills Rare - Injection site reactions - mainly mild in nature
Injury, poisoning, and proceduralcomplications	Rare - Radiation toxicity - Radiation recall

Combination use in breast cancer

The frequency of grade 3 and 4 haematological toxicities, particularly neutropaenia, increases when gemcitabine is usedin combination with paclitaxel. However, the increase in these adverse reactions is not associated with an increasedincidence of infections or haemorrhagic events. Fati- gue and febrile neutropaenia occur more frequently whengemcitabine is used in combination with paclitaxel. Fatigue, which is not associated with anaemia, usually resolves afterthe first cycle.

Grade 3 and 4 Adverse Events Paclitaxel versus gemcitabine plus paclitaxel				
	Number (%) of Patients			
	Paclitaxel arm (N=259)		Gemcitabine plus Paclitaxel arm (N=262)	
	Grade 3	Grade 4	Grade 3	Grade 4
Laboratory				
Anaemia	5 (1.9)	1 (0.4)	15 (5.7)	3 (1.1)
Thrombocytopaenia	0	0	14 (5.3)	1 (0.4)
Neutropaenia	11 (4.2)	17 (6.6)*	82 (31.3)	45 (17.2)*
Non-laboratory				
Febrile neutropaenia	3 (1.2)	0	12 (4.6)	1(0.4)
Fatigue	3 (1.2)	1 (0.4)	15 (5.7)	2 (0.8)
Diarrhoea	5 (1.9)	0	8 (3.1)	0
Motor neuropathy	2 (0.8)	0	6 (2.3)	1 (0.4)
Sensory neuropathy	9 (3.5)	0	14 (5.3)	1 (0.4)

*Grade 4 neutropaenia lasting for more than 7 days occurred in 12.6% of patients in the combination arm and 5.0% of patients in the paclitaxel arm.

Combination use in bladder cancer

Grade 3 and 4 Adverse Events MVAC versus Gemcitabine plus cisplatin				
	Number (%) of Patients			
	MVAC* arm (N=196)		Gemcitabine plus cisplatin arm (N=200)	
	Grade 3	Grade 4	Grade 3	Grade 4
Laboratory				
Anaemia	30 (16)	4 (2)	47 (24)	7 (4)
Thrombocytopaenia	15 (8)	25 (13)	57 (29)	57 (29)
Non-laboratory				

Nausea and vomiting	37 (19)	3 (2)	44 (22)	0 (0)
Diarrhoea	15 (8)	1 (1)	6 (3)	0 (0)
Infection	19 (10)	10 (5)	4 (2)	1 (1)
Stomatitis	34 (18)	8 (4)	2 (1)	0 (0)

*Methotrexate, Vinblastine, Doxorubicin and Cisplatin

Combination use in ovarian cancer

Grade 3 and 4 Adverse Events Carboplatin versus Gemcitabine plus carboplatin				
	Number (%) of Patients			
	Carboplatin arm (N=174)		Gemcitabine plus carboplatin arm (N=175)	
	Grade 3	Grade 4	Grade 3	Grade 4
Laboratory				
Anaemia	10(5.7)	4 (2.3)	39 (22.3)	9 (5.1)
Neutropaenia	19(10.9)	2(1.1)	73(41.7)	50 (28.6)
Thrombocytopaenia	18(10.3)	2(1.1)	53(30.3)	8 (4.6)
Leucopaenia	11(6.3)	1(0.6)	84(48.0)	9 (5.1)
Non-laboratory				
Haemorrhage	0(0.0)	0(0.0)	3(1.8)	0(0.0)
Febrile neutropaenia	0(0.0)	0(0.0)	2(1.1)	0(0.0)
Infection without neutropaenia	0(0.0)	0(0.0)	0(0.0)	1(0.6)

Sensory neuropathy was also more frequent in the combination arm than with single agent Carboplatin

4.9 Overdose

There is no known antidote for overdose of gemcitabine. Doses as high as 5700 mg/m² have been administered by intravenous infusion over 30-minutes every 2 weeks with clinically acceptable toxicity. In the event of suspected overdose, the patient should be monitored with appropriate blood counts and receive supportive therapy, as necessary.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents/pyrimidine analogues.

ATC code: L01BC05

Cytotoxic activity in cell cultures

Gemcitabine shows significant cytotoxic effects against a variety of cultured murine and human tumour cells. Its action is phase-specific such that gemcitabine primarily kills cells that are undergoing DNA synthesis (S-phase) and, under certain circumstances, blocks the progression of cells at the junction of the G1/S phase boundary. In vitro, the cytotoxic effect of gemcitabine is dependent on both concentration and time.

Antitumoral activity in preclinical models.

In animal tumour models, antitumoural activity of gemcitabine is schedule-dependent.

When gemcitabine is administered daily, high mortality among the animals but minimal antitumoural activity is observed. If, however, gemcitabine is given every third or fourth day, it can be administered in nonlethal doses with substantial antitumoural activity against a broad spectrum of mouse tumours.

Mechanism of action

Cellular metabolism and mechanism of action: Gemcitabine (dFdC), which is a pyrimidine antimetabolite, is metabolised intracellularly by nucleoside kinase to the active diphosphate (dFdCDP) and triphosphate (dFdCTP) nucleosides. The cytotoxic effect of gemcitabine is due to inhibition of DNA synthesis by two mechanisms of action by dFdCDP and dFdCTP. First, dFdCDP inhibits ribonucleotide reductase, which is uniquely responsible for catalysing the reactions that produce deoxynucleoside triphosphates (dCTP) for DNA synthesis. Inhibition of this enzyme by dFdCDP reduces the concentration of deoxynucleosides in general and, in particular, dCTP. Second, dFdCTP competes with dCTP for incorporation into DNA (self-potentialion).

Likewise, a small amount of gemcitabine may also be incorporated into RNA. Thus, the reduced intracellular concentration of dCTP potentiates the incorporation of dFdCTP into DNA. DNA polymerase epsilon lacks the ability to eliminate gemcitabine and to repair the growing DNA strands. After gemcitabine is incorporated into DNA, one additional nucleotide is added to the growing DNA strands. After this addition there is essentially a complete inhibition in further DNA synthesis (masked chain termination). After incorporation into DNA, gemcitabine appears to induce the programmed cell death process known as apoptosis.

Clinical data

Bladder cancer

A randomised phase III study of 405 patients with advanced or metastatic urothelial transitional cell carcinoma showed no difference between the two treatment arms, gemcitabine/cisplatin versus methotrexate/vinblastine/adriamycin/cisplatin (MVAC), in terms of median survival (12.8 and 14.8 months respectively, p=0.547), time to disease progression (7.4 and 7.6 months respectively, p=0.842) and response rate (49.4% and 45.7% respectively, p=0.512). However, the combination of gemcitabine and cisplatin had a better toxicity profile than MVAC.

Pancreatic cancer

In a randomised phase III study of 126 patients with advanced or metastatic pancreatic cancer, gemcitabine showed a statistically significant higher clinical benefit response rate than 5-fluorouracil (23.8% and 4.8% respectively, p=0.0022). Also, a statistically significant prolongation of the time to progression from 0.9 to 2.3 months (log-rank p<0.0002) and a statistically significant prolongation of median survival from 4.4 to 5.7 months (log-rank p<0.0024) was observed in patients treated with gemcitabine compared to patients treated with 5-fluorouracil.

Non small cell lung cancer

In a randomised phase III study of 522 patients with inoperable, locally advanced or metastatic NSCLC, gemcitabine in combination with cisplatin showed a statistically significant higher

response rate than cisplatin alone (31.0% and 12.0%, respectively, p<0.0001). A statistically significant prolongation of the time to progression, from 3.7 to 5.6 months (log-rank p<0.0012) and a statistically significant prolongation of median survival from 7.6 months to 9.1 months (log-rank p<0.004) was observed in patients treated with gemcitabine/cisplatin compared to patients treated with cisplatin.

In another randomised phase III study of 135 patients with stage IIIB or IV NSCLC, a combination of gemcitabine and cisplatin showed a statistically significant higher response rate than a combination of cisplatin and etoposide (40.6% and 21.2%, respectively, p=0.025). A statistically significant prolongation of the time to progression, from 4.3 to 6.9 months (p=0.014) was observed in patients treated with gemcitabine/cisplatin compared to patients treated with etoposide/cisplatin.

In both studies it was found that tolerability was similar in the two treatment arms.

Ovarian carcinoma

In a randomised phase III study, 356 patients with advanced epithelial ovarian carcinoma who had relapsed at least 6 months after completing platinum based therapy were randomised to therapy with gemcitabine and carboplatin (Gcb), or carboplatin (Cb). A statistically significant prolongation of the time to progression of disease, from 5.8 to 8.6 months (log-rank p= 0.0038) was observed in the patients treated with GCb compared to patients treated with Cb. Differences in response rate of 47.2% in the GCb arm versus 30.9% in the Cb arm (p=0.0016) and median survival 18 months (GCb) versus 17.3 (Cb) (p=0.73) favoured the GCb arm.

Breast cancer

In a randomised phase III study of 529 patients with inoperable, locally recurrent or metastatic breast cancer with relapse after adjuvant/neoadjuvant chemotherapy, gemcitabine in combination with paclitaxel showed a statistically significant prolongation of time to documented disease progression from 3.98 to 6.14 months (log-rank p=0.0002) in patients treated with gemcitabine/paclitaxel compared to patients treated with paclitaxel.

After 377 deaths, the overall survival was 18.6 months versus 15.8 months (log rank p=0.0489, HR 0.82) in patients treated with gemcitabine/paclitaxel compared to patients treated with paclitaxel and the overall response rate was 41.4% and 26.2% respectively (p= 0.0002).

5.2 Pharmacokinetic properties

The pharmacokinetics of gemcitabine have been examined in 353 patients in seven studies. The 121 women and 232 men ranged in age from 29 to 79 years. Of these patients, approximately 45% had non-small cell lung cancer and 35% were diagnosed with pancreatic cancer. The following pharmacokinetic parameters were obtained for doses ranging from 500 to 2,592 mg/m² that were infused from 0.4 to 1.2 hours.

Absorption

Peak plasma concentrations (obtained within 5 minutes of the end of the infusion) were 3.2 to 45.5 µg/ml. Plasma concentrations of the parent compound following a dose of 1,000 mg/ m2/30-minutes are greater than 5 µg/ml for approximately 30-minutes after the end of the infusion, and greater than 0.4 µg/ml for an additional hour.

Distribution

The volume of distribution of the central compartment was 12.4 l/m² for women and 17.5 l/m² for men (inter-individual variability was 91.9%). The volume of distribution of the peripheral compartment was 47.4 l/m². The volume of the peripheral compartment was not sensitive to gender. The plasma protein binding was considered to be negligible.

Half-life: This ranged from 42 to 94 minutes depending on age and gender. For the recommended dosing schedule, gemcitabine elimination should be virtually complete within 5 to 11 hours of the start of the infusion. Gemcitabine does not accumulate when administered once weekly.

Biotransformation

Gemcitabine is rapidly metabolised by cytidine deaminase in the liver, kidney, blood and other tissues. Intracellular metabolism of gemcitabine produces the gemcitabine mono, di and triphosphates (dFdCMP, dFdCDP and dFdCTP) of which dFdCDP and dFdCTP are considered active. These intracellular metabolites have not been detected in plasma or urine. The primary metabolite, 2'-deoxy-2', 2'-difluorouridine (dFdU), is not active and is found in plasma and urine.

Elimination

Systemic clearance ranged from 29.2 l/hr/m² to 92.2 l/hr/m² depending on gender and age (inter-individual variability was 52.2%). Clearance for women is approximately 25% lower than the values for men. Although rapid, clearance for both men and women appears to decrease with age. For the recommended gemcitabine dose of 1000 mg/m² given as a 30- minute infusion, lower clearance values for women and men should not necessitate a decrease in the gemcitabine dose. Urinary excretion: Less than 10% is excreted as unchanged drug.

Renal clearance was 2 to 7 l/hr/m².

During the week following administration, 92 to 98% of the dose of gemcitabine administered is recovered, 99% in the urine, mainly in the form of dFdU and 1% of the dose is excreted in faeces.

dFdCTP kinetics

This metabolite can be found in peripheral blood mononuclear cells and the information below refers to these cells. Intracellular concentrations increase in proportion to gemcitabine doses of 35-350 mg/m2/30-minutes, which give steady state concentrations of 0.4-5 µg/ ml. At gemcitabine plasma concentrations above 5 µg/ml, dFdCTP levels do not increase, suggesting that the formation is saturable in these cells.

Half-life of terminal elimination: 0.7-12 hours.

dFdU kinetics

Peak plasma concentrations (3-15 minutes after end of 30-minute infusion, 1000 mg/m²):

28-52 µg/ml. Trough concentration following once weekly dosing: 0.07-1.12 µg/ml, with no apparent accumulation. Triphasic plasma concentration versus time curve, mean half-life of terminal phase 65 hours (range 33-84 hr).

Formation of dFdU from parent compound: 91%-98%.

Mean volume of distribution of central compartment: 18 l/m² (range 11-22 l/m²).

Mean steady state volume of distribution (Vss): 150 l/m² (range 96-228 l/m²).

Tissue distribution: Extensive.

Mean apparent clearance: 2.5 l/hr/m² (range 1-4 l/hr/m²).

Urinary excretion: All.

Gemcitabine and paclitaxel combination therapy

Combination therapy did not alter the pharmacokinetics of either gemcitabine or paclitaxel.

Gemcitabine and carboplatin combination therapy

When given in combination with carboplatin the pharmacokinetics of gemcitabine were not altered.

Renal impairment

Mild to moderate renal insufficiency (GFR from 30 ml/min to 80 ml/min) has no consistent, significant effect on gemcitabine pharmacokinetics.

6. Pharmaceutical particulars

6.1 List of excipients

Sodium hydroxide (for pH adjustment)

Hydrochloric acid (for pH adjustment)

Water for Injections

6.2 Incompatibilities

This medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

Unopened vial

24 months

In-Use: Further dilution

After dilution, chemical and physical in-use stability has been demonstrated for:

Diluent	Target Concentration	Storage Conditions	Time period
0.9% sodium chloride solution for infusion	0.1 mg/ml and 26 mg/ml	2-8°C in the absence of light in non-PVC (polyolefin)infusion bags	24 hours
0.9% sodium chloride solution for infusion	0.1 mg/ml and 26 mg/ml	2-8°C in the absence of light in PVC infusion bags	24 hours
0.9% sodium chloride solution for infusion	0.1 mg/ml and 26 mg/ml	25°C under normal lighting conditions in PVC infusion bags	6 hours
5% glucose solu- tion for infusion	0.1 mg/ml and 26 mg/ml	25°C under normal lighting conditions in PVC infusion bags	6 hours

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless dilution has taken place in controlled and validated aseptic conditions.

6.4 Special precautions for storage

Store in a refrigerator (2°C-8°C).

For storage conditions after dilution of the medicinal product.

6.5 Nature and contents of container

1 g/26.3 ml presentation

Each carton contains 1 vial (30 ml/20 mm flint moulded USP Type I glass vial with reinforced Non-PVC Base, with 20 mm grey bromobutyl omniflex plus coated rubber stoppers (RFS) with 20 mm aluminium flip off red colour seals).

Each vial of the 1g presentation contains 26.3 ml concentrate.

6.6 Special precautions for disposal and other handling

Handling

The normal safety precautions for cytostatic agents must be observed when preparing and disposing of the infusion solution. Handling of the concentrate should be done in a safety box and protective coats and gloves should be used. If no safety box is available, the equipment should be supplemented with a mask and protective glasses.

If the preparation comes into contact with the eyes, this may cause serious irritation. The eyes should be rinsed immediately and thoroughly with water. If there is lasting irritation, a doctor should be consulted. If the solution is spilled on the skin, rinse thoroughly with water.

Instructions for dilution

An approved diluent for gemcitabine solution is sodium chloride 9 mg/ml (0.9%) solution for injection (without preservative).

- Use the aseptic technique during any dilution of gemcitabine for intravenous infusion administration.
- Parenteral medicinal products should be inspected visually for particulate matter and discolouration prior to administration. If particulate matter is observed, do not administer.
- Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. Marketing authorisation holder

Manufactured by:

Shilpa Medicare Limited

Unit IV, Plot No's S-20 to S-26, Pharma SEZ, TSIC, Green Industrial Park, Polepally (V),

Jadcherla (M), Mahabubnagar (Dt), 509301, Telangana, India.

Name & Address of Product Registration Holder (PRH):

Unimed Sdn Bhd

53, Jalan Tembaga SD5/2B, Bandar Sri Damansara,52200 Kuala Lumpur, Malaysia.

8. Registration No: MAL24036035ACZ

9. Date of Revision

07.2024

45xxxxx