





Therefore, the recommended dose of docetaxel in those patients with elevated liver function test (LFTs) is 75 mg/m<sup>2</sup> and LFTs should be measured at baseline and before each cycle.

For patients with serum bilirubin levels <1.5xULN and ALT and AST >3.5 times the ULN concurrent with serum alkaline phosphatase levels >6 times the ULN, no dose-reduction can be recommended and docetaxel should not be used unless clinically indicated.

In combination with cisplatin and 5-fluorouracil for the treatment of patients with gastric adenocarcinoma, the pivotal clinical study excluded patients with ALT and/or AST >1.5 times ULN associated with alkaline phosphatase >2.5 times ULN and bilirubin >1 times ULN; for these patients, no dose-reductions can be recommended and docetaxel should not be used unless clinically indicated. No data are available in patients with hepatic impairment treated by docetaxel in combination in the other indications.

#### Patients with renal impairment

There are no data available in patients with severely impaired renal function treated with docetaxel.

Docetaxel 10 mg/ml solution for infusion contains Macrogol 300 which may increase the risk of nephrotoxicity in patients with renal impairment.

#### Neutrocytopenia

The development of severe peripheral neutrocytosis requires a reduction of dose.

#### Cardiac toxicity

Heart failure has been observed in patients receiving docetaxel in combination with trastuzumab, particularly following anthracycline (doxorubicin or epirubicin)-containing chemotherapy. This may be moderate to severe and has been associated with death.

When patients are candidates for treatment with docetaxel in combination with trastuzumab, they should undergo baseline cardiac assessment. Cardiac function should be further monitored during treatment (e.g. every three months) to help identify patients who may develop cardiac dysfunction. For more details see summary of product characteristics of trastuzumab.

Ventricular arrhythmia including ventricular tachycardia (sometimes fatal) has been reported in patients treated with docetaxel in combination regimens including doxorubicin, 5-fluorouracil and/or cyclophosphamide.

Baseline cardiac assessment is recommended.

#### Eye disorders

Cystoid macular oedema (CMO) has been reported in patients treated with docetaxel. Patients with impaired vision should undergo a prompt and complete ophthalmological examination. In case CMO is diagnosed, docetaxel treatment should be discontinued and appropriate treatment initiated.

#### Second primary malignancies

Second primary malignancies have been reported when docetaxel was given in combination with anticancer treatments known to be associated with second primary malignancies. Second primary malignancies (including acute myeloid leukaemia, myelodysplastic syndrome and non-Hodgkin lymphoma) may occur several months or years after docetaxel-containing therapy. Patients should be monitored for second primary malignancies.

#### Tumour lysis syndrome

Tumour lysis syndrome has been reported with docetaxel after the first or the second Patient at risk of tumour lysis syndrome (e.g. with renal impairment, hyperuricaemia, bulky tumour, rapid progression) should be closely monitored. Correction of dehydration and treatment of high uric acid levels are recommended prior to initiation of treatment.

#### Other

Contraceptive measures must be taken by both men and women during treatment and for men at least 6 months after cessation of therapy.

The concomitant use of docetaxel with strong CYP3A4 inhibitors (e.g. ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, neflavinir, ritonavir, saquinavir, telitremvir and voriconazole) should be avoided.

Additional cautions for use in **adjunct treatment of breast cancer**

#### Complicated neutropenia

For patients who experience complicated neutropenia (prolonged neutropenia, febrile neutropenia or infection), G-CSF and dose reduction should be considered.

#### Gastrointestinal reactions

Symptoms such as early abdominal pain and tenderness, fever, diarrhoea, with or without neutropenia, may be early manifestations of serious gastrointestinal toxicity and should be evaluated and treated promptly.

#### Concurrent heart failure (CHF)

Patients should be monitored for symptoms of congestive heart failure during therapy and during the follow-up period. In patients treated with the TAC regimen for node positive breast cancer, the risk of CHF has been shown to be higher during the first year after treatment.

#### Patients with 4+ nodes

As the benefit observed in patient with 4+ nodes was not statistically significant on disease-free survival (DFS) and overall survival (OS), the positive benefit risk ratio for TAC in patients with 4+ nodes was not fully demonstrated in the final analysis.

#### Elderly

##### Caution for use in **adjunct treatment of breast cancer**

There are limited data available in patients >70 years of age or docetaxel use in combination with doxorubicin and cyclophosphamide.

##### Caution for use in **castration-resistant prostate cancer**

Of the 333 patients treated with docetaxel every three weeks in a prostate cancer study (TAX 327), 209 patients were 65 years of age or greater and 68 patients were older than 75 years. In patients treated with docetaxel every three weeks, the incidence of related nail disorders occurred at a rate >10% higher in patients who were 65 years of age or greater compared to younger patients. The incidence of related fever, diarrhoea, anaemia and peripheral oedema occurred at rates >10% higher in patients who were 75 years of age or greater versus less than 65 years.

##### Caution for use in **hormone-sensitive prostate cancer**

Of the 545 patients treated with docetaxel every 3 weeks in a hormone-sensitive prostate cancer study (STAMPEDE), 296 patients were 65 years of age or older, and 48 patients were 75 years of age or older. More patients aged ≥65 years in the docetaxel arm reported hypersensitivity reaction, neutropenia, anaemia, fluid retention, dyspnoea, and nail changes when compared to the patients aged less than 65 years. None of those increases in frequency reached 10% difference with the control arm. In patients who were 75 years of age or older, when compared to younger patients, neutropenia, anaemia, diarrhoea, dyspnoea and upper respiratory tract infection were reported with a greater incidence (at least 10% higher).

##### Caution for use in **gastric adenocarcinoma cancer**

Among the 300 (221 patients in the phase III part of the study and 79 patients in the phase II part) patients treated with docetaxel in combination with cisplatin and 5-fluorouracil in the gastric cancer study, 74 were 65 years of age or older and 4 patients were 75 years of age or older. The incidence of serious adverse events was higher in older people compared to younger patients. The incidence of the following adverse events (all grades): lethargy, stomatitis, neutropenic infection occurred at rates >10% higher in patients who were 65 years of age or older compared to younger patients.

Older people treated with TCF should be closely monitored.

#### Exciipient information

20mg/2 ml vial

Each 2 ml vial of concentrate contains 364 mg anhydrous ethanol which is equivalent to less than 10 ml beer or 4 ml wine.

80 mg/8 ml vial

Each 8 ml vial of concentrate contains 1455 mg anhydrous ethanol which is equivalent to less than 37 ml beer or 15 ml wine.

Each 16 ml vial of concentrate contains 2911 mg anhydrous ethanol which is equivalent to less than 73 ml beer or 30 ml wine.

An example of ethanol exposure based on maximum single daily dose is as follows:

- Administration of 18 l of docetaxel 10 mg/ml concentrate for solution for infusion to an adult weighing 70 kg would result in exposure to 46.5 mg/kg of ethanol which may cause a rise in blood alcohol concentration (BAC) of 7.8 mg/100 ml.

For comparison, for an adult drinking a glass of wine or 500 ml of beer, the BAC is likely to be about 50 mg/100 ml.

Co-administration with medicines containing e.g. propylene glycol or ethanol may lead to accumulation of ethanol and induce adverse effects, particularly in young children with low or immature metabolic capacity.

The amount of ethanol in this medicine is not likely to have an effect in adults and adolescents, and its effects in children are not likely to be noticeable. It may have some effects, such as somnolence, in neonates and young children.

The ethanol content in this medicinal product should be carefully considered in the following patient groups who may be at higher risk of ethanol-related adverse effects:

- Pregnant or breast-feeding women
- Patients with liver disease
- Patients with epilepsy
- Patients suffering from alcoholism.

Consideration should be given to possible effects on the central nervous system.

#### J. Interactions with Other Medications

The amount of docetaxel in this medicinal product is not may alter the effects of other medicinal products.

The in vitro studies have shown that the metabolism of docetaxel may be modified by the concomitant administration of compounds which induce, inhibit or are metabolised by (and thus may inhibit the enzyme competitively) cytochrome P450-3A4 such as ciclosporin, ketoconazole and erythromycin. As a result, caution should be exercised when treating patients with these medicinal products as concomitant therapy since there is a potential for a significant interaction.

In case of combination with CYP3A4 inhibitors, the occurrence of docetaxel adverse reactions may increase, as a result of reduced metabolism. If the concomitant use of a strong CYP3A4 inhibitor (e.g. ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, neflavinir, ritonavir, saquinavir, telitremvir and voriconazole) cannot be avoided, a close clinical surveillance is warranted and a dose-adjustment of docetaxel may be suitable during the treatment with the strong CYP3A4 inhibitor In a pharmacokinetic study with 7 patients, the co-administration of docetaxel with the strong CYP3A4 inhibitor ketoconazole led to a significant decrease in docetaxel clearance by 49%.

Docetaxel pharmacokinetics in the presence of prednisone was studied in patients with metastatic prostate cancer. Docetaxel is metabolised by CYP3A4 and prednisone is known to induce CYP3A4. No statistically significant differences in prednisone or the pharmacokinetics of docetaxel was observed. Docetaxel is a highly protein bound (>95%) Although the presence of *in vivo* interaction of docetaxel with concomitantly administered medicinal product has not been investigated formally, *in vitro* interactions with highly protein bound agents such as erythromycin, diphtherydiamine, progesterone, propofene, phenytoin, salicylate, sulfamethoxazole and sodium valproate did not affect protein binding of docetaxel. In addition, dexamethasone did not affect protein binding of docetaxel. Docetaxel did not influence the binding of digoxin.

The pharmacokinetics of docetaxel, doxorubicin and cyclophosphamide were not influenced by the co-administration. Limited data from a single uncontrolled study were suggestive of an interaction between docetaxel and carboplatin. When combined to docetaxel, the clearance of carboplatin was about 50% lower than values previously reported for carboplatin monotherapy.

#### K. Statement on usage during pregnancy and lactation

##### Pregnancy

There is no information on the use of docetaxel in pregnant women. Docetaxel has been shown to be both embryotoxic and fetotoxic in rabbits and rats, and to reduce fertility in rats. As with other cytotoxic antineoplastic pro-drugs, docetaxel can cause foetal harm when administered to pregnant women. Therefore, docetaxel must not be used during pregnancy unless clearly indicated.

Women of childbearing age receiving docetaxel should be advised to avoid becoming pregnant, and to inform the treating physician immediately should this occur.

##### Breast-feeding

Docetaxel is a lipophilic substance but it is not known whether it is excreted in human milk. Consequently, because of the potential for adverse reactions in nursing infants, breast feeding must be discontinued for the duration of docetaxel therapy.

##### Contraception in males and females

An effective method of contraception should be used during treatment.

##### Fertility

In non-clinical studies, docetaxel had gonotoxic effects and may alter male fertility. Therefore, men being treated with docetaxel are advised not to father a child and to stop fathering for up to 6 months after treatment and to seek advice on conservation of sperm prior to treatment.

#### L. Adverse Effects Undesirable Effects

##### Summary of the safety profile for all indications

The adverse reactions considered to be possibly or probably related to the administration of docetaxel have been obtained in:

- 1312 and 121 patients who received 100 mg/m<sup>2</sup> and 75 mg/m<sup>2</sup> of docetaxel as a single agent respectively.
- 259 patients who received docetaxel in combination with doxorubicin.
- 486 patients who received docetaxel in combination with cisplatin.
- 92 patients treated with docetaxel in combination with trastuzumab.
- 255 patients who received docetaxel in combination with capecitabine.
- 332 patients (TAX 327) who received docetaxel in combination with prednisone or prednisolone (clinically important treatment-related adverse events are presented).
- 1276 patients (744 and 532 in TAX316 and GEECAM 9805 respectively) who received docetaxel in combination with doxorubicin and cyclophosphamide (clinically important treatment-related adverse events are presented).
- 300 gastric adenocarcinoma patients (221 patients in the phase III part of the study and 79 patients in the phase II part) who received docetaxel in combination with cisplatin and 5-fluorouracil (clinically important treatment-related adverse events are presented).
- 174 and 251 head and neck cancer patients who received docetaxel in combination with cisplatin and 5-fluorouracil (clinically important treatment-related adverse events are presented).

These reactions were described using the NCI Common Toxicity Criteria (grade 3 = G3; grade 3–4 = G3/4; grade 4 = G4), the COSTART and the MedDRA terms. Frequencies are defined as: very common (>10); common (>1/100 to <1/10); uncommon (>1/1,000 to <1/100); rare (>1/10,000 to <1/1,000); very rare (<1/10,000), not known (cannot be estimated from available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

The most commonly reported adverse reactions of docetaxel alone are: neutropenia (which was reversible and not cumulative; the median day to nadir was 7 days) and the median duration of severe neutropenia (<500 cells/mm<sup>3</sup> was 7 days), anaemia, alopecia, nausea, vomiting, stomatitis, diarrhoea and asthenia. The severity of adverse events of docetaxel may be increased when given in combination with other chemotherapeutic agents.

For combination with trastuzumab, adverse events (all grades) reported to >10% are displayed. There was an increased incidence of SAEs (40% vs. 31%) and Grade 4 AEs (3% vs. 2%) in the trastuzumab combination are compared to docetaxel monotherapy.

For combination with capecitabine, the most frequent treatment-related undesirable effects (G2/3) reported in a phase III study in breast cancer patients failing anthracycline treatment are presented (see capecitabine summary of product characteristics).

For combination with ADT and with prednisone or prednisolone (STAMPEDE study), adverse events occurring over the 6 cycles of treatment with docetaxel and having at least 2% higher incidence in the docetaxel treatment arm by comparison to the control arm, are presented, using the CTCAE grading scale.

The following adverse reactions are frequently observed with docetaxel:

##### Immune system disorders

Hypersensitivity reactions have generally occurred within a few minutes following the start of the infusion of docetaxel and were usually mild to moderate. The most frequently reported symptoms were flushing, rash with or without pruritus, chest tightness, back pain, dyspnoea and fever or chills. Severe reactions were characterised by hypotension and/or bronchospasm or generalised rash/erythema.

##### Neurology system disorders

The development of severe peripheral neuropathy requires a reduction of dose Mild to moderate neuro-sensory signs are characterised by paraesthesia, dysesthesia or pain including burning. Neuro-sensory events are mainly characterised by weakness.

##### Skin and subcutaneous tissue disorders

Reversible cutaneous reactions have been observed and were generally considered as mild to moderate. Reactions were characterised by rash including localized eruptions mainly on the feet and hands (including severe hand and foot syndromes), but also on the arms, face, thorax, and frequently associated with pruritus. Eruptions generally occurred within one week after the docetaxel infusion. Less frequently, severe symptoms such as eruptions followed by desquamation which rarely lead to interruption or discontinuation of docetaxel treatment were reported. Severe nail disorders are characterised by hypo- or hyperpigmentation and sometimes pain and onycholysis.

##### General disorders and administration site conditions

Infusion site reactions were generally mild and consisted of hyperpigmentation, inflammation, redness or dryness of the skin, phlebitis or extravasation and swelling of the vein.

Fluid retention includes events such as peripheral oedema and less frequently pleural effusion, pericardial effusion, ascites and weight gain. The peripheral oedema usually starts at the lower extremities and may become generalised with a weight gain of 3 kg or more. Fluid retention is cumulative in incidence and severity.

##### Tabulated list of adverse reactions in breast cancer for docetaxel 100 mg/m<sup>2</sup> single agent

| MedDRA system organ classes                          | Very common adverse reactions                                                                              | Common adverse reactions                                                                                                                 | Uncommon adverse reactions  |
|------------------------------------------------------|------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------|
| Infections and infestations                          | Infection (G3/4: 5.7%); including sepsis and pneumonia; fatal in 1.7%                                      | Infection associated with G4 neutropenia (G3/4: 4.6%)                                                                                    |                             |
| Blood and lymphatic system disorders                 | Neutropenia (G4: 76.4%); Anaemia (G3/4: 4.9%); Febrile neutropenia                                         | Thrombocytopenia (G4: 0.2%)                                                                                                              |                             |
| Immune system disorders                              | Hypersensitivity (G3/4: 5.3%)                                                                              |                                                                                                                                          |                             |
| Metabolism and nutrition disorders                   | Anorexia                                                                                                   |                                                                                                                                          |                             |
| Nervous system disorders                             | Peripheral sensory neuropathy (G3: 4.7%); Peripheral motor neuropathy (G3/4: 4%); Dysgeusia (severe: 0.0%) |                                                                                                                                          |                             |
| Cardiac disorders                                    |                                                                                                            | Arrhythmia (G3/4: 0.7%)                                                                                                                  | Cardiac failure             |
| Respiratory, thoracic and mediastinal disorders      | Dyspnoea (severe: 2.7%)                                                                                    |                                                                                                                                          |                             |
| Gastrointestinal disorders                           | Stomatitis (G3/4: 5.2%); Diarrhoea (G3/4: 4%); Nausea (G3/4: 4%); Vomiting (G3/4: 3.4%)                    | Constipation (severe: 0.2%); Abdominal pain (severe: 1%); Gastrointestinal haemorrhage (severe: 0.3%)                                    | Oesophagitis (severe: 0.4%) |
| Skin and subcutaneous tissue disorders               | Alopecia: Skin reactions (G3/4: 5.9%); Nail disorders (severe: 2.6%)                                       |                                                                                                                                          |                             |
| Musculoskeletal and connective tissue disorders      | Myalgia (severe: 1.4%)                                                                                     | Arthralgia                                                                                                                               |                             |
| General disorders and administration site conditions | Fluid retention (severe: 6.5%); Asthenia (severe: 11.2%); Pain                                             | Infusion site reaction; Non-cardiac chest pain (severe: 0.4%)                                                                            |                             |
| Investigations                                       |                                                                                                            | G3/4 Blood bilirubin increased (<5%); G3/4 Blood alkaline phosphatase increased (<4%); G3/4 AST increased (<2%) G3/4 ALT increased (<2%) |                             |

Description of selected adverse reactions in breast cancer for docetaxel 100 mg/m<sup>2</sup> single agent

##### Blood and lymphatic system disorders

Rare: Needing episodes associated with grade 3/4 thrombocytopenia

##### Not known: Leucopenia

##### Neurology system disorders

Reversibility data are available among 35.3% of patients who developed neurotoxicity following docetaxel treatment at 100 mg/m<sup>2</sup> as single agent. The events were spontaneously reversible within 3 months.

##### Skin and subcutaneous tissue disorders

Very rare: one case of alopecia non-reversible at the end of the study. 73% of the cutaneous reactions were reversible within 21 days.

##### General disorders and administration site conditions

The median cumulative dose to treatment discontinuation was more than 1,000 mg/m<sup>2</sup> and the median time to fluid retention reversibility was 16.4 weeks (range 0 to 42 weeks). The onset of moderate and severe retention is delayed (median cumulative dose: 818.9 mg/m<sup>2</sup>) in patients with predemarcation compared with patients without predemarcation (median cumulative dose: 489.7 mg/m<sup>2</sup>); however, it has been reported in some patients during the early course of therapy.

##### Tabulated list of adverse reactions in non-small cell lung cancer for docetaxel 75 mg/m<sup>2</sup> single agent

| MedDRA system organ classes                          | Very common adverse reactions                                                             | Common adverse reactions                 |
|------------------------------------------------------|-------------------------------------------------------------------------------------------|------------------------------------------|
| Infections and infestations                          | Infections (G3/4: 5%)                                                                     | Febrile neutropenia                      |
| Blood and lymphatic system disorders                 | Neutropenia (G4: 91.7%); Anaemia (G3/4: 10.8%); Thrombocytopenia (G3/4: 1.7%)             |                                          |
| Immune system disorders                              |                                                                                           | Hypersensitivity (not severe)            |
| Metabolism and nutrition disorders                   | Anorexia                                                                                  |                                          |
| Nervous system disorders                             | Peripheral sensory neuropathy (G3/4: 0.8%)                                                | Peripheral motor neuropathy (G3/4: 2.5%) |
| Cardiac disorders                                    |                                                                                           | Arrhythmia (no severe)                   |
| Vascular disorders                                   |                                                                                           | Hypotension                              |
| Gastrointestinal disorders                           | Nausea (G3/4: 3%); Stomatitis (G3/4: 1.7%); Vomiting (G3/4: 0.8%); Diarrhoea (G3/4: 1.7%) | Constipation                             |
| Skin and subcutaneous tissue disorders               | Alopecia: Skin reaction (G3/4: 0.8%)                                                      | Nail disorders (severe: 0.8%)            |
| Musculoskeletal and connective tissue disorders      |                                                                                           | Myalgia                                  |
| General disorders and administration site conditions | Asthenia (severe: 12.4%); Fluid retention (severe: 0.8%); Pain                            |                                          |
| Investigations                                       |                                                                                           | G3/4 Blood bilirubin increased (<2%)     |

##### Tabulated list of adverse reactions in breast cancer for docetaxel 75 mg/m<sup>2</sup> in combination with doxorubicin

| MedDRA system organ classes                          | Very common adverse reactions                                                           | Common adverse reactions                                                                  | Uncommon adverse reactions                         |
|------------------------------------------------------|-----------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|----------------------------------------------------|
| Infections and infestations                          | Infection (G3/4: 7.2%)                                                                  |                                                                                           |                                                    |
| Blood and lymphatic system disorders                 | Neutropenia (G4: 91.7%); Anaemia (G3/4: 9.4%); Febrile neutropenia                      | Thrombocytopenia (G3/4: 0.8%)                                                             |                                                    |
| Immune system disorders                              |                                                                                         | Hypersensitivity (G3/4: 1.2%)                                                             |                                                    |
| Metabolism and nutrition disorders                   | Anorexia                                                                                |                                                                                           |                                                    |
| Nervous system disorders                             | Peripheral sensory neuropathy (G3: 0.8%)                                                | Peripheral motor neuropathy (G3/4: 0.4%)                                                  |                                                    |
| Cardiac disorders                                    |                                                                                         | Cardiac failure; Arrhythmia (no severe)                                                   |                                                    |
| Vascular disorders                                   |                                                                                         |                                                                                           | Hypotension                                        |
| Gastrointestinal disorders                           | Nausea (G3/4: 5%); Stomatitis (G3/4: 7.8%); Diarrhoea (G3/4: 6.2%); Vomiting (G3/4: 2%) | Constipation                                                                              |                                                    |
| Skin and subcutaneous tissue disorders               | Alopecia: Nail disorders (severe: 0.6%); Skin reaction (no severe)                      | Myalgia                                                                                   |                                                    |
| Musculoskeletal and connective tissue disorders      |                                                                                         |                                                                                           |                                                    |
| General disorders and administration site conditions | Asthenia (severe: 9.5%); Fluid retention (severe: 1.2%); Pain                           | Infusion site reaction                                                                    |                                                    |
| Investigations                                       |                                                                                         | G3/4 Blood bilirubin increased (<2.5%); G3/4 Blood alkaline phosphatase increased (<2.5%) | G3/4 AST increased (<1%); G3/4 ALT increased (<1%) |

##### Tabulated list of adverse reactions in non-small cell lung cancer for docetaxel 75 mg/m<sup>2</sup> in combination with cisplatin

| MedDRA system organ classes                          | Very common adverse reactions                                                             | Common adverse reactions                                         | Uncommon adverse reactions                                                  |
|------------------------------------------------------|-------------------------------------------------------------------------------------------|------------------------------------------------------------------|-----------------------------------------------------------------------------|
| Infections and infestations                          | Infection (G3/4: 5.7%)                                                                    |                                                                  |                                                                             |
| Blood and lymphatic system disorders                 | Neutropenia (G4: 51.7%); Anaemia (G3/4: 6.9%); Thrombocytopenia (G3/4: 0.5%)              | Febrile neutropenia                                              |                                                                             |
| Immune system disorders                              |                                                                                           | Hypersensitivity (G3/4: 2.5%)                                    |                                                                             |
| Metabolism and nutrition disorders                   | Anorexia                                                                                  |                                                                  |                                                                             |
| Nervous system disorders                             | Peripheral sensory neuropathy (G3: 3.7%); Peripheral motor neuropathy (G3/4: 2%)          |                                                                  |                                                                             |
| Cardiac disorders                                    |                                                                                           | Arrhythmia (G3/4: 0.7%)                                          | Cardiac failure                                                             |
| Vascular disorders                                   |                                                                                           | Hypotension (G3/4: 0.7%)                                         |                                                                             |
| Gastrointestinal disorders                           | Nausea (G3/4: 9.6%); Vomiting (G3/4: 7.8%); Diarrhoea (G3/4: 6.2%); Stomatitis (G3/4: 2%) | Constipation                                                     |                                                                             |
| Skin and subcutaneous tissue disorders               | Alopecia: Nail disorders (severe: 0.7%); Skin reaction (G3/4: 0.2%)                       |                                                                  |                                                                             |
| Musculoskeletal and connective tissue disorders      | Myalgia (severe: 0.5%)                                                                    |                                                                  |                                                                             |
| General disorders and administration site conditions | Asthenia (severe: 9.9%); Fluid retention (severe: 1.3%); Fever (G3/4: 1.2%)               | Infusion site reaction                                           |                                                                             |
| Investigations                                       |                                                                                           | G3/4 Blood bilirubin increased (2.1%); G3/4 ALT increased (1.3%) | G3/4 AST increased (0.5%); G3/4 Blood alkaline phosphatase increased (1.3%) |

##### Tabulated list of adverse reactions in breast cancer for docetaxel 100 mg/m<sup>2</sup> in combination with trastuzumab

| MedDRA system organ classes                          | Very common adverse reactions                                                                                                     | Common adverse reactions |
|------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------|--------------------------|
| Blood and lymphatic system disorders                 | Neutropenia (G3/4: 32%); Febrile neutropenia (includes neutropenia associated with fever and antibodies not to neutropenic signs) |                          |
| Metabolism and nutrition disorders                   | Anorexia                                                                                                                          |                          |
| Psychiatric disorders                                | Insomnia                                                                                                                          |                          |
| Nervous system disorders                             | Paraesthesia; Headache; Dysgeusia; Hypoesthesia                                                                                   |                          |
| Vascular disorders                                   | Lymphedema                                                                                                                        | Cardiac failure          |
| Respiratory, thoracic and mediastinal disorders      | Epistaxis; Pharyngolaryngeal pain; Nasopharyngitis; Dyspnoea; Cough; Rhinitis                                                     |                          |
| Gastrointestinal disorders                           | Nausea; Diarrhoea; Vomiting; Constipation; Stomatitis; Dyspepsia; Abdominal pain                                                  |                          |
| Skin and subcutaneous tissue disorders               | Alopecia; Erythema; Rash; Nail disorders                                                                                          |                          |
| Musculoskeletal and connective tissue disorders      | Myalgia; Arthralgia; Pain in extremity; Bone pain; Back pain                                                                      |                          |
| General disorders and administration site conditions | Asthenia; Oedema peripheral; Pyrexia; Fatigue; Muscular inflammation; Pain; Infusion like illness; Chest pain; Chills             | Lethargy                 |
| Investigations                                       | Weight increased                                                                                                                  |                          |

##### Description of selected adverse reactions in breast cancer for docetaxel 100 mg/m<sup>2</sup> in combination with trastuzumab

Symptomatic cardiac failure was reported in 2.2% of the patients who received docetaxel plus trastuzumab compared to 0% of patients given docetaxel alone.

In the docetaxel plus trastuzumab arm, 64% had received a prior anthracycline as adjuvant therapy compared with 55% in the docetaxel arm alone.

##### Blood and lymphatic system disorders

Very common: Haematological toxicity was increased in patients receiving trastuzumab and docetaxel, compared with docetaxel alone (32% grade 3/4 neutropenia versus 22% using NCI-CTC criteria. Note that this is likely to be an underestimation since docetaxel alone at a dose of 100 mg/m<sup>2</sup> is known to result in neutropenia in 97% of patients, 76% grade 4, based on nadir blood counts. The incidence of febrile neutropenia/neutropenic sepsis was also increased in patients treated with Herceptin plus docetaxel (23% versus 17% for patients treated with docetaxel alone).

##### Tabulated list of adverse reactions in breast cancer for docetaxel 75 mg/m<sup>2</sup> in combination with capecitabine

| MedDRA system organ classes                                                                                                                                              | Very common adverse reactions                                                                                                                        | Common adverse reactions                                                                 |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|
| Infections and infestations                                                                                                                                              | Infection (G3/4: 3.3%)                                                                                                                               | Oral candidiasis (G3/4: <1%)                                                             |
| Blood and lymphatic system disorders                                                                                                                                     | Neutropenia (G3/4: 6.7%); Anaemia (G3/4: 10%)                                                                                                        | Thrombocytopenia (G3/4: 0.6%); Febrile neutropenia                                       |
| Metabolism and nutrition disorders                                                                                                                                       | Anorexia (G3/4: 1%); Decreased appetite                                                                                                              | Dehydration (G3/4: 2%)                                                                   |
| Nervous system disorders                                                                                                                                                 | Dysgeusia (G3/4: <1%); Paraesthesia (G3/4: 1%)                                                                                                       | Dizziness; Headache (G3/4: <1%); Neuropathy peripheral                                   |
| Eye disorders                                                                                                                                                            | Lacrimation increased                                                                                                                                |                                                                                          |
| Respiratory, thoracic and mediastinal disorders                                                                                                                          | Pharyngolaryngeal pain (G3/4: 2%)                                                                                                                    | Dyspnoea (G3/4: 1%); Cough (G3/4: <1%); Dyspnoea (G3/4: <1%)                             |
| Gastrointestinal disorders                                                                                                                                               | Nausea (G3/4: 1.8%); Diarrhoea (G3/4: 14%); Stomatitis (G3/4: 0%); Vomiting (G3/4: 4%); Constipation (G3/4: 1%) Abdominal pain (G3/4: 2%); Dyspepsia | Abdominal pain upper; Dry mouth                                                          |
| Skin and subcutaneous tissue disorders                                                                                                                                   | Hand-foot syndrome (G3/4: 24%); Alopecia (G3/4: 0.6%); Nail disorders (G3/4: 2%)                                                                     | Dermatologic Rash erythematous (G3/4: <1%); Nail discoloration; Onychomycosis (G3/4: 1%) |
| Musculoskeletal and connective tissue disorders                                                                                                                          | Myalgia (G3/4: 2%) Arthralgia (G3/4: 1%)                                                                                                             | Pain in extremity (G3/4: <1%); Back pain (G3/4: 1%)                                      |
| General disorders and administration site conditions                                                                                                                     | Asthenia (G3/4: 3%); Pyrexia (G3/4: 1%); Fatigue/weakness (G3/4: 2%); Oedema peripheral (G3/4: 0.6%)                                                 | Leathargy; Pain                                                                          |
| Investigations                                                                                                                                                           |                                                                                                                                                      | Weight decreased; G3/4 Blood bilirubin increased (0%)                                    |
| Tabulated list of adverse reactions in metastatic castration-resistant prostate cancer for docetaxel 75 mg/m <sup>2</sup> in combination with prednisone or prednisolone |                                                                                                                                                      |                                                                                          |
| MedDRA system organ classes                                                                                                                                              | Very common adverse reactions                                                                                                                        | Common adverse reactions                                                                 |
| Infections and infestations                                                                                                                                              | Infection (G3/4: 3.3%)                                                                                                                               |                                                                                          |
| Blood and lymphatic system disorders                                                                                                                                     | Neutropenia (G3/4: 3.2%); Anaemia (G3/4: 4.9%)                                                                                                       | Thrombocytopenia (G3/4: 0.6%); Febrile neutropenia                                       |
| Immune system disorders                                                                                                                                                  | Anorexia (G3/4: 0.6%)                                                                                                                                | Hypersensitivity (G3/4: 0.6%)                                                            |
| Metabolism and nutrition disorders                                                                                                                                       |                                                                                                                                                      |                                                                                          |
| Nervous system disorders                                                                                                                                                 | Peripheral sensory neuropathy (G3/4: 12%); Dysgeusia (G3/4: 0%)                                                                                      | Peripheral motor neuropathy (G3/4: 0.3%)                                                 |
| Eye disorders                                                                                                                                                            |                                                                                                                                                      | Lacrimation increased (G3/4: 0.6%)                                                       |
| Cardiac disorders                                                                                                                                                        |                                                                                                                                                      | Cardiac left ventricular function decrease (G3/4: 0.3%)                                  |
| Respiratory, thoracic and mediastinal disorders                                                                                                                          |                                                                                                                                                      | Epididymis (G3/4: 0%); Dyspnoea (G3/4: 0.6%); Cough (G3/4: 0.3%)                         |