

Essliver Capsule

Essential Phospholipids 300 mg
(Phospholipids from soya-beans)

Composition:

Each capsule contains:
Essential Phospholipids 300 mg
Source of capsule shell is bovine Gelatin.

Dosage Form: Solid oral capsule

Product Description:

White cap with white body, size "0" hard gelatin capsules. Brownish cream colored powder.

Pharmacological Action:

Pharmacodynamics Properties:

ATC CODE: V03AX - OTHER THERAPEUTIC PRODUCTS

Among the pharmacodynamic properties were reported hepatoprotective effects found in numerous experimental models into acute liver damage, for example induced by ethanol, alcy alcohol, carbontetrachloride, paracetamol and galactosamine. Moreover, in chronic models (ethanol, thioacetamide, organic solvents) was seen also the inhibition of steatosis and fibrosis. As active principle has been suggested accelerated membrane regeneration and stabilisation, inhibited lipid peroxidation and inhibited collagen synthesis. Specific investigations into human Pharmacodynamics are not available.

Pharmacokinetics Properties:

Animal experiments into the pharmacokinetics showed that more than 90% of the orally applied radiolabelled soya-bean phospholipids are absorbed in the small intestine. Most of it is split by phospholipase A to 1-acyllysophosphatidylcholine, 50% of which is reacylated immediately into polyunsaturated phosphatidylcholine still during the process of absorption in the intestinal mucosa. This polyunsaturated phosphatidylcholine reaches the blood via the lymph pathway and from there mainly bound to HDL - it passes in particular to the liver. Tests into human pharmacokinetics were performed a.o. with radioactively labeled dilinoleoylphosphatidylcholine (3H and 14C). The choline moiety was 3H-labeled and the linoleic acid had the 14C-label. The maximum 3H concentration was achieved after 6 to 24 hours and amounted to 19.9% of the dose. The half-life for the choline component was 66 hours. The maximum 14C concentration was achieved after 4 to 12 hours and amounted to 27.9% of the dose. The half-life for this component was 32 hours. In the faeces were found 2% of the 3H and 4.5% of the 14C label, in the urine 6% of the 3H and only minor amount of the 14C label. These results show that both isotopes are absorbed to over 90% in the intestine.

Indications :

Nutritional support in the management of damaged liver (due to chronic disease, liver cirrhosis, fatty liver and intoxication by hepatotoxic substances)

Dosage & Administration:

In the beginning take 3 times daily 2 capsules (600 mg phospholipids from soya-beans).
Not more than 6 capsules a day should be taken (1800 mg phospholipids from soya-beans).
After some time the dose may be reduced to 3 times daily 1 capsule (300 mg phospholipids from soya-beans) (maintenance dose).

Contraindications:

Known hypersensitivity to soya-bean preparations or to any of the excipients.

Interactions with Other Medicaments:

An interaction of Essential Phospholipids with anticoagulants cannot be excluded; therefore, dose adjustment of the anticoagulant might be necessary. The doctor should be consulted in case of simultaneous application.

Warning & Precautions:

Special Precautions: Due to the content in soya bean oil, Essential Phospholipids may provoke severe allergic reactions. Essential Phospholipids therapy is not a substitute for the avoidance of the noxious agent causing liver damage (e.g., alcohol). In chronic hepatitis, the adjuvant therapy with soya bean Phospholipids is justified only when improved subjective well-being manifests during therapy. The physician should be consulted when complaints aggravate or when other unclear complaints occur.

Use for Pregnancy & Lactation:

Preparations from soya beans are largely used in human food, and so far, no events have been reported that would suggest any risk during pregnancy. Specific results from investigation are not available, therefore, the use of Essential Phospholipids is not recommended during pregnancy and lactation.

Side Effects:

For the evaluation of any adverse effects, the following incidences are considered: Very frequent (>1/10), frequent (>1/100 to <1/10), occasional (>1/1,000 to <1/100), rare (>1/10,000 to <1/1,000), very rare (<1/10,000). Occasionally, the administration of Essliver may provoke gastrointestinal disorders such as stomach complaints, soft stool and diarrhoea. On very rare occasions, allergic reactions may occur, such as exanthema and urticaria. The doctor or pharmacist should be informed about any undesirable effect not listed in the previously mentioned. Administration of Essliver should be discontinued in the event of one of the previously mentioned side effects, especially if hypersensitivity reactions occur. The patient should consult the doctor to decide about the severity and any measures that might be necessary.

Symptoms and Treatment of Overdose:

Neither overdose reaction nor symptom of intoxication has been reported to date with Essential Phospholipids.

Effects on Ability to Drive and Use Machine:

Normal use of the product is not known to have any effects on ability to drive or use machines.

Storage :

Store below 30°C.

Active Ingredients : Essential Phospholipids

Inactive Ingredients : Disodium EDTA, Butylated hydroxyl Toluene, Purified Talcum & Aluminum Magnesium Silicate Hydrate,

Route of Administration: For oral use

Shelf Life: 36 months

Presentation :

3 x 10 Capsules

MALNO.

Serial No.:*****

Date of Revision: 06/04/2023

Product Registration Holder

Healol

74-3, Jalan Wangsa Delima 6,
KLSC Wangsa Maju, 53300,
Kuala Lumpur, Malaysia



Manufactured in India by:

Nabros Pharma Pvt. Ltd.

Survey No. 110/A/2, Amit Farm,
Jain Upasrya, Near Coca Cola Factory,
N.H. No. 8, Kajipura - 387411, Kheda,
INDIA

06/04/2023

Size : 150 x 240 mm

ARTWORK CODE