

# SN GENTA 300 KIU/G

WATER SOLUBLE POWDER  
(FOR VETERINARY USE ONLY)

## COMPOSITION

Gentamycin Sulphate ..... 300,000 IU/g

## PRODUCT DESCRIPTION

The finished product is white or almost white color powder.

## TARGET SPECIES

Pigs

## PHARMACODYNAMIC

Gentamicin is a bactericidal aminoglycoside antibacterial, whose antimicrobial action results from acting on the 30 S subunit of the ribosomes, preventing protein synthesis. Prevents the entire initiation phase, disrupting the ordering of messenger RNA and causing an incorrect reading of the genetic code by the transfer RNA. It also disrupts the permeability of the bacterial membrane.

It is active against: *Escherichia coli*, *Brachyspira hyodysenteriae*

Anaerobic organisms are resistant. Cross resistance occurs between members of the group.

## PHARMACOKINETIC

Gentamicin is barely absorbed orally and consequently plasma concentrations are practically undetectable. It remains in the gastrointestinal tract and its distribution to the other organs and tissues is reduced.

## INDICATIONS

Pigs: In weaned animals.

Treatment of digestive infections caused by non-invasive *Escherichia coli* strains sensitive to gentamicin.

Treatment of swine dysentery caused by microorganisms sensitive to gentamicin.

## WARNINGS AND PRECAUTIONS

**Special warnings for each target species:**

None.

**Special precautions for use:**

**i. Special precautions for use in animals:**

Good clinical practice requires that treatment be based on sensitivity tests of bacteria isolated from diseased animals. If this is not possible, treatment should be based on local (regional, farm level) epidemiological information on the sensitivity of different strains of bacterial species commonly involved in the infectious process.

The use of the drug under conditions other than those recommended in the label/ package insert may increase the

prevalence of bacteria resistant to gentamicin and decrease the efficacy of treatment with aminoglycosides as a consequence of the appearance of cross-resistance.

When using this drug, the official recommendations (national or regional) on the use of antimicrobials should be considered.

## ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Aminoglycosides can cause hypersensitivity (allergy) after inhalation, ingestion, or skin contact. Avoid skin contact with this product if you know you are sensitized or if you have been advised not to work with such preparations. People with known hypersensitivity to gentamicin or any of the excipients should avoid contact with the veterinary medicinal product.

The medicine can be irritating to the respiratory tract and some of the dust produced can be inhaled. The medicine must be handled with care to avoid dust production. Use in a well-ventilated area away from drafts. Avoid inhaling dust during incorporation into water. Use personal protective equipment consisting of a dust mask, when handling the veterinary medicine.

This product can be irritating to the eyes and mucous membranes. Avoid direct contact with the skin and eyes. Wear personal protective equipment consisting of gloves, work overalls and approved safety glasses when handling the veterinary medicinal product. If accidental contact with skin or eyes occurs, immediately wash the affected area with plenty of water. If irritation persists, see a doctor immediately and show the package leaflet or label.

Do not smoke, eat or drink when handling the veterinary medicine.

Wash hands after use.

Due to the reproductive and / or fetus toxicity of gentamicin and suspected potential harm to fertility and / or the fetus of one of the excipients, the product cannot be prepared or administered by pregnant women.

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak

## CONTRAINDICATIONS

Do not use in cases of hypersensitivity to gentamicin or to any of the excipients.

## ADMINISTRATION & DOSAGE

Administration in drinking water.

Administer 2,000 IU of gentamicin sulfate / kg bw (equivalent to 66.7 mg of medicine for every 10 kg of bw) for 3 consecutive days.

The weight of the animals should be determined as accurately as possible to avoid insufficient dosing.

Water consumption depends on the clinical situation of the animal and the time of year. To ensure a correct dosage, the concentration of gentamicin in the water will be adjusted taking into account daily consumption.

Based on the recommended dose, the number and weight of animals to be treated, the exact daily dose of the medicine should be calculated using the following formula:

$$\frac{\text{g of medicine / liter of drinking water}}{\text{IU of active substance in g of medicine (IU / g) x Average water consumption (liters / day)}} = \frac{\text{Active substance dose (IU / kg bw / day) x mean weight of animals (kg)}}{\text{IU of active substance in g of medicine (IU / g) x Average water consumption (liters / day)}}$$

Medicated water must be prepared daily.

## INTERACTION WITH OTHER MEDICAMENTS

Do not administer together with other aminoglycosides or bacteriostatic antibiotics.

## PREGNANCY AND LACTATION

Its use is not recommended during pregnancy or lactation.

## OVERDOSE AND TREATMENT

No information.

## SIDE EFFECT

Intestinal dysbiosis may occur.

## WITHDRAWAL TIME

Meat: 14 days.

**SHELF LIFE:** 36 months

**SHELF LIFE after first opening & reconstitution:** 24 hours

## STORAGE

Store in temperature not exceeding 30°C. Strictly avoid light & heat exposure.

## PACKAGING

The powder filled 1 kg in aluminum foil.

**DATE OF REVISION:** 04 Apr 2023

Product registration holder & Manufactured by:



**SHENNONG ANIMAL HEALTH (M) SDN BHD**

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,  
Taman Perindustrian USJ 1,  
47610 Subang Jaya, Selangor.