

EM-TAIKOMIX 10% GRANULES

Description and Composition

EM-TAIKOMIX 10% GRANULES is a pink, free-flowing granules. Each kg contains 100g Florfenicol as its active ingredient and 1.5g Calcium Propionate as preservative.

Pharmacodynamics

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic. However, bactericidal activity has been demonstrated *in-vitro* against *Pasteurella multocida* when florfenicol is present at concentrations above the MIC for 4 to 12 hours.

In-vitro testing has shown that florfenicol is active against the bacterial pathogens most commonly isolated in respiratory diseases in pigs, including *Pasteurella multocida*.

A total of 193 *Pasteurella multocida* isolates from the respiratory tract of swine were collected between 2002 and 2003 in France, Spain, Greece, Germany, the United Kingdom and Belgium. The Minimal Inhibitory Concentration (MIC) of florfenicol against the target pathogen ranges from 0.25 to 1µg / ml with a MIC90 of 0.5µg / ml. The only mechanisms of chloramphenicol resistance that are known to have significant clinical relevance are CAT (Chloramphenicol Acetyl Transferase)-mediated inactivation and efflux-pump resistance. Of these, only some of the efflux mediated resistance would also confer resistance to florfenicol and thus have the potential to be affected by florfenicol use in animals.

Pharmacokinetics

After administration to pigs by gavage at 10mg / kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5 µg/ml were reached approximately 3 hours after dosing. The terminal half-life was between 3 and 4 hours. When pigs were given free access, for 5 days, to feed medicated premix at the recommended dose of 10mg / kg serum florfenicol concentrations exceeds 1µg / ml for more than 16 hours each day of treatment.

Florfenicol is well absorbed when administered orally and following distribution it is rapidly eliminated in the urine and faeces in a ratio of 3:1. A fraction is excreted unchanged and the rest is metabolised into 5 major metabolites. After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

Indication

For the treatment and metaphylaxis of swine respiratory disease caused by *Pasteurella multocida* susceptible to florfenicol in infected herds. The presence of the disease in the herd must be established before the product is used.

Recommended Dose

To be administered orally, in medicated feeding stuff.

Pig (fattening pigs): 10mg florfenicol per kg body weight (bw) (equivalent to 100mg product) per day administered for 5 consecutive days.

For a daily feed intake of 50g / kg body weight, this dosage corresponds to a rate of incorporation of 2kg premix per ton of feed, i.e. 200ppm florfenicol.

Route of Administration

For oral administration via feed.

Contraindication

Do not administer to boars intended for breeding.

Do not use in case of hypersensitivity to the active substance or any of the excipients.

Warning and Precautions

Special warnings for each target species

Animals showing a decreased appetite and/or a poor general condition should be treated by the parenteral route.

Special precautions for use

Special precautions for use in animals

The product should be used in conjunction with susceptibility testing and take into account official and local policy relating to the use of antimicrobials.

This premix is intended for the manufacturing of solid medicated feed and cannot be used as it is; the incorporation rate of the premix in feed cannot be lower than 2kg / ton.

Treatment should not exceed 5 days.

In a field clinical study, within a week after the administration of the last dose, the incidence of pigs presenting either mild

depression and/or mild dyspnea and/or pyrexia (40°C) was approx. 20% in the initially severely ill animals.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Skin sensitisation may occur.

Avoid skin contact.

Do not handle this product in case of known sensitization to active substance or any of the excipients.

Handle this product with care to avoid exposure during incorporation of premix into feed and administration of feed to animals, taking all recommended precautions.

Wear either a disposable half-mask respirator or a non-disposable respirator with a filter, chemically resistant gloves, protective coveralls and goggles while incorporating the premix into feed.

Wear gloves and do not smoke, eat, or drink when handling the product or medicated feed.

Wash hands thoroughly with soap and water after use of the product or medicated feed.

Rinse thoroughly with water in case of exposure.

If you develop symptoms following exposure such as skin rash, you should seek medical advice and take the package leaflet or the label with you.

Other precautions

Manure from treated swine must be stored for a minimum of one month before being spread and incorporated in fields.

Interaction with Other Medicaments

None known.

Pregnancy and Lactation

The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

Therefore the use is not recommended during pregnancy and lactation.

Side Effects

Commonly observed adverse effects are diarrhoea perianal inflammation and rectal eversion. Increased serum calcium may also be observed. These effects are transient, resolving on cessation of treatment.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))

- common (more than 1 but less than 10 animals in 100 animals treated)

- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)

- rare (more than 1 but less than 10 animals in 10,000 animals treated)

- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

Symptoms and Treatment of Overdose

In the event of overdose, a reduction in food and water consumption, together with a decrease in body weight may be observed. There may be an increase in refused feed and an increase in serum calcium.

Storage Condition

Store below 30°C.

Shelf Life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Periods

Meat and offal: 14 days

Packing

10kg, 25kg.

Revised date: 31.3.2026

	Manufacturer & Product Registration Holder:
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com