

CLOPIFEED 250G/KG POWDER

Description and Composition

CLOPIFEED 250G/KG POWDER is an off-white powder. Each kg contains 250g of Clopidol as its active ingredient.

Pharmacodynamics

Clopidol, also known as meticlorpindol or clopidol, is a pyridinol with broad coccidiostatic activity against early development of *Eimeria* spp. by inhibiting mitochondrial energy production in sporozoites and trophozoites.

Pharmacokinetics

There are few data on the metabolism of meticlorpindol in poultry, but it has been shown that in chickens the compound is absorbed from the GI tract to a significant extent and distributed widely to tissues, but is not extensively metabolized.

Indication

Chickens (broilers and layer replacements): Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*.

Turkeys: Aid in the prevention of leucocytozoonosis caused by *Leucocytozoon smithi*.

Recommended Dose

Chickens (broilers and layer replacements):

113.5g Clopidol per ton (907kg) of feed. Do not feed to chickens over 16 weeks of age. If intended for use as caged layers: 227g Clopidol per ton (907kg) of feed. Feed up to 16 weeks of age continuously as the sole ration.

Turkeys:

For turkeys grown for meat purposes only: Administer 113.5g or 227g (0.0125 or 0.025 percent) Clopidol per ton (907kg) of feed continuously as the sole ration depending upon management practices, degree of exposure and amount of feed eaten.

Route of Administration

For oral administration via feed.

Contraindication

Do not feed to laying hens. Do not feed to chickens reared for laying after 16 weeks of age. Do not administer clopidol to birds intended for breeding purposes.

Warning and Precautions

Special precautions for use:

Special precautions for use in animals

Do not use this medicated feed for treatment of outbreaks of Coccidiosis.

Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a 2-day period.

Special precautions to be taken by the person administering the veterinary medicinal product

When mixing, direct contact with the skin and mucous membranes should be avoided. Accidental ingestion should be avoided. Wear overalls, safety glasses, mask and impervious gloves when handling or mixing the product. Contaminated clothing should be removed and any splashes on to the skin should be washed off immediately. If accidental eye contact occurs, immediately rinse thoroughly with water. Seek medical advice if irritation persists. Wash hands after use.

Interaction with Other Medicaments

Do not mix with other anticoccidials.

Pregnancy and Lactation

Meticlorpindol fed to laying hens is deposited in egg albumen at concentrations about twice that found in egg yolk, and residues can persist for several weeks when the medication is given at high doses.

Side Effects

None under correct use of the product.

Symptoms and Treatment of Overdose

Unlikely to occur if use the product as per instructed.

Storage Condition

Store below 30°C.

Shelf life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Period

Chicken (broilers) and turkeys: 5 days before slaughter if given at the level of 0.025 percent in feed or reduce level to 0.0125 percent.

Packing

10kg and 25kg.

Revision date: 7.12.2023

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No.1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com