

LINCO SPECSURE POWDER

Description and Composition

LINCO SPECSURE POWDER is a white to off-white powder. Each gram contains 222mg Lincomycin (as hydrochloride) and 444mg of Spectinomycin (as hydrochloride) as its active ingredients.

Pharmacodynamics

Lincomycin is an antibiotic derived from *Streptomyces lincolnensis*. It belongs to the lincosamide group which also includes pirlimycin and clindamycin. Lincomycin is active mainly against Gram-positive bacteria. Lincomycin exerts its antibacterial action by inhibiting RNA-dependent protein synthesis by acting on the 50S subunit of the ribosome.

Spectinomycin is an aminocyclitol antibiotic produced by *Streptomyces spectabilis*. It exerts its bacteriostatic effect by binding to the 30S subunit of bacterial ribosomes and inhibiting the translation of protein synthesis.

Pharmacokinetics

Lincomycin: The pharmacokinetics of lincomycin have not apparently been extensively studied in veterinary species. The drug is rapidly absorbed from the gut, but only about 30–40% of the total dose is absorbed. Peak serum levels are attained about 2–4 hour after oral dosing. Lincomycin is distributed into most tissues. Lincomycin is bound from 57–72% to plasma proteins, depending on the drug's concentration. Lincomycin is partially metabolized in the liver. Unchanged drug and metabolites are excreted in the urine, feces and bile.

Spectinomycin: After oral administration only about 7% of the dose is absorbed, but the drug that remains in the GI tract is active. Tissue levels of absorbed drug are lower than those found in the serum. Absorbed drug is excreted via glomerular filtration into the urine mostly unchanged.

Indication

Poultry: As an aid in the prevention and treatment of Chronic Respiratory Disease (CRD, Air sacculitis) associated with mycoplasmal and coliform infections of growing chickens.

Recommended Dose

Poultry: Add 150g of product to 120 litres of drinking water (provides 833mg antibiotic activity per litre). For automatic medicators set to deliver 7.8ml of stock solution per litre of drinking water, prepare stock solution by dissolving 150g of product in 0.9 litres of water.

Prevention: To prevent losses due to CRD, administer at the above dosage for the first 3 to 5 days of life and again for one or two days at about 4 weeks of age. The longer period (5 days) of initial administration is recommended if disease has been a problem in the past on the premises.

Treatment: Begin treatment at the above dosage at the first sign of disease. Administer as the sole source of drinking water for 5 to 7 days.

Fresh medicated drinking water should be provided daily.

Route of Administration

By oral administration via drinking water.

Contraindication

Do not allow rabbits, hamsters, guinea pigs, horses, dairy cattle, or other ruminants access to feeds and/or drinking water containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects or metabolic disorders (i.e. Ketosis in dairy cattle).

Warning and Precautions**Special warning for each target species:**

Treated animals must not be slaughtered for use in food for at least 72 hours for poultry after the latest treatment with this drug. Not be used in laying birds.

Special precautions for use:Special precautions for use in animals

Do not allow rabbits, hamsters, guinea pigs, horses, dairy cattle, or other ruminants access to feeds and/or drinking water containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects or metabolic disorders (i.e. Ketosis in dairy cattle).

Special precautions to be taken by the person administering the veterinary medicinal product

Keep out of reach of children.

When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes.

Interaction with Other Medicaments

Antagonism has been reported when spectinomycin is used with chloramphenicol or tetracycline. Following drug interactions have either been reported or are theoretical in animals receiving lincomycin and may be of significance in veterinary patient: Cyclosporine, erythromycin, kaolin and neuromuscular blocking agents.

Pregnancy and Lactation

Must not be used in laying birds.

Side Effects

None known.

Symptoms and Treatment of Overdose

None known.

Storage Condition

Store below 30°C.

Shelf life

3 years. To be used within 24 hours after dilution.

Withdrawal Period

72 hours

Not be used in laying birds.

Packing

100g, 200g, 500g, 1kg

Revision date: 24.3.2023

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No.1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com