

NEO-VET -(Neomycin 20% w/w Water Soluble Powder)

CONTROLLED MEDICINE

Composition:

Each 1 kg contains:

Active Ingredient:

Neomycin (as neomycin sulphate) ... 200mg

Product description:

A light yellow colour crystalline powder

Target species:

Goat, cattle, sheep , swine and turkey

Pharmacokinetics

Neomycin is poorly absorbed from the gastrointestinal tract. Absorption from the gastrointestinal tract can be significant in neonates. 90% of neomycin is excreted in the faeces after oral administration.

Pharmacodynamics

Neomycin is an antibiotic from the aminoglycoside family. Aminoglycosides have a broad antibacterial spectrum with good activity against Gram negative species, especially *Escherichia coli* and less activity against Gram positive species. This class of antimicrobials has no effect against anaerobic bacteria.

Neomycin binds to the 30S subunit of the bacterial ribosome which disturbs the reading of the constituent code of the RNA messenger, and finally the synthesis bacterial protein. At high concentrations, it has been shown that aminoglycosides damage the cell wall, conferring bactericidal and bacteriostatic properties.

The resistance mechanisms are complex and differ between aminoglycoside molecules. Four mechanisms of resistance have been identified: changes of the ribosome, reduction of permeability, inactivation by enzymes and substitution of the molecular target. The common mechanism of resistance is the production of aminoglycoside modifying enzymes. These resistance mechanisms can be located in mobile genetics elements increasing the likelihood of spread of aminoglycoside resistance as well as co and cross-resistance. The level of resistance of pathogenic *E coli* towards neomycin in calves in Europe ranges between 20 and 50 %.

Indication:

Goats

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in goats.

Cattle (excluding veal calves)

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves).

Swine

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in swine.

Sheep

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in sheep.

Turkeys (growing)

For the control of mortality associated with *E. coli* organisms susceptible to neomycin sulfate in growing turkeys.

Dosage and Administration

Goats

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in goats.

10 milligrams of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram/77mg Neo-vet powder per kilogram) in divided doses for a maximum of 14 days.

Add to drinking water. Not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days. Discontinue treatment 3 days prior to slaughter in goats. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Cattle (excluding veal calves)

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves).

10 milligrams of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram/77mg Neo-vet powder per kilogram) in divided doses for a maximum of 14 days.

Add to drinking water. Not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days. Discontinue treatment 1 day prior to slaughter in cattle. A withdrawal period has not been established for use in pre ruminating calves. Do not use in calves to be processed for veal. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Swine

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in swine.

10 milligrams of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram/77mg Neo-vet powder per kilogram) in divided doses for a maximum of 14 days.

Add to drinking water. Not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days. Discontinue treatment 3 days prior to slaughter in swine. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Sheep

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in sheep.

10 milligrams of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram/77mg Neo-vet powder per kilogram) in divided doses for a maximum of 14 days.

Add to drinking water. Not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days. Discontinue treatment 2 days prior to slaughter in sheep. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Turkeys (growing)

For the control of mortality associated with *E. coli* organisms susceptible to neomycin sulfate in growing turkeys.

10 milligrams of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram/77mg Neo-vet powder per kilogram) for 5 days.

Add to drinking water. Not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 5 consecutive days. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Contraindication:

Do not use in known cases of hypersensitivity to the active substance, to aminoglycosides or to any of the excipients.

Do not use in cases of intestinal obstruction.

Warning and Precautions:

Special warnings for each target species

Medicated drinking water intake can be affected by the severity of the disease. In case of insufficient intake of water, cattle and pigs should be treated parenterally.

Special precautions for use in animals

Powder for oral solution that is to be dissolved in water and cannot be used as it is.

Special care should be taken when considering to administer the product to the newborn calf due to the known higher gastrointestinal absorption of neomycin in neonates. This higher absorption could lead to an increased risk of oto- and nephrotoxicity. The use of the product in neonates should be based on the benefit/risk determination from the attending veterinarian.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria. Official, national and regional antimicrobial policies should be taken into account when the product is used.

Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to neomycin and may decrease the effectiveness of treatment with aminoglycosides due to the potential for cross resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after use.

People with known hypersensitivity to aminoglycosides should avoid contact with the veterinary medicinal product.

If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Other precautions

None.

Interactions with other medications

General anaesthetics and muscle relaxing products increase the neuro-blocking effect of aminoglycosides. This may cause paralysis and apnoea.

Special care should be taken when using concurrently with diuretics and potentially oto- or nephrotoxic substances.

Pregnancy and lactation:

Laboratory studies in animals have not produced any evidence of teratogenic effects of neomycin.

The safety of the veterinary medicinal product has not been evaluated in the target species.

Use only according to the benefit/risk assessment by the responsible veterinarian.

Adverse Effect:

None known

Overdose and Treatment:

Nephrotoxic and/or ototoxic effects may occur in case of accidental overdose.

Withdrawal period:

Cattle excluding veal calves: 1 day before slaughter.

Sheep: 2 days before slaughter.

Swine and goats: 3 days before slaughter.

Turkey: zero days before slaughter.

Proposed Shelf Life

Shelf Life as packaged for Sale :3 years

Shelf Life after first opening of container :7 days

Shelf life after reconstitution or dilution

The solution is discarded after 24 hours

Storage Condition :

Storage below 30°C

Protect from light and moisture.

For veterinary use only

Keep away from children

Jauhi daripada kanak-kanak

BATCH NUMBER
MANUFACTURED DATE
EXPIRY DATE
MAL NUMBER

MARKET AUTHORIZATION HOLDER
FIJILIAM ENTERPRISE SDN.BHD
100 & 101,Taman A.S.T, Jalan Sungai Ujong
70200 Seremban, Negeri Sembilan,
Malaysia

MANUFACTURER:
Nam Pharma Sdn Bhd
5,Lebuh Perusahaan
Klebang 11,Taman
Perindustrian
Antarabangsa IGB
31200,
CHEMOR PERAK
MALAYSIA

Date of Revision: February 2024