

XYLANIL 100MG/ML INJECTION

Description and Composition

XYLANIL 100MG/ML INJECTION is a colourless to pale pink clear solution. Each ml contains 100mg of Xylazine (as hydrochloride) as its active ingredient, 1.8mg Methyl Paraben and 0.2mg Propyl Paraben as its preservatives.

Pharmacodynamics

A potent alpha2-adrenergic agonist, xylazine is classified as a sedative / analgesic with muscle relaxant properties. Although xylazine possesses several of the same pharmacologic actions as morphine, it does not cause CNS excitation in cats, horses or cattle, but causes sedation and CNS depression. In horses, the visceral analgesia produced has been demonstrated to be superior to that produced by meperidine, butorphanol or pentazocine.

Pharmacokinetics

Absorption is rapid following IM injection, but bioavailabilities are incomplete and variable. Bioavailabilities of 40-48% in the horse and 52-90% in the dog have been reported after IM administration.

In horses, the onset of action following IV dosage occurs within 1-2 minutes with a maximum effect 3-10 minutes after injection. The duration of effect is dose dependent but may last for approximately 1.5 hours. The serum half-life after a single dose of xylazine is approximately 50 minutes in the horse and recovery times generally take from 2-3 hours.

In dogs and cats, the onset of action following an IM or SC dose is approximately 10-15 minutes, and 3-5 minutes following an IV dose. The analgesic effects may persist for only 15-30 minutes, but the sedative actions may last for 1-2 hours depending on the dose given. The serum half-life of xylazine in dogs has been reported as averaging 30 minutes. Complete recovery after dosing may take from 2-4 hours in dogs and cats.

Indication

Cats, dogs and horses: To produce sedation, as an analgesic, and as a preanesthetic to local or general anesthesia.

Recommended Dose

Cats, dogs and horses: 1.1mg Xylazine per kg body weight (equivalent to 0.5ml product per 45kg body weight) intravenously or 2.2mg Xylazine per kg body weight (equivalent to 1.0ml product per 45kg body weight) intramuscularly.

Route of Administration

To be given by intravenous or intramuscular injection in horses, dogs and cats.

Contraindication

Xylazine is contraindicated in animals receiving epinephrine or having active ventricular arrhythmias.

Warning and Precautions

Special warnings for each target species

Use with extreme caution in animals with preexisting cardiac dysfunction, hypotension or shock, respiratory dysfunction, severe hepatic or renal insufficiency, preexisting seizure disorders, or if severely debilitated.

May induce premature parturition; it should generally not be used in the last trimester of pregnancy.

Do not use xylazine in conjunction with other tranquilizers.

Avoid intra-arterial injection as it may cause severe seizures and collapse in horses.

Special precautions for use

Special precautions for use in animals:

Not for use in horses intended for food.

Horses have been known to kick after a stimulatory event (usually auditory); use caution. The addition of opioids (e.g., butorphanol) may help temper this effect, but may cause increased risks for hypotension or ileus development.

May inhibit gastrointestinal motility, use with caution in patients treated for intestinal impactions.

Use cautiously in horses during the vasoconstrictive development phase of laminitis as xylazine has been shown to reduce digital flow of blood for about 8 hours after administration.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Horses have been known to kick after a stimulatory event (usually auditory); use caution.

Interaction with Other Medicaments

Do not use xylazine in conjunction with other tranquilizers.

The combination use of acepromazine with xylazine is generally considered safe, but there is potential for additive hypotensive effects and this combination should be used cautiously in animals susceptible to hemodynamic complications.

CNS depressant agents, Other (barbiturates, narcotics, anesthetics, phenothiazines, etc.) may cause additive CNS depression if used with xylazine. Dosages of these agents may need to be reduced.

The use of epinephrine with or without the concurrent use of halothane with xylazine may induce the development of ventricular arrhythmias.

case of a horse developing colic-like clinical signs after reserpine and xylazine has been reported. Until more is known about this potential interaction, use of these two agents together should be avoided.

Pregnancy and Lactation

Limited information was located on the safety of xylazine in pregnancy; apparently, there are no reports of teratogenicity in animals. May induce premature parturition; it should generally not be used in the last trimester of pregnancy.

Side Effects

Emesis is generally seen within 3-5 minutes after xylazine administration in cats and occasionally in dogs.

To prevent aspiration, do not induce further anesthesia until this time has lapsed. Dogs may develop bloat from aerophagia that may require decompression. Because of gaseous distention of the stomach, xylazine's use before radiography can make test interpretation difficult. Large animals may become ataxic following dosing and caution should be observed.

Cats: Emesis, muscle tremors, bradycardia with partial A-V block, reduced respiratory rate, movement in response to sharp auditory stimuli, & increased urination.

Dogs: Muscle tremors, bradycardia with partial A-V block, reduced respiratory rate, movement in response to sharp auditory stimuli, emesis, bloat from aerophagia which may require decompression.

Horses: Muscle tremors, bradycardia with partial A-V block, reduced respiratory rate, movement in response to sharp auditory stimuli, sweating, increased intracranial pressure or decreased mucociliary clearance

Symptoms and Treatment of Overdose

In the event of an accidental overdosage, cardiac arrhythmias, hypotension, and profound CNS and respiratory depression may occur. Seizures have also been reported after overdoses. There has been much interest in using alpha-blocking agents as antidotes or reversal agents to xylazine. Yohimbine, atipamezole, and tolazoline have been suggested for use alone and in combination to reverse the effects of xylazine or speed recovery times. To treat the respiratory depressant effects of xylazine toxicity,

mechanical respiratory support with respiratory stimulants (e.g., doxapram) have been recommended for use.

Storage Condition

Store below 30°C. Protect from light.

Shelf life

3 years.

Following withdrawal of first dose, use the product within 28 days. Any unused material should be discarded.

Packing

20ml, 50ml, 100ml, 250ml.

Revised date: 30.8.2023

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com