

Important information. Please read carefully.

TYMORA

Mouthwash Solution 0.15% w/v

COMPOSITION

Each 15ml contains

Benzydamine Hydrochloride 22.5mg

Methyl Hydroxybenzoate 0.1% w/v

Ethanol 96% 1.18% w/v

PHARMACODYNAMICS

Mechanism of action:

The indazole analogue benzydamine has physicochemical properties and pharmacological activities which differ from those of the aspirin-like NSAIDs. Unlike aspirin-like NSAIDs which are acids or metabolized to acids, benzydamine is a weak base. In further contrast, benzydamine is a weak inhibitor of the prostaglandin synthesis. Only at concentration of 1mM and above benzydamine effectively inhibits cyclooxygenase and lipoxygenase enzyme activity. It mostly exerts its effects through inhibition of the synthesis of pro-inflammatory cytokines including tumor necrosis factor- α (TNF- α) and Interleukin-1 β (IL-1 β) without significantly affecting other pro-inflammatory (IL-6 and 8) or anti-inflammatory cytokines (IL-10, IL-1 receptor antagonist). Further mechanisms of action are hypothesized including the inhibition of the oxidative burst of neutrophils as well as membrane stabilization as demonstrated by the inhibition of granule release from neutrophils and the stabilization of lysosomes. The local anaesthetic activity of the compound has been related to an interaction with cationic channels.

Pharmacodynamic effects:

Benzydamine specifically acts on the local mechanisms of inflammation such as pain, oedema or granuloma. Benzydamine topically applied demonstrates anti-inflammatory activity reducing oedema as well as exudate and granuloma formation. Further, it exhibits analgesic properties if pain is caused by an inflammatory condition and local anaesthetic activity. Hyperthermia, which is indicative of systemic functional involvement, is poorly affected by benzydamine.

PHARMACOKINETICS

Oral doses of benzydamine are well absorbed and plasma drug concentrations reach a peak fairly rapidly and then decline with a half-life of about 13 hours. Less than 20% of the drug is bound to plasma proteins. Although local drug concentrations are relatively large, the systemic absorption of mouthwash-gargle doses of benzydamine is relatively low compared to oral doses. This low absorption should greatly diminish the potential for any systemic drug side-effects when benzydamine is administered by this route. Benzydamine is metabolized primarily by oxidation, conjugation and dealkylation.

INDICATIONS

Tymora Solution is indicated for the temporary relief of painful conditions of the oral cavity including tonsillitis, sore throat, radiation mucositis, aphthous ulcers, post orosurgical and periodontal procedures.

DOSAGE AND ADMINISTRATION

For oromucosal use only. To be used as a gargle or oral rinse.

- Tymora Solution should generally be used undiluted, but if stinging occurs it may be diluted with water.
- The solution should be expelled from the mouth after use.
- Uninterrupted treatment should not exceed seven days.

Dosage in Adult

When used as a gargle, the usual dose is 15ml which should be gargled for at least 30 seconds at 1½ to 3 hourly intervals, as needed.

When used as a rinse for oral lesions, the usual dose is 15ml which should be held in the mouth and swirled around for at least 30 seconds, with repeat use every 1½ to 3 hours throughout the day.

Dosage in Children

5 – 15ml as a gargle if able to do so, or as an oral rinse.

With Impaired Renal Function

Since absorbed benzydamine and its metabolites are excreted in the urine, the possibility of systemic effects should be considered in patients with severe renal impairment.

With Impaired Liver Function

Since absorbed benzydamine is highly metabolised in the liver, the possibility of systemic effects should be considered in patients with severe hepatic impairment.

CONTRAINDICATIONS

Tymora Solution is contra-indicated in patients with known hypersensitivity to the active substance benzydamine hydrochloride and its excipients.

WARNINGS AND PRECAUTIONS

- Benzydamine use is not advisable in patients with hypersensitivity to acetylsalicylic acid or other NSAIDs.
- Caution should be exercised in patients suffering from or with a previous history of bronchial asthma.
- Tymora should generally be used undiluted, but if 'stinging' occurs the rinse may be diluted with water.
- Avoid contact with eyes.
- Tymora contains methyl hydroxybenzoate that may cause allergic reactions (possibly delayed).
- The alcohol (ethanol) content in Tymora is 1.18% w/v which is equivalent to 177 mg per 15ml of solution.
- Tymora is not recommended in children under 6 years of age because lack of sufficient clinical experience.

DRUG INTERACTIONS

None known.

PREGNANCY AND LACTATION

Pregnancy

Benzydamine solution should not be used in pregnancy unless considered essential by the physician.

Breast-feeding

Benzydamine solution should not be used in lactation unless considered essential by the physician.

SIDE EFFECTS

Benzydamine is generally well tolerated and side-effects are minor.

Local Adverse Reactions

The most commonly reported reaction is oral numbness (2.6%). Occasional burning or stinging sensation may occur and has been reported in 1.4% of treated cases. Other local adverse effects were less common and included dryness or thirst (0.2%), tingling (0.2%), warm feeling in mouth and altered sense of taste (<0.1%).

Systemic Adverse Reactions:

These were very uncommon and never of a serious nature. They consisted mainly of nausea, vomiting, retching, gastro-intestinal disorders (0.4%), dizziness (0.1%), headache and drowsiness (<0.1%). Hypersensitivity reactions occur very rarely but may be associated with pruritus, rash, urticaria, photodermatitis and occasionally laryngospasm.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Symptoms:

Benzydamine solution is unlikely to cause adverse systemic effects, even if accidental ingestion should occur. Intoxication is only to be expected if large quantities of benzydamine solution are swallowed (> 300mg).

Symptoms associated with ingested overdose of benzydamine are mainly gastrointestinal symptoms and symptoms of the central nervous system. Most frequent gastrointestinal symptoms are nausea, vomiting, abdominal pain, and esophageal irritation. Symptoms of the central nervous system include dizziness, hallucinations, agitation, anxiety, and irritability.

Management:

In acute overdose only symptomatic treatment is possible. Patients should be kept under close observation and supportive treatment should be given. Adequate hydration must be maintained.

STORAGE CONDITION

Store below 30°C. Protect from light.

SHELF LIFE

The expiry date is indicated on the packaging.

PRODUCT DESCRIPTION, DOSAGE FORM AND PACKAGING AVAILABLE

A clear, green solution with menthol flavour.

Available in pack sizes of 60ml, 100ml and 120ml with measuring cup.

For further information, please consult your physician or pharmacist.

Revision Date: 31-05-2024

Manufacturer and Product Registration Holder:

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5 Cheng Industrial Estate 75250 Melaka, Malaysia.

TYxxxx-001

05.24

Font Size: 10 Times New Roman (for registration purpose)