



SUMMARY OF PRODUCT CHARACTERISTICS
DUTABIT PLUS 0.5/0.4
Dutasteride 0.5mg and Tamsulosin Hydrochloride 0.4mg Hard Capsules
R_p Only

NAME OF THE MEDICINAL PRODUCT

: Dutasteride 0.5mg and Tamsulosin Hydrochloride 0.4mg Hard Capsules

1.TRADE NAME OF PRODUCT

: DUTABIT PLUS 0.5/0.4

STRENGTH:

: 0.5 mg and 0.4 mg.

PHARMACEUTICAL DOSAGE FORM

: Hard Capsules.

2.QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains 0.5 mg of dutasteride and 0.4 mg tamsulosin hydrochloride.
For the full list of excipients, see section 6.1.

3.PHARMACEUTICAL FORM

Size '00', hard shell capsules having opaque orange cap imprinted with 'DTT' in black ink and opaque brown body imprinted with '0.5/0.4' in black ink containing one dutasteride soft gelatin capsule and white to off white Tamsulosin Hydrochloride beadlets.
The product code imprinting details for the Dutasteride and Tamsulosin Hydrochloride Capsules 0.5 mg and 0.4 mg is as below:
Dutasteride and Tamsulosin Hydrochloride Capsules 0.5 mg and 0.4 mg
Cap - 'DTT'
Body – '0.5/0.4'

4.CLINICAL PARTICULARS

4.1 Therapeutic indications

DUTABIT PLUS is indicated as combination therapy for the treatment of moderate to severe symptoms of benign prostatic hyperplasia (BPH).
DUTABIT PLUS reduces the risk of acute urinary retention and the need for surgery in patients with moderate to severe symptoms of BPH.

Dosage/administration

Patients >21 years old (including elderly patients)
The recommended dose of DUTABIT PLUS is one capsule daily (0.5 mg / 0.4 mg).
DUTABIT PLUS must always be taken with a glass of water, up to 30 minutes after a meal. It must never be taken on an empty stomach. Ideally it should be taken at the same time of the day (this means with the same meal).
The capsules should be swallowed whole and not chewed or opened, as contact with the capsule contents may result in irritation of the oropharyngeal mucosa.
Special dosage instructions
Elderly patients:
No age-related dose adjustment is required.
Renal impairment
There is no experience available in patients with renal impairment. DUTABIT PLUS should therefore be used with caution especially in patients with severe renal impairment (creatinine clearance <10ml/min)
Hepatic impairment
There is no experience available in patients with hepatic impairment. DUTABIT PLUS should be used with caution in patients with mild to moderate hepatic impairment. In patients with severe hepatic impairment, the use of DUTABIT PLUS is contraindicated.
Patients ≤21 years old:
DUTABIT PLUS is not indicated in patients ≤21 years old.

4.3 Contraindications

- patients with a history of orthostatic hypotension.
- severe hepatic impairment.
- women, patients ≤21 years old.
- hypersensitivity to tamsulosin, dutasteride, other ingredients of the product or to other 5 α- reductase inhibitors.

4.4 Special Warnings & Precautions for use

Prior to initiating treatment with DUTABIT PLUS the patient should be examined to rule out other causes of the symptoms.
DUTABIT PLUS should only be prescribed after a careful benefit/risk assessment due to the risk of adverse events from both active components, and after careful consideration of alternative treatment options (e.g. monotherapies).
Patients with large residual urinary volume and/or severely impaired urine flow should be monitored carefully for acute or chronic urinary retention.
As the active principle dutasteride is absorbed through the skin, contact with leaking capsules should be avoided. In the event of contact with leaking capsules the area affected should be washed thoroughly with soap and water at once. Women should not handle crushed or broken DUTABIT PLUS capsules if they are pregnant or could become pregnant, due to the possibility of absorption of dutasteride and the possible risk to a male foetus.

Effects on prostate-specific antigen (PSA) and the detection of prostate cancers

Before initiating treatment with DUTABIT PLUS tests to rule out prostate cancer including a digital rectal examination should be carried out. These tests should be repeated at regular intervals during the treatment.
The PSA serum level is an important parameter for early detection of prostate cancer.
Treatment with DUTABIT PLUS leads to an average reduction of 50% in PSA serum levels within 6 months (with great inter-individual fluctuations, standard deviation 30%). This is why PSA levels within the normal range in patients taking DUTABIT PLUS do not rule out prostate cancer.
For this reason, after six months of treatment with DUTABIT PLUS, PSA levels must be assessed again and the result of this measurement used as the baseline value for future measurements. Any confirmed rise in PSA levels compared to the lowest value determined during treatment with DUTABIT PLUS may indicate the presence of prostate cancer (particularly high grade tumour) or lack of compliance and must therefore be evaluated carefully, even if values are still within the normal range for men who have not been treated with 5α-reductase inhibitors. A comparison with previous PSA values should be used to interpret the PSA value in patients treated with dutasteride. After discontinuation of the treatment the PSA levels return to baseline values within 6 months.
The ratio of free to total PSA remains unchanged by DUTABIT PLUS . If the doctor uses the percentage of free PSA as a marker to detect prostate cancer, no adjustment of the value is necessary during treatment with DUTABIT PLUS.

Breast cancer in men

Data on a possible relationship between long-term therapy with dutasteride and the occurrence of breast cancer in men are currently inadequate. Prescribers should instruct their patients to promptly report any changes in their breast tissue such as lumps or nipple discharge. Currently it is not clear if there is a causal relationship between the occurrence of male breast cancer and long term use of dutasteride.

Hypotension

During treatment with an adrenergic alpha-1 receptor blocker such as tamsulosin orthostatic hypotension may occur which in rare cases can result in syncope. Caution is advised in particular in patients who exhibited an excessive haemodynamic reaction during previous treatment with an alpha-1 receptor blocker and in patients receiving antihypertensive therapy. In theory, when Tamsulosin hydrochloride is co-administered with anaesthetics, PDE5 inhibitors or other adrenergic alpha-1 receptor blockers, there is a risk of potentiation of hypotensive effects. DUTABIT PLUS must not therefore be used in combination with other adrenergic alpha-1 receptor blockers, and should only be used with caution in combination with PDE5 inhibitors.
Patients beginning treatment with DUTABIT PLUS should be cautioned to sit or lie down at the first signs of orthostatic hypotension (dizziness) until the symptoms have resolved.

Cardiac failure under combined therapy with tamsulosin

Cases of (acute or chronic) cardiac failure under combined therapy with dutasteride and an alpha blocker (mainly tamsulosin) were observed more frequently than under a monotherapy with dutasteride or an alpha blocker (incidence under dutasteride 0.1%, under alpha blocker 0.2%, under combined therapy 0.6%). However, there was no difference between the three treatment groups for all of the undesirable effects in the cardiovascular system. Up until now there have been no grounds to suspect a causal connection between the medication and the manifestation of cardiac failure, especially since the majority of the patients affected presented predisposing illnesses such as arterial hypertension or coronary heart disease.
In some cases the symptoms of heart failure did not occur until after more than one year of treatment.
Caution is advised in patients with coronary heart disease owing to the possible antihypertensive effects of tamsulosin (see below).

Intraoperative floppy iris syndrome

Intraoperative floppy iris syndrome (IFIS) has been observed during ophthalmological interventions (cataract and glaucoma surgery) in some patients treated with adrenergic alpha-1 receptor blockers, such as tamsulosin. IFIS is characterised by a flaccid iris, progressive intraoperative miosis despite premedication with standard mydriatics, and potential prolapse of the iris towards the incisions for phacoemulsification. IFIS may increase the risk of intraoperative and postoperative ophthalmological complications, and the surgeon should be prepared, if necessary, to adjust his surgical techniques accordingly.
It is recommended that treatment with tamsulosin is not initiated in patients who are about to undergo eye surgery. However, it has not been proven that discontinuation of tamsulosin 1-2 weeks prior to the intervention is beneficial. There have been some reports on IFIS in patients who discontinued tamsulosin some time prior to the operation.

Effects on the reproductive system

The relevance of the effects of dutasteride on semen characteristics for an individual patient's fertility is not known.

The effects of Tamsulosin hydrochloride on sperm count or function have not been examined.

In the case of combined administration of dutasteride and tamsulosin there was an increased incidence of undesirable effects in the organ system "reproductive system and breast", particularly at the start of the treatment (i.e. during the first 6-12 months) compared to therapy with dutasteride or tamsulosin alone.

Hepatic impairment

The effects of hepatic impairment on the pharmacokinetics of dutasteride have not been investigated. As dutasteride undergoes extensive metabolism and has a half- life of 3-5 weeks, DUTABIT PLUS should be used with caution in patients with mild to moderate hepatic impairment (see section 4.2 and 4.3). Since also for tamsulosin there are no data regarding severe hepatic impairment available, DUTABIT PLUS is contraindicated for severe liver impairment.

CYP3A4 inhibitors

Co-administration of tamsulosin and CYP3A4 inhibitors may increase tamsulosin exposure. In particular, there is a risk for significantly increased tamsulosin exposure in the case of CYP2D6 slow metabolisers who are being treated concurrently with potent CYP3A4 inhibitors. As polymorphism for CYP2D6 is generally unknown outside clinical studies, DUTABIT PLUS should basically not be co-administered with potent CYP3A4 inhibitors (e.g. ketoconazole, itraconazole, voriconazole, clarithromycin, indinavir, nelfinavir, ritonavir, saquinavir).

DUTABIT PLUS should only be co-administered with caution with moderate CYP3A4 inhibitors (e.g. erythromycin, fluconazole, verapamil, diltiazem). This applies in particular if the patient is being concurrently treated with a potent or moderate CYP2D6 inhibitor (such as paroxetine) or if concurrent treatment is with a CYP3A4 inhibitor that also inhibits CYP2D6 (such as amiodarone, cimetidine, imatinib).

Renal impairment:

The effects of renal impairment on the pharmacokinetics of tamsulosin have not been investigated in patients with creatinine clearance <10 ml/min. DUTABIT PLUS should therefore be used with caution in such patients. For dutasteride no pharmacokinetic data are available in patients with impaired renal function. Due to the minimal renal elimination, there is no meaningful impact expected.

Swallowing disorders

DUTABIT PLUS has not been studied in patients with swallowing disorders (e.g. stenosing Oesophageal changes or neurological disorder saccompanied by impaired oesophageal motility). Caution is therefore advised in such patients.

Discontinuation of DUTABIT PLUS

On discontinuing DUTABIT PLUS the prostate can go back to the size it was prior to the treatment. Patients should therefore be suitably monitored for the recurrence of symptomatic BPH.

Increased Risk of High-Grade Prostate Cancer

In men aged 50 to 75 years with a prior negative biopsy for prostate cancer and a baseline PSA between 2.5 ng/mL and 10.0 ng/mL taking DUTABIT PLUS in the 4-year Reduction by Dutasteride of Prostate Cancer Events (REDUCE) trial, there was an increased incidence of Gleason score 8-10 prostate cancer compared with men taking placebo DUTABIT PLUS 1.0% versus placebo 0.5%). In a 7-year placebo controlled clinical trial with another 5-alpha reductase inhibitor (finasteride 5 mg, PROSCAR), similar results for Gleason score 8-10 prostate cancer were observed (finasteride 1.8% versus placebo 1.1%).

5-alpha reductase inhibitors may increase the risk of development of high-grade prostate cancer. Whether the effect of 5-alpha reductase inhibitors to reduce prostate volume, or study related factors, impacted the results of these studies has not been established.

Prostate cancer (in particular high grade tumours)

Studies in men at increased risk of prostate cancer revealed a higher incidence of Gleason 8 – 10 prostate cancers in men treated with 5α-reductase inhibitors (dutasteride or finasteride) compared to placebo. The relationship between 5 α-reductase inhibitors and higher grade prostate cancer is not clear yet. Men taking DUTABIT PLUS should be regularly evaluated for the manifestation of prostate cancer, including PSA testing.

Leaking capsules

Dutasteride is absorbed through the skin, therefore, women, children and adolescents must avoid contact with leaking capsules. If contact is made with leaking capsules, the contact area should be washed immediately with soap and water.

Excipients

This medicinal product contains the colouring agent sunset yellow (E-110), which may cause allergic reactions. This medicinal product contains lecithin derived from soya oil. If you are allergic to peanut or soya, do not use this medicinal product.

4.5 Interactions with other medicinal products & other form of interactions

There have been no drug interaction studies for DUTABIT PLUS . The following statements reflect the information available on the individual components.

Pharmacokinetic interactions

Interactions with CYP450 enzyme system

Dutasteride

Dutasteride is metabolized by CYP3A4. A reduction in clearance by concomitant administration of CYP3A4 inhibitors is regarded as clinically irrelevant in view of the wide therapeutic window. In vitro, dutasteride is not metabolised by CYP1A2, CYP2A6, CYP2D6, CYP2E1, CYP2C8, CYP2C9, CYP2C19 and CYP2B6. Under in vitro conditions dutasteride has no inhibitory effect on cytochrome P450 enzymes.

Tamsulosin

Concomitant administration of tamsulosin with CYP3A4 or CYP2D6 enzyme inhibitors May Leadto increased Tamsulosin exposure. Concomitant administration of ketoconazole (a potent CYP3A4 inhibitor) resulted in an increase of the C and AUC max of tamsulosin by a factor of 2.2 and 2.8 respectively.

Concomitant administration of paroxetine (a potent CYP2D6 inhibitor) caused an increase of the C and AUC of tamsulosin by a factor of 1.3 and 1.6 respectively. The max effects of co-administration of both a CYP3A4 and a CYP2D6 inhibitors with tamsulosin have not been evaluated clinically, however there is a potential for significant increase in tamsulosin exposure.

Interactions with oral anticoagulants

Dutasteride

Dutasteride did not show any interaction with the plasma protein binding of warfarin, acenocoumarol or phenprocoumon in vitro.

Tamsulosin

No study has been conducted on interactions between tamsulosin hydrochloride and warfarin. Tamsulosin has no effect on the pharmacokinetics or efficacy of acenocoumarol in healthysubject. The effect of acenocoumarol on the pharmacokinetics of tamsulosin has not been investigated. There are no interaction studies with phenprocoumon. The INR values of patients on oral anticoagulation therapy should be closely monitored for 2-3 months when starting or stopping treatment with DUTABIT PLUS .

Other pharmacokinetic interactions

Dutasteride

No clinically significant pharmacokinetic or pharmacodynamic interactions have been observed between dutasteride and tamsulosin, terazosin, warfarin, digoxin and cholestyramine.

In vitro, dutasteride did not displace diazepam or phenytoin from plasma protein binding and in turn was not displaced by them either.

Tamsulosin

Concomitant administration of tamsulosin hydrochloride and furosemide produced an 11-12% reduction in the C max and AUC of tamsulosin hydrochloride; however these Max changes are clinically irrelevant and no dose adjustment is necessary. Tamsulosin had no effect on the pharmacokinetics of atenolol, digoxin, enalapril, nifedipine or theophylline.

Pharmacodynamic interactions

Tamsulosin

In studies on hypertonic patients whose blood pressure was stable with atenolol, enalapril or nifedipine, tamsulosin did not have a relevant effect on blood pressure compared to placebo.

4.6 Fertility Pregnancy & Lactation

DUTABIT PLUS is contraindicated in women.

No studies have been conducted with DUTABIT PLUS to investigate the effect on pregnancy and lactation. The following statements reflect the information available on the individual components.

Pregnancy

5α-reductase inhibitors can inhibit the development of the external genitalia of male foetuses.

As dutasteride is absorbed through the skin, contact (especially with leaking capsules) should be avoided during pregnancy. It is not known whether a male foetus may be adversely affected if his mother is exposed to the semen of a patient being treated with dutasteride.

Breast-feeding

It is not known whether Dutasteride or Tamsulosin hydrochloride are excreted in human breast milk.

Fertility

Dutasteride has been reported to affect semen characteristics (reduction in sperm count, semen volume, and sperm motility) in healthy men. The possibility of reduced male fertility cannot be excluded.

Effects of Tamsulosin hydrochloride on sperm counts or sperm function have not been evaluated.

4.7. Effects on ability to drive and use machine

The effect of Dutasteride & Tamsulosin hydrochloride capsules on the ability to perform tasks that require judgement, motor or cognitive skills have not been conducted. However, patients should be informed about the possible occurrence of symptoms related to orthostatic hypotension (such as dizziness) and visual disturbances when taking DUTABIT PLUS that could impair the ability to drive and to use machines.

4.8 Undesirable effects

The most common undesirable effects during combined administration of Dutasteride and Tamsulosin hydrochloride

A/s: 280 x 420 mm ■ Black

 AUROBINDO Packaging Development		Product Name	Component	Item Code	Date & Time
		DUTABIT PLUS	Leaflet	P1530903	10.07.2024 & 02.10 pm
		Country	Version No.	Reason Of Issue	Reviewed / Approved by
Team Leader		Kiran	Dimensions (mm)	 30903	
Initiator		Shirisha	280 x 420 mm		
Artist		Advnt (Rakesh)	Pharma Code: 30903		
Additional Information: Supersede Code: P1519885					

related to the organ system "Reproductive system and breast". These undesirable effects were observed more frequently with the combination therapy than with the respective monotherapies and in each case were more frequent in the first year of treatment than in the subsequent years of treatment. The undesirable effects, which were observed either during one of the two monotherapies or in the co-administration study, are listed below by organ system and frequency.

Neoplasms

Unknown : Breast cancer in men

Immune system disorders

Uncommon : Rash,Pruritus,Utricaria

Rare : Angioedema

Ver Rare : Stevens-Johnson syndrome

Psychiatric disorders

common : Decreased libido

Very Rare : Depressive mood

Nervous system disorders

Common : Dizziness

Rare : Syncope

Eyes

Unknown : Blurred vision, visual disturbances

Following market launch there have also been reports of a relationship between the occurrence of what is known as Intraoperative Floppy Iris Syndrome (IFIS) during cataract and glaucoma surgery and treatment tamsulosin.

Cardiovascular disorders

Uncommon : Palpitations, orthostatic hypotension

Unknown : Tachycardia, arrhythmias (e.g. atrial fibrillation)

Respiratory disorders

Uncommon : Rhinitis

Unknown : Epistaxis, dyspnoea

Gastrointestinal disorders

Uncommon : Nausea, constipation, diarrhoea, vomiting, dry mouth

Skin and subcutaneous tissue disorders

Unknown : Exfoliative dermatitis, erythema multiforme

Reproductive system and breast disorders

Common : Impotence, ejaculation disorders (such as retrograde ejaculation), gynaecomastia, breast tenderness.

Very rare : Priapism, testicular pain and testicular swelling

Unknown : Persistent sexual dysfunction (such as impotence, ejaculationdisorders and decreased libido) even after discontinuation of therapy

General disorders

Uncommon : Asthenia

4.9. Overdose

No data are available with regard to overdosage with DUTABIT PLUS . The following statements reflect the information available on the individual components.

Dutasteride

There is no specific antidote for dutasteride. Therefore, if an overdose is suspected appropriate symptomatic treatment should be initiated.

Tamsulosin

Cases of acute overdose with 5mg tamsulosin have been reported. Acute hypotension (systolic blood pressure 70 mmHg), vomiting and diarrhoea were observed in these patients.

In cases of acute hypotension occurring after overdose with tamsulosin hydrochloride cardiovascular support should be given. Normalisation of blood pressure and heart rate may be accomplished by lying the patient down. If this is inadequate, volume expanders and if necessary vasopressors can then be administered. Since tamsulosin exhibits strong plasma protein binding, dialysis is unlikely to be of benefit in removing tamsulosin from the body. Vomiting may be induced to reduce absorption. If a large dose has been taken, administration of activated charcoal and osmotic laxatives may be useful.

5. PHARMACOLOGICAL PROPERTIES

5.1. Pharmacodynamic properties

Pharmacotherapeutic group: Alpha-adrenoreceptor antagonists, ATC code: G04CA52

Mechanism of action:

DUTABIT PLUS is a combination of two drugs with complementary mechanisms of action to improve symptoms in patients with BPH: dutasteride, a dual 5 α - reductase inhibitor (5-ARI) and tamsulosin hydrochloride, an antagonist of α 1a-adrenoreceptors

Dutasteride

Dutasteride inhibits the Type 1 and 2 5 α - reductase isoenzymes which are responsible for the conversion of testosterone to 5 α -dihydrotestosterone (DHT) and therefore causes a fall in the DHT level in the circulation and in the prostate. DHT is the main androgen responsible for hyperplasia of the glandular prostate tissue.

Dutasteride reduces the size of the prostate, relieves the symptoms, improves urine flow and reduces the risk of acute urinary retention and the need for surgery.

Tamsulosin

Tamsulosin is a selective α 1-adrenoreceptor blocker for the symptomatic treatment of functional symptoms of benign prostatic hyperplasia. It binds selectively and competitively to postsynaptic α 1-adrenoreceptors (mainly the alpha1a subtype—approximately 75% of the α 1-receptors in the prostate are of the α 1a subtype) responsible for contraction of the smooth muscle of the prostate and urethra. Tamsulosin thereby reduces smooth muscle tension in the prostate and urethra. This increases maximum urinary flow rate and reduces urinary tract obstruction. It also improves the complex of irritative and obstructive symptoms in which bladder instability and tension of the smooth muscles of the lower urinary tract play an important role.

Effects on DHT/testosterone:

The peak effect of dutasteride in the sense of reduction of DHT is dose-dependent and occurs within 1-2 weeks. After taking dutasteride (0.5 mg/day) for one or two weeks, the mean DHT concentrations in the serum were reduced by 85% and 90% respectively, after one year by 94% and after 2 years by 93%. After discontinuation of treatment the DHT serum concentrations associated with the clinical effects return to the baseline values within about 4 months.

The mean rise in the serum testosterone level was 19% after both one and two years. The testosterone concentrations fluctuated within the normal physiological range

Safety Pharmacodynamics

Adrenergic alpha-1 receptor blockers can reduce blood pressure by lowering peripheral resistance.

5.2. Pharmacokinetic properties

Absorption

Dutasteride

After oral administration the peak serum concentration of dutasteride is attained within 1-3 hours. Absolute bioavailability compared with a 2-hour intravenous infusion is approx. 60%.

Tamsulosin

Tamsulosin hydrochloride is absorbed from the intestine and is almost completely available. After a single dose following a meal, tamsulosin plasma concentrations reach their peak after 6 hours.

Tamsulosin hydrochloride has linear absorption kinetics, with achievement of steady state concentrations by about the fifth day of once-a-day dosing. The rate of absorption of tamsulosin hydrochloride is reduced by a recent meal. Uniformity of absorption can be promoted by the patient always taking tamsulosin hydrochloride 30 minutes after the same meal each day.

Tamsulosin plasma levels exhibit considerable inter-individual variability with both single and multiple dosing.

Distribution

Dutasteride

Dutasteride has a large volume of distribution (300-500 l) and is highly plasma proteinbound (> 99.5%; to albumin and 1-acid glycoprotein). The steady-state serum concentration (C_{ss}) of approx. 40 ng/ml at a dose of 0.5 mg dutasteride a day is C_s achieved after 6 months. After 1 and 3 months the dutasteride serum levels reach 65% and about 90% respectively of the steady-state concentration. The steady-state concentration is also reached in seminal fluid after 6 months.

After administration for 12 months the dutasteride concentration in seminal fluid was on average 3.4 ng/ml (range 0.4-14 ng/ml), i.e. on average 11.5% of the serum level.

Tamsulosin

The volume of distribution of tamsulosin hydrochloride is low (approx. 0.2l/kg). Tamsulosin hydrochloride is extensively bound to human plasma proteins (94% to 99%), primarily -1 acid glycoprotein (AAG).

Metabolism

Dutasteride

Dutasteride undergoes extensive metabolism. It is mainly hydroxylated and dehydrogenated to inactive metabolites. There are 4 major metabolites and 6 subsidiary metabolites. Hydroxylation takes place in vitro via CYP3A4.

Tamsulosin

Tamsulosin hydrochloride is extensively metabolised slowly by cytochrome P450 enzymes in the liver. In vitro results

indicate that CYP3A4 and CYP2D6, and to a minor extent some other CYP isoenzymes, are involved in the metabolism of tamsulosin. However, the majority is present in the plasma in the form of the unchanged active ingredient. None of the metabolites is more active or more toxic than the parent substance. The metabolites of Tamsulosin hydrochloride undergo extensive conjugation to glucuronide or sulphate prior to renal excretion.

Elimination

Dutasteride

5.4% of the dose administered is excreted as unchanged dutasteride in the faeces, and the rest in the form of metabolites. Only traces of the unchanged substance (less than 0.1% of the dose) can be detected in the urine. Dutasteride clearance is low. The elimination half-life is 3-5 weeks.

Tamsulosin

Following oral administration of tamsulosin in the form of modified release capsules, the apparent elimination half life in the fed state after a single dose is approximately 10 hours and in the steady state is approximately 13 hours. Approximately 10% of the substance is excreted unchanged in urine.

Kinetics of special patient populations

No pharmacokinetic studies have been conducted with DUTABIT PLUS on special patient populations. The following statements reflect the information available on the individual components.

Elderly

Dutasteride

In the age groups >50-65 years and >70 years there were no statistically significant differences in pharmacokinetics.

Tamsulosin

Findings indicate that overall exposure is approx. 40% increased in patients aged 55-75 years compared with young males.

Renal impairment

Dutasteride

The effects of renal impairment on the pharmacokinetics of dutasteride have not been investigated. As less than 0.1% of the dose is excreted in the urine, no effect on the serum levels is to be expected.

Tamsulosin

No clinically relevant changes in the pharmacokinetics of tamsulosin were observed in patients with mild to moderate (30 CLcr < 70 mL/min/1.73m²) or moderate to severe Lcr(10 CLcr < 30 mL/min/1.73m²) renal impairment compared with subjects with normal Lcr renal function. However, patients with terminal renal disease (CLcr < 10 mL/min/1.73m²) Cr have not been studied.

Hepatic impairment

Dutasteride

The effects of hepatic impairment on the pharmacokinetics of dutasteride have not been investigated.

Tamsulosin

In patients with mild to moderate hepatic dysfunction (Child-Pugh classification: Grades A and B), no clinically relevant changes in the pharmacokinetics of tamsulosin hydrochloride were observed. Tamsulosin hydrochloride has not been studied in patients with severe hepatic dysfunction (Child-Pugh C).

6.PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Composition of Hard Capsule shell

In Cap:

Hypermellose

Carrageenan

Potassium chloride

Titanium dioxide (E171)

Sunset yellow FCF (FD&C Yellow 6) (E 110)

Iron oxide red (E172)

In Body:

Hypermellose

Carrageenan

Potassium chloride

Titanium dioxide (E171)

Iron oxide black (E172)

Iron oxide red (E172)

Contents in Dutasteride soft capsules:

Glycerol monocaprylocaprate (Type-I)

Butylhydroxytoluene (E321)

Soft capsule shell:

Gelatin (160 bloom)

Glycerol

Titanium dioxide (E171)

Iron oxide yellow (E172)

Contents in Tamsulosin Hydrochloride Beadlets

Cellulose, Microcrystalline (Grade-101)

Talc

Methacrylic acid-ethyl acrylate copolymer (1:1) dispersion 30 percent (contains 0.7% Sodium Laurilsulfate and 2.3% Polysorbate 80)

Triacetin

Calcium Stearate

Printing ink (Black)

Shellac (E904)

Black iron oxide (E172)

Potassium hydroxide

6.2Incompatibilities

Not applicable.

6.3Shelf life

Please refer outer package for expiry date.

6.4 Special precautions for storage

Do not store above 30°C.

6.5 Nature and contents of container

Blister packs: 30 hard capsules (3 x 10's blister).

6.6 Special precautions for disposal and other handling

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7.MANUFACTURER & PRODUCT REGISTRATION HOLDER:

Manufacturer:

Aurobindo Pharma Limited,

Unit VII, Special Economic Zone Tsiiic,

Plot No SJ Sy Nos 411/P 425/P 434/P 435/P And 458/P

Green Industrial Park, Jadcherla Manda/ Mahabubnagar District,

Polepally Village, 509302, India

Product Registration Holder in Malaysia:

Healol Pharmaceuticals Sdn Bhd,

74-3, Jalan Wangsa Delima 6,

KLSC Wangsa Maju, 53300,

Kuala Lumpur,

Malaysia.

8.DATE OF REVISION OF THE TEXT

08 July 2024.