

Important information. Please read carefully.

TYMORA

Oromucosal Spray 0.3% w/v

COMPOSITION

Contains 3mg Benzydamine Hydrochloride/ml. Each bottle (15ml) contains 88 actuations. The volume/actuation is 0.17ml, with dose/actuation of 0.51mg Benzydamine Hydrochloride. Methyl Parahydroxybenzoate 0.5mg/ml as preservative. Ethanol 96% 81mg/ml.

PHARMACODYNAMICS

Benzydamine specifically acts on the local mechanisms of inflammation such as pain, oedema or granuloma. Benzydamine topically applied demonstrates anti-inflammatory activity reducing oedema as well as exudate and granuloma formation. In addition, it exhibits analgesic properties if pain is caused by an inflammatory condition and local anaesthetic activity. Hyperthermia, which is indicative of systemic functional involvement, is poorly affected by benzydamine.

PHARMACOKINETICS

Benzydamine is rapidly absorbed from the gastrointestinal tract and maximum plasma levels reached after 2-4 hours upon oral administration. The most important aspect of the tissue distribution of Benzydamine is its tendency to concentrate at the site of inflammation. About half of the Benzydamine is excreted unchanged via the kidney at a rate of 10% of the dose within the first 24 hours. The remainder is metabolised, mostly to N-Oxide.

INDICATIONS

Tymora Oromucosal Spray 0.3% W/V is indicated for the temporary relief of painful conditions of the mouth and throat including tonsillitis, sore throat, radiation mucositis, aphthous ulcers, pharyngitis, swelling, redness, inflammatory conditions, post orosurgical and periodontal procedures.

DOSAGE AND ADMINISTRATION

For oromucosal use only.

Adults and children over 12 years: 2-4 sprays (1-2 mg) directly onto the sore/inflamed area and swallow gently. Repeat every 1 1/2 to 3 hours as necessary.

Children (6-12 years): 2 sprays (1 mg) directly onto sore/inflamed area and swallow gently. Repeat every 1 1/2 to 3 hours as necessary.

Children under 6 years: Not recommended.

Uninterrupted treatment should not exceed seven days, unless under medical supervision.

Instructions for use

1. To use the spray, hold the bottle in an upright position.
2. The first time you use the spray, take the transparent cup off. Put the cup with spray arm on the bottle. Point the spray arm away from you and anyone else. Then press the white plunger down firmly several times until a fine spray appears out of the end of the spray arm. Your spray is now ready to use.
3. Point the spray arm at the sore part of your throat or mouth and press the white plunger down firmly. One press releases one spray of medicine.
4. Apply the correct number of sprays according to the instruction above. Then wipe the end of the spray arm with a clean tissue. This helps to prevent your spray blocking.

5. If your spray blocks or stops working properly, return it to your pharmacist. Do not push anything into the end of the spray arm to try to clear a blockage.
6. No additional cleaning is required in use of the product.

CONTRAINDICATIONS

Tymora Oromucosal Spray 0.3% W/V is contra-indicated in patients with known hypersensitivity to the active substance benzydamine hydrochloride or to other components/excipients (either one).

WARNINGS AND PRECAUTIONS

If one of the indicated adverse effects occurs, application of the medication must be suspended.

Benzydamine use is not advisable in patient with hypersensitivity to salicylates (e.g. acetylsalicylic acid and salicylic acid) or other NSAIDs.

Bronchospasm may be precipitated in patients suffering from or with a previous history of bronchial asthma. Caution should be exercised in these patients.

Avoid contact with eyes.

If the condition is aggravated or not improved use should cease.

Methyl parahydroxybenzoate may cause allergic reactions (possibly delayed).

This medicine contains 13mg of alcohol (ethanol) in each spray. The amount in 1 spray of this medicine is equivalent to less than 1ml beer or 1ml wine. The small amount of alcohol in this medicine will not have any noticeable effects.

This medicine contains less than 1 mmol sodium (23mg) per dose of 4 sprays, that is say essentially "sodium free"

DRUG INTERACTIONS

There are no known drug interactions with benzydamine

PREGNANCY AND LACTATION

Pregnancy

Tymora Oromucosal Spray 0.3% W/V should not be used in pregnancy unless considered essential by the physician. There is no evidence of a teratogenic effect in animal studies.

Breast-feeding

Tymora Oromucosal Spray 0.3% W/V should not be used in lactation unless considered essential by the physician.

SIDE EFFECTS

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness

The following rate values have been used: Very common ($\geq 1/10$), Common ($\geq 1/100$ to $<1/10$), Uncommon ($\geq 1/1,000$ to $<1/100$), Rare ($\geq 1/10,000$ to $<1/1,000$) and Very rare ($<1/10,000$), not known (cannot be estimated from the available data).

The most common side effects are numbness and a stinging feeling in the mouth.

System organ class	Frequency	Adverse reaction
Immune system disorders	Not known	Anaphylactic reactions, Hypersensitivity reactions.
Respiratory, thoracic and mediastinal disorders	Very rare	Laryngospasm or bronchospasm
Gastrointestinal disorders	Uncommon	Oral numbness and a stinging feeling in the mouth. The stinging has been reported to disappear upon continuation of the of the treatment, however if it persists it is recommended that treatment be discontinued.
Skin and subcutaneous tissue disorders	Very rare	Pruritus, urticaria, photosensitivity reaction and rash
	Not known	Angioedema

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Symptoms

Tymora Oromucosal Spray 0.3% W/V is unlikely to cause adverse systemic effects, even if accidental ingestion should occur. No special measures are required. Intoxication is only expected in case of accidental ingestion of large quantities of benzydamine (> 300 mg)

Symptoms associated with overdose of ingested benzydamine are mainly gastrointestinal symptoms and symptoms of the central nervous system. Most frequent gastrointestinal symptoms are nausea, vomiting, abdominal pain and oesophageal irritation. Symptoms of the central nervous system include dizziness, hallucinations, agitation, anxiety and irritability.

Treatment

In acute overdose only symptomatic treatment is possible. Patients should be kept under close observation and supportive treatment should be given. Adequate hydration must be maintained.

STORAGE CONDITION

Store below 30°C.

SHELF LIFE

The expiry date is indicated on the packaging.
Discard 6 months after first opening.

PRODUCT DESCRIPTION, DOSAGE FORM AND PACKAGING AVAILABLE

Oromucosal spray solution.

Clear and colourless solution with cherry taste and flavour in HPDE bottle equipped with mechanical dosing pump.

Available in 15ml (88 actuations) multidose spray bottle with a dosing pump.

For further information, please consult your physician or pharmacist.

Revision Date: 13-05- 2024

Manufactured by:
Laboratorium Sanitatis, S.L.
Parque Tecnológico de Álava,
C/Leonardo da Vinci 11,
01510 Miñano (Álava), Spain

Product Registration Holder:
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate 75250 Melaka, Malaysia.

TYxxxx-001

05.24

Font size: Times New Roman 11 (for registration purpose)