

STORAGE CONDITION

Store below 30°C. Protect from moisture.

PRODUCT DESCRIPTION

White effervescent granules with lemon lime flavor, produce clear to slightly cloudy solution when dissolved in water.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK**

For further information, please consult your pharmacist or physician.

Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn. Bhd.

1-5 Cheng Industrial Estate 75250 Melaka, Malaysia.

Contract Manufactured by:

Unison Nutraceuticals Sdn. Bhd.

No. 13, Jalan Tasik Utama 52, Kawasan Perindustrian Tasik Utama,
75450 Ayer Keroh, Melaka, Malaysia.

Revision Date: 13-Aug-2025

001

giomie**Utinol**

EFFERVESCENT GRANULES

Urinary Alkalinizer

Composition:

Each sachet contains:

Sodium Bicarbonate	1.76g
Sodium Citrate	0.63g
Citric Acid	0.72g
Tartaric Acid	0.89g

PHARMACODYNAMICS

Sodium Bicarbonate acts as alkalinizing agent which increases the excretion of free bicarbonate ions in the urine, and thus raises the urinary pH. A rise in urinary pH increases the solubility of cysteine in the urine. By maintaining alkaline urine, the actual dissolution of uric acid stones may be accomplished. Sodium Bicarbonates acts as antacid as well by neutralizing or buffering existing quantities of stomach acid but has no direct effect in its output.

Sodium Citrate is metabolized to bicarbonates after absorption, thus providing similar pharmacological effect as per Sodium Bicarbonate.

Citric Acid and Tartaric Acid both having transient effect as neutralizing or buffering agents.

PHARMACOKINETICS

After absorption, Sodium Citrate is metabolized to bicarbonates. Sodium Bicarbonate is excreted through kidney and also via lung by forming carbon dioxide. Sodium Citrate, Citric Acid and Tartaric Acid are excreted through urine. Less than 5% of the citrates are excreted in the urine unchanged.

INDICATIONS

Utinol is indicated for the relieve of discomfort caused by mild urinary tract infections (UTI), in which provides symptomatic relief of dysuria. It may enhance the action of certain antibiotics especially sulphonamide. It helps to prevent crystallization of uric acid (urates) in gout therapy and promoting dissolution of uric acid stones (kidney stones). This alkalinizer can be used for relieving symptoms associated with gastric hyperacidity.

DOSAGE AND ADMINISTRATION

For oral use only.

Dissolve 1-2 sachets in a glass (200ml) of water.

Four times daily or as prescribed.

CONTRAINDICATIONS

Contraindicated in patients with severe renal impairment or renal failure, hypernatremia, metabolic or respiratory alkalosis, hypocalcaemia, hypochlorhydria, Addison's disease, acute dehydration, heat cramps, overt and occult cardiac failure, and severe myocardial damage.

Not recommended in concomitant use with hexamine mandelate or hexamine hippurate therapy because an acidic urine is needed.

Avoid concomitant use with quinolone antibiotics as to prevent crystalluria in alkaline urine.

WARNING AND PRECAUTIONS

Should not be taken by patients on a sodium restricted diet.

Caution should be taken in patients with peptic ulceration and/or renal abnormalities as to avoid the condition of metabolic alkalosis. Should be given extremely cautiously to patients with heart failure, oedema, renal impairment, hypertension, eclampsia, and or aldosteronism.

DRUG INTERACTIONS

Use of urinary alkalinizer theoretically may result in decreased therapeutic effect of chlorpropamide, lithium, salicylates and tetracyclines; increased therapeutic effect of amphetamines, ephedrine, and pseudoephedrine.

Antacid: Concurrent use of antacids with Sodium Citrate and Sodium Bicarbonate may promote development of calcium stones in patients with uric acid stones and may also cause hypernatremia. Concurrent use of aluminium containing antacids with salts can increase aluminium absorption, possibly resulting in acute aluminium toxicity, especially in patients with renal insufficiency.

Quinolones: Citrates may reduce the solubility of ciprofloxacin, norfloxacin, or ofloxacin in the urine. Patients should be observed for signs of crystalluria and nephrotoxicity.

PREGNANCY AND LACTATION

Use in pregnancy and during lactation should be considered only when the possible benefits outweigh the potential risks.

SIDE EFFECTS

Sodium Bicarbonate given orally can cause stomach cramps, belching, and flatulence. Rarely, spontaneous rupture of the stomach has been reported.

Tartrate component of this product may not be completely absorbed. Due to this, it may exert mild laxative effect. Prolonged and excessive use may cause a systemic alkalosis and or hypernatremia.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Excessive use of bicarbonate or bicarbonate-forming compounds may lead to hypokalaemia and metabolic alkalosis, especially in patients with impaired renal function. Symptoms include mood changes, tiredness, slow breathing, muscle weakness, and irregular heart-beat. Muscle hypertonicity, twitching, and tetany may develop, especially in hypocalcaemic patients.

Treatment of metabolic alkalosis associated with bicarbonate overdose consists mainly of appropriate correction of fluid and electrolyte balance. Replacement of calcium, chloride, and potassium ions may be of particular importance.

SHELF-LIFE

The expiry date is indicated on the packaging.