

ESmonen 20% W/W Premix Powder

KEEP OUT OF REACH OF CHILDREN/ JAUHKAN DARI CAPAIAN KANAK-KANAK

Controlled Medicine/ Ubat Terkawal

Product name: ESmonen 20% w/w Premix Powder

Active Ingredients: 100g:20g monensin (20,000,000U)

Dosage form: powder

Product description: Brown or dark brown, Powder

Pharmacodynamics: Monensin is a monovalent ionophore anticoccidial drug with a broad-spectrum anticoccidial effect. Its mechanism of killing coccidia is by interfering with the normal penetration of K⁺ and Na⁺ in the insect cells, so that a large amount of Na⁺ enters the cells. In order to balance the osmotic pressure, a large amount of water enters the coccidial cells, causing swelling and death. It is effective against coccidiosis in chickens (*Eimeria acervulina*, *E.necatrix*, *E.tenella*, *E.maxima*, *E.mivati*, *E.acervulina*, *E.brunetti*). The peak period of action of monensin is in the first 2 days of the life cycle of coccidia. It has an inhibitory effect on sporozoites and first generation schizonts. The best effect is the second day after coccidia infection. In addition, monensin also has a strong antibacterial effect on gram-positive bacteria such as *Staphylococcus aureus*, *Streptococcus* and *Clostridium perfringens*.

Pharmacokinetics: Monensin is seldom absorbed in the intestines of normal cattle and poultry, and the drugs and metabolites excreted through feces account for about 99% of the administered dose. Metabolism tests of monensin in cattle and rats show that monensin can be metabolized in tissues into a variety of metabolites after a small amount of monensin is absorbed. The metabolites excreted in feces are the same as liver metabolites.

Indication: As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. mitis*, *E. necatrix*, *E. maxima*, *E. tenella*, and *E. brunetti* in broiler chickens

Target species: Broiler chickens

Recommended dose: Thoroughly mix 500 g (0.50 kg) of ESmonen in one metric tonne (1000 kg) of feed to provide monensin at 100 g monensin activity per metric tonne of feed.

Route of Administration: Must be thoroughly mixed in feeds before use. Recommended that ESmonen be mixed with a small quantity of one of the feed ingredients before it is incorporated in the finished feed.

Contraindication: 1. Do not feed to replacement and laying chickens or turkeys. 2. Poultry consuming monensin should not be treated with products containing tiamulin.

Warning: When mixing and handling Monensin premix, use protective clothing, impervious gloves and a dust mask.

Operators should wash thoroughly with soap and water after handling.

Cautions: 1. Do not use Monensin premix for treatment of outbreaks of coccidiosis. 2. Do not exceed recommended levels as reduced average daily gains may result. 3. Do not allow canines, horses or other equines, or guinea fowl access to formulations containing monensin. Ingestion of monensin by these species has been fatal. 4. Some species of turkey coccidia may be monensin tolerant. 5. Not to be used after expiration date.

Interaction of Other Medicaments: May be used in feeds containing the pellet binding agents bentonite (2%), Attapulgate (2%), Kaolin (2.5%), Lignin Sulfonate (4%), Carboxy Methylcellulose (0.1%) or Agri-Colloid.

Pregnancy and Lactation: Do not feed to laying chickens or turkeys

Sides effects: None known

Symptoms and treatment of overdose: None

Storage condition: Store in a cool, dry place below 30°C.

Shelf life:2 years

Shelf life after first opening of container:3 months

Shelf life after reconstitution or dilution:3 months

ATCVet Code: QP51BB03

Withdraw period: No pre-slaughter withdrawal is required when this drug product is used according to the label directions.

Maximum Residual : Residue label: monensin. Chicken:Muscle 10µg/kg, Fat 100µg/kg, Kidney 10µg/kg, Liver 10 µg/kg

Disposal containers instructions: Collect and reclaim or dispose in sealed containers at licensed waste disposal site. Dispose of contents/container in accordance with local/regional/national/international regulations. Review federal, provincial, and local government requirements prior to disposal.

Pack size: 25 kgs

Batch Number:

Manufacturing Date:

Expiration Date:

**Manufactured by:****Zhejiang Esigma Biotechnology Co.,Ltd.**

Address:No. 3 Chunchao West Road, Changan Town,Haining City, Zhejiang, China

**Product registration holder:****Pahang Pharmacy SDN BHD (10826-K)**

Address: Lot 5979 Jalan Teratai, 5 1/2 Mile Off Jalan Meru, 41050 Klang, Selangor Darul Ehsan, Malaysia

Date of revision of package insert: 22 Aug. 2024