

PRODUCT LITERATURE

Product Name:

PVP-WORMFREE SUSPENSION 250MG/5ML

Each 5 ml contains:

Active Ingredient:

Pyrantel Pamoate 725 mg (equivalent to Pyrantel base USP 250 mg)

Preservative:

Propyl Paraben 0.01%w/v
Methyl Paraben 0.1%w/v

Product Description:

A yellow coloured suspension

Target Species:

Dogs

Pharmacodynamics:

Pyrantel pamoate may be used for the treatment of hookworms and roundworms in humans and pets with effects that can vary according to parasite species and resistance status. Pyrantel is most commonly used in horses for ascarids, large and small strongyles and oxyurids (and at a double dose for Anoplocephala). In dogs and cats, pyrantel is used for common GI nematodes, except whipworms.

Pharmacokinetics:

Because pyrantel pamoate is poorly absorbed from the GI tract, its antiparasitic effects remain active in the lower GI tract of dogs, horses and cats.

Indication:

To prohibit the reinfections of *Toxocara canis*.

Recommended Dose:

Equivalent of 2.27 milligrams of pyrantel base per pound of body weight.

Administer 1 teaspoon (5 mL) for each 4.5kg of body weight.

- a) Administer to puppies at 2, 3, 4, 6, 8, and 10 weeks of age.
- b) Administer to lactating bitches 2 to 3 weeks after whelping.
- c) Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals.

Consult your Veterinarian for diagnosis and treatment.

Route of Administration: Oral

Contraindications:

Do not use in cases of known hypersensitivity to the active substances or to any of the excipients and simultaneously with piperazine and levamisole.

Warning & Precautions:

Pyrantel Pamoate should not be used in pets that are allergic to it or pets that are severely weakened or frail. It is considered safe to use in nursing animals and should be used with caution (follow the dosing instructions carefully) in pregnant animals.

Interactions with Other Medicaments:

Pyrantel should not be used concurrently with morantel or levamisole because of the similarity mechanisms of action and toxicity.

Pregnancy & Lactation:

Pyrantel Pamoate is safe to use in pregnancy and lactation and is frequently used in these situations to minimize the worm burdens of neonatal animals.

Side effects:

With recommended dosages, reports of side/adverse effects in association with administration of the tetrahydropyrimidines are rare.

Symptoms & Treatment of Overdose:

Pyrantel has a moderate margin of safety. Dosages up to, approximately 7 times recommended generally result in no toxic reactions. In horses, doses of 20 times those recommended yielded no adverse effects. Chronic dosing of pyrantel pamoate in dogs resulted in symptoms when given at 50 mg/kg/day, but not at 20 mg/kg/day over 3 months. Symptoms of toxicity that could possibly be seen include increased respiratory rates, profuse sweating (in animals able to do so), ataxia or other cholinergic effects.

Storage Conditions:

Store below 30°C in a dry place and protected from light.

Dosage Form:

Oral Solution/Suspension

Packaging:

Packed in 60ml and 1Litre in White HDPE Plastic bottle.

Disposal of Container:

To be disposed as per local authority requirement.

Manufactured & Distributed by:

Pelangi Vet Pharma Sdn Bhd. (665097K)

Lot 8, PERDA Industrial Park,
Lorong IKS, Simpang Ampat B
14100 Simpang Ampat,
S.P.S Pulau Pinang, Malaysia

Date of revision: 21/06/2024