

VK05UM-240730
VK King Kong Balm Insert
Size: 84mm(L) x 150mm(H)

84mm



REG. NO.: MALxxxxxxxX

ACTIVE INGREDIENT: Camphor 11%w/w, Menthol 8%w/w, Methyl Salicylate 5%w/w

PRODUCT DESCRIPTION
White colour ointment with characteristic smell

INDICATION: Provide symptomatic relief for mild to moderate muscular and joint aches and pains, muscle cramps, shoulder aches and stiff neck.

DIRECTION OF USE
Rub on affected area 2 to 3 times a day.

ROUTE OF ADMINISTRATION: Topical

PHARMACODYNAMICS
Camphor: Camphor is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol, methyl salicylate, it becomes ideal for neuralgia and other painful areas.

Menthol: Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl Salicylate: Methyl salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins and cause platelet dysfunction. Methyl salicylate is used topically as a counter irritant. Upon application, it is absorbed through the skin and is applied for the relief of the pain in rheumatic conditions and painful muscle or joints.

PHARMACOKINETICS
Camphor is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure. It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine. Camphor crosses the placenta.

After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl Salicylate may be absorbed through intact skin. The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity. Absorption characteristics of salicylates vary with the dose, formulation and route of administration. Percutaneous absorption is enhanced by exercise, heat, occlusion or disruption of the integrity of the skin or application to large areas of skin. Both the rate and extent of absorption increases after repeated application; increasing the bioavailability. Methyl salicylate is extensively metabolised to salicylic acid in the dermal and subcutaneous tissues after topical application. At therapeutic levels, the half-life of salicylates is 2- 4 hours. As salicylate level increase into the toxic range, the half-life can be greater than 18 hours.

CONTRAINDICATIONS: Do not use in patients with aspirin or salicylate idiosyncrasy. Patient with known hypersensitivity to any of the ingredients in the product.

WARNINGS AND PRECAUTIONS: This product is contraindicated in children below 2 years of age. Caution must be exercised when older children are treated.

It is dangerous to place any camphor containing product into the nostril of children. A small amount applied this way may cause immediate collapse.

Initially apply a small amount to a limited skin area and

observe whether there is discomfort or itch before applying to larger skin area. Discontinues use if irritation develops. Avoid contact with the eyes and mucous membranes. Do not apply to open wounds or damaged skin. Avoid excessive use during pregnancy and lactation. Do not apply over large areas of the body or bandage or use with heating pads. If symptoms persist, consult your doctor.

For external use only. Keep out of reach of children. *Jauhkan daripada capaian kanak-kanak.*

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES: Not known

INTERACTIONS WITH OTHER MEDICAMENTS: This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joints pains may increase the chances of bleeding.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION: Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

ADVERSE EFFECTS/ UNDESIRABLE EFFECTS: This product can cause convulsions. Headache, dizziness, nausea and vomiting, skin irritation, contact dermatitis, rash, itching, redness or swelling, burning or stinging sensation may occur.

May cause hypersensitivity/ allergic reactions in some individuals with sensitive skin.

SYMPTOMS AND TREATMENT OF OVERDOSE:
Symptoms of methyl salicylate overdose
Salicylate intoxication can occur after ingestion or topical application of methyl salicylate. Mild chronic salicylate intoxication or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage. Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.

Symptoms of camphor/ menthol overdose
Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness and coma; instant collapse in infants after the local application of menthol or camphor to their nostrils. Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self- limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema and status epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure and death.

Treatment
The stomach should be emptied by gastric lavage or administration of oral activated charcoal. Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalemia and dehydration if present. Any convulsion must be controlled first through supportive care including anticonvulsant therapy.

STORAGE CONDITIONS: Store below 30°C. Protect from direct sunlight.

SHELF LIFE: Please refer to the packaging. Do not use beyond the expiry date.

DOSAGE FORM: Ointment

PACKING: 10g, 15g, 20g, 25g, 30g

150mm

FRONT

84mm



DESKRIPSI PRODUK

Balsem berwarna putih dengan bau ramuan khas.

INDIKASI
Memberi kelegaan kepada sakit-sakit otot dan sendi, kekejangan otot, sakit bahu dan sakit tengkuk.

CARA PENGGUNAAN
Sapukan pada kawasan yang terjejas 2 hingga 3 kali sehari.

ARAHAN PENGGUNAAN: Topikal

KONTRAINDIKASI: Jangan gunakan pada pesakit yang menggunakan aspirin atau salicylate idiosyncrasy dan pengguna yang mempunyai rangsangan hipersensitiviti terhadap mana-mana bahan ramuan dalam produk ini.

AMARAN DAN LANGKAH BERJAGA-JAGA: Balsem ini tidak sesuai untuk kanak-kanak di bawah umur 2 tahun. Perhatian hendaklah diambil untuk penggunaan kanak-kanak yang berumur 2 tahun ke atas.

Produk ini mengandungi camphor dan adalah merbahaya kepada kanak-kanak sekiranya disapukan ke dalam bahagian hidung. Walaupun hanya sedikit, cara sapuan ini mungkin akan menyebabkan kanak-kanak pengsan serta-merta.

Mungkin akan memberi rangsangan pada kulit bagi pengguna yang berkulit sensitif. Cubu sapukan sedikit di bahagian kulit terhad, sebelum disapukan pada bahagian kulit yang meluas. Hentikan penggunaannya jika mengalami gatal-gatal yang serius dan elakkan

sentuhan pada mata, selaput lendir atau luka. Hadkan penggunaan untuk ibu yang mengandung atau ibu yang menyusukan anak. Jangan sapukan pada permukaan badan yang luas, dibalut atau bahagian badan yang menggunakan pad pemanas. Sekiranya simptom berterusan, sila rujuk kepada doktor.

Untuk penggunaan luar sahaja.
Jauhkan daripada capaian kanak-kanak.

KESAN TERHADAP PEMANDU KENDERAAN DAN PENGOPERASI MESIN: Tidak diketahui

INTERAKSI DENGAN UBAT LAIN: Produk ini mengandungi methyl salicylate dan apabila disapu pada kulit, ia akan menyerap ke dalam saluran darah melalui kulit. Bagi pesakit yang menggunakan warfarin, penggunaan berlebihan pada otot-otot atau sendi mungkin akan meningkatkan peluang pendarahan.

KENYATAAN PENGGUNAAN PRODUK SEMASA KEHAMILAN DAN PENYUSUAN: Untuk ibu yang mengandung atau ibu yang menyusukan anak, sila rujuk kepada doktor sebelum menggunakannya.

KESAN SAMPINGAN/ KESAN TIDAK DIINGINI: Produk ini mungkin menyebabkan kekejangan, sakit kepala, pening, loya dan muntah, kerengsaan kulit, dermatitis kontak, ruam, gatal, kemerahan atau bengkak, sensasi terbakar atau menyengat mungkin berlaku.

Boleh menyebabkan hipersensitiviti/ reaksi alergic pada beberapa individu dengan kulit sensitif.

CARA PENYIMPANAN: Simpan di tempat kering pada suhu di bawah 30°C dan daripada sinaran cahaya matahari.

JANGKA HAYAT: Sila rujuk kepada label luaran. Jangan gunakan selepas tarikh luput.

PEMBUNGUHAN: 10g, 15g, 20g, 25g, 30g

150mm

金刚牌万金油

产品描述：白色膏体，具有特征性的气味。

主治：帮助缓解轻至中度的肌肉和关节疼痛、肌肉痉挛、肩痛和颈部僵硬的症状。

使用方法：涂抹于患处，一天2至3次。

使用说明：局部使用

禁忌：对阿司匹林或水杨酸盐过敏的患者禁用。对产品中任何成分已知过敏的患者禁用。

警告和注意事项：本产品不适用于2岁以下的儿童。在治疗年龄较大的儿童时必须小心。

将含有樟脑的产品放入儿童的鼻孔内是危险的。这种使用方式，即使是少量的樟脑都可能会导致儿童立即崩溃。

在大范围涂抹皮肤之前，先在小范围的皮肤上涂抹少量本产品进行测试，观察是否有不适或瘙痒。如果出现不良反应，则停止使用。避免与眼睛和黏膜接触。不要涂抹在伤口或受损的皮肤上。怀孕和哺乳期间避免

Product Registration Holder:
Jasei Healthcare Sdn Bhd
658-D, Jalan Bukit Melaka 1/1,
Taman Bukit Melaka, Bukit Beruang,
Mukim Bukit Baru, 75450 Melaka Tengah,
Melaka, Malaysia.

Manufacturer:
Wellmetech Pharmaceutical Industries Sdn Bhd
Lot B058, Kota Samarahan Industrial Park,
94300 Samarahan, Kuching, Malaysia.

Date of Revision: July 2024

REAR