



GUTRIL-S 500MG CAPSULE
GLUCOSAMINE SULPHATE

Active Ingredient:

Each capsule contains:

Glucosamine Sulphate Sodium Chloride 628mg (equivalent to Glucosamine Sulphate 500mg)

Derived from seafood.

Description: White colour powder is filled in green size 0 bovine capsule.
No marking on the capsule.

Pharmacodynamic:

Glucosamine can stimulate the production of cartilage- building compounds, restore damaged cells of cartilage tissue and has some anti-inflammatory abilities.

Pharmacokinetic:

Glucosamine

Absorption

After oral administration, bioavailability is low due to first-pass hepatic metabolism 26%. The gastrointestinal absorption is close to 90%.

Distribution

Glucosamine is not protein-bound, but rather incorporates into plasma proteins (primarily globulins). Volume of distribution: 2.5 Liters.

Metabolism

- Liver extensive

The first-pass effect in the liver in which more than 70% of glucosamine is metabolized.

Excretion

Renal Excretion, 10%

Feces, 11%

Part of a dose of glucosamine sulfate is eliminated as carbon dioxide via expired air.

Indication: Used as adjuvant therapy for osteoarthritis.

Recommended D

Light or moderate osteoarthritis symptoms:

One capsule taken twice daily for at least 6 weeks or according to medical prescription.

Severe osteoarthritis symptoms:

Initial therapy: 1 capsule to be taken 3 times daily is recommended for at least 8 weeks.

Follow- up therapy:

Maintenance therapy should be followed for 3 – 4 months (as according to medical prescription) by administration of 1 capsule taken twice daily.

The treatment of osteoarthritis should be repeated every other 6 months or less (according to medical prescription).

Route of Administration: Oral

Contraindication:

Hypersensitivity to glucosamine sulphate or patients who are allergic to shellfish. Glucosamine sulphate should not be used in patients taking warfarin (or other coumarin anticoagulants) as increased effect during concomitant treatment with glucosamine has been reported.

Glucosamine sulphate should not be used in patients with impaired renal and/ or liver function, as no dose recommendations can be given since no studies have been performed.

Patients taking oral tetracycline should not take glucosamine at the same time, as concurrent treatment with glucosamine may increase the absorption and serum concentrations of tetracyclines.

Warning & Precautions:

A doctor must be consulted to rule out the presence of joint diseases for which other treatment should be considered.

A doctor must be consulted if this medicine intended for long term use.

As the risk of diabetogenic effects is unknown and until further evidence becomes available, caution should be exercised in patients with diabetes or a predisposition for diabetes.

Therefore, in patients with impaired glucose tolerance, pre-diabetics and diagnosed diabetics (Type I and II), monitoring of the blood glucose levels should be increased, as deemed necessary, before start of treatment, periodically during treatment and at the end of the treatment, as determined by the primary Healthcare Professional.

Where relevant, monitoring of insulin requirements is also recommended before start of treatment, periodically during treatment and at the end of the treatment under medical

In patients with a known risk factor for cardiovascular disease, monitoring of the blood lipid levels is recommended during treatment with Glucosamine sulphate since hypercholesterolemia has been observed in a few patients treated with glucosamine. The results should be a factor in determining whether treatment is continued after 2-3 months.

A report on exacerbated asthma symptoms triggered after initiation of glucosamine therapy has been described (symptoms resolved after withdrawal of glucosamine). Asthmatic patients starting on glucosamine should therefore be aware of potential worsening of symptoms.

Effect on Ability to Drive and Use Machines: Not known

Interactions with Other Medicaments:

Hypoglycaemic agents

Close monitoring of blood sugar levels is recommended for diabetics on hypoglycaemic agents.

Warfarin

There are limited data on possible drug interactions with glucosamine, but increments in the INR parameter have been reported with oral vitamin K antagonists. Patients treated with oral vitamin K antagonists should therefore be closely monitored at the time of initiation or termination of glucosamine therapy.

Tetracyclines

Concurrent treatment with glucosamine may increase the absorption and serum concentration of tetracyclines, but the clinical relevance of this interaction is probably limited. If the patient is taking oral tetracycline they should not take glucosamine at the same time.

Pregnancy & Lactation:

Pregnancy

There are no adequate data from the use of glucosamine in pregnant women. From animal studies only insufficient data are available. Glucosamine sulphate should not be used during pregnancy.

Breastfeeding

There are no data available on the excretion of glucosamine into human milk. The use of glucosamine sulphate during breastfeeding is therefore not recommended as there is no data on the safety for the newborn.

Side Effects:

- Cardiovascular

Peripheral oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

- Central nervous system

Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

- Gastrointestinal

Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

- Skin

Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

Symptoms & Treatment of Overdose:

Signs and symptoms of accidental or intentional overdose might include headache, dizziness, disorientation, arthralgia, nausea, vomiting, diarrhea or constipation. In case of overdose, treatment with glucosamine should be discontinued.

Storage Condition:

This product should be stored below 30°C in a dry place, away from direct sunlight and moisture.

Keep out of reach of children.

Jauhkan daripada capaian kanak-kanak.

Shelf Life: 2 years

Packing: 60 capsules in white HDPE bottle

Distributed by Product Registration Holder:

Janipro (M) Sdn. Bhd. (838162-X)

42 Jalan Seksyen 1/21, Taman Kajang Utama,
43000 Hulu Langat, Selangor, Malaysia.

Manufacturer:

Syarikat Wen Ken Drug Sdn. Bhd.

24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.

Date of Revision: Feb 2025