

# PACKING INSERT

Registration Number:

For animal use only  
KEEP OUT OF REACH OF CHILDREN

**Product name:** SalinoCox 12% w/w Premix Granules

**Active substance Strength:** 100g:12g (12,000,000U Salinomycin)

**Dosage form:** Granules

**Product description:** Light to dark brown granule

**Pharmacodynamics:** Salinomycin is active against certain Gram-positive bacteria, while Gram-negative species are resistant. The use of Salinomycin as a feed additive at the proposed concentration is unlikely to increase shedding of *Salmonella*, *E. coli* and *Campylobacter* or to induce resistance and cross-resistance to antimicrobials important in human and animal therapy.

**Pharmacokinetics:** Salinomycin sodium is absorbed to a certain extent in the chicken and extensively metabolised. Unchanged Salinomycin represents a very small fraction of the metabolites in tissue and excreta. Many metabolites, predominantly mono- and multi-hydroxylated, have been identified in tissues and excreta. The metabolites in excreta show a higher degree of hydroxylation than in the liver. Salinomycin-related metabolites have a reduced ionophoric activity when compared with Salinomycin. Salinomycin is considered the MR; ratios MR to total residue are available for all relevant tissues for one and 6 h withdrawal. No residues in eggs are expected provided that the proposed maximum dose and duration of administration are respected.

**Indication:** For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella* in broiler chickens and in replacement pullets intended for use as caged layers.

**Target species:** Broiler chicken

**Recommended dose:** Thoroughly mix 500 g of SALINOCOX 12% to each tonne of chicken feed. This produces a salinomycin level of 60g/tonne in the finished feed which should be fed continuously as the sole ration.

**Route of Administration:** Thoroughly mix with the feed

**Interactions:** Other anticoccidial drugs should not be added to feeds containing SalinoCox, SalinoCox should not be used concurrently with Tiamulin or other antibiotics.

**Contraindication:** This product must not be fed to horses or any other animals not included in the label claim.

**Warning and Precautions:** Harmful if swallowed. May irritate the eyes. Avoid contact with eyes. Do not inhale dust. When opening the container and mixing into stockfeed wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves, goggles and disposable dust mask. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing

**Adverse effect:** Salinomycin sodium is safe for chickens for fattening at 70 mg/kg complete feed, for chickens reared for laying at 50 mg/kg complete feed in the first 12 weeks of life.

**Pregnancy and lactation:** DO NOT USE in birds laying eggs for human consumption or within 7 days of laying, where eggs or egg products are to be used for human consumption or processing.

**Incompatibility:** The simultaneous use of SalinoCox and certain antibiotic drugs (e.g. tiamulin) is contraindicated.

**Sides effects:** None known

**Overdose and treatment:** Recommended use levels should not be exceeded.

**Storage condition:** Keep sealed and stored in a place below 25°C

**Shelf Life:** 2 years

**Withdraw period:** MEAT: 0 days

EGGS: DO NOT USE in birds laying eggs for human consumption or within 7 days of laying where eggs or egg products are to be used for human consumption or processing.

Maximum Residual: Residue marker: salinomycin

Chicken: Egg 20 µg/kg Edible offal 500 µg/kg Muscle 100 µg/kg

**Packaging available:** 25kg/bag

Batch Number:

Manufacturing Date:

Expiration Date:

Manufactured by:

**SHANDONG SHENGLI BIOENGINEERING CO., LTD**

Address: No.118, Tongji Road, High-tech Development Zone, Jining, Shandong Province, China

Product Registration Holder:

**Pahang Pharmacy SDN BHD (10826-K)**

Address: Lot 5979, Jalan Teratai, 5 ½ Mile Off Jalan Meru, 41050 Klang, Selangor Darul Ehsan, Malaysia