

25 mm

FRONT

Pharmacode

ベノフロン

Date of revision: 16 July 2025

VENOFLO **Film-Coated Tablet 500mg**

26,25 mm

Pharmacode

1. NAME OF THE MEDICINAL PRODUCT

Venoflon Film-Coated Tablet 500mg

2. SPECIAL NOTICE AND RECOMMENDATION

Keep out of reach of children.

Read the package insert carefully before use.

3. COMPOSITION

Each film-coated tablet contains:

Micronized purified flavonoid fraction corresponding to:

Diosmin.....450 mg

Flavonoids expressed as hesperidin.....50 mg

4. PHARMACEUTICAL FORM

Film-coated tablet.

Brown-yellow, oval-shaped film-coated tablet, biconvex and plain on both sides.

5. INDICATIONS

(A) Treatment of symptoms related to venolymphatic insufficiency (heavy legs, pain, early morning restless legs).

(B) Treatment of functional symptoms related to acute hemorrhoidal attack.

6. ADMINISTRATION AND DOSAGE

Administration

Oral route.

Dosage

Venolymphatic insufficiency: 2 tablets daily in 2 divided doses, mid-day and evening at mealtimes.

Acute hemorrhoidal attack: 6 tablets daily (in 3 divided doses) for the first 4 days, then 4 tablets daily (in 2 divided doses) for the next 3 days.

Paediatric population: The safety and efficacy of Venoflon Film-Coated Tablet 500mg in children and adolescents under 18 years of age have not been established.

7. CONTRAINDICATIONS

Known hypersensitivity to any ingredient in the formula.

8. SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Hemorrhoidal attack: The administration of this product does not preclude treatment for other anal conditions. The treatment must be short-term. If symptoms do not subside promptly, a proctological examination should be performed and the treatment should be reviewed.

9. PREGNANCY AND LACTATION

Pregnancy

There are no or limited amount of data from the use of micronized purified flavonoid fraction in pregnant women.

Animal studies do not indicate reproductive toxicity. As a precautionary measure, it is preferable to avoid the use of Venoflon Film-Coated Tablet 500mg during pregnancy.

Lactation

It is unknown whether the active substance/metabolites are excreted in human milk.

A risk to the newborns/infants cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from Venoflon Film-Coated Tablet 500mg therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

10. EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

None known.

11. DRUG INTERACTIONS

No interaction studies have been performed. No clinically relevant drug interaction has been reported to date from post-marketing experience on the product.

12. ADVERSE REACTIONS

The adverse reactions reported consist mainly in gastrointestinal events (diarrhoea, dyspepsia, nausea, vomiting).

Adverse reactions listed below are classified according to frequency and system organ class.

Nervous system disorders: Rare: dizziness, headaches, malaise.

Gastrointestinal disorders: Common: diarrhoea, dyspepsia, nausea, vomiting; Uncommon: colitis; Not known: abdominal pain.

Skin and subcutaneous tissue disorders: Rare: rash, pruritus, urticarial; Not known: isolated face, eyelid and lip oedema. Exceptionally, Quincke's oedema.

13. OVERDOSAGE AND MANAGEMENT

Symptoms

The most frequently reported adverse events in overdose cases were gastrointestinal events (such as diarrhoea, nausea, abdominal pain) and skin events (such as pruritus, rash).

Management

Management of overdose should consist in treatment of clinical symptoms.

Venoflon Film-Coated Tablet 500mg_Insert
Dimension: 150 x 210 mm (W x L)
Code: PI20250716

25 mm

BACK

Pharmacode

Date of revision: 16 July 2025

26,25 mm

Pharmacode

14. PHARMACODYNAMIC PROPERTIES

Pharmacotherapeutic group: Vasoprotectives; capillary stabilizing agents; bioflavonoids.
ATC code: C05C A53

In pharmacology: Venoflon Film-Coated Tablet 500mg exerts a dual action on the venous return system: at vein and venule level, it increases parietal tone and exerts an anti-stasis action; at the microcirculatory level, it reinforces capillary resistance and normalizes capillary permeability.

Dose/effect relationship: Statistically significant dose-effect relationships have been demonstrated for the following venous plethysmography parameters: capacitance, distensibility and emptying time. The best dose-effect ratio is obtained with 2 tablets.

Venotonic activity: It increases venous tone: venous occlusion plethysmography with a mercury strain gauge revealed a reduction in venous emptying time.

Microcirculatory activity: In patients with signs of capillary fragility, it increases capillary resistance as measured by angiostrometry.

Efficacy and clinical safety: The medicinal product has therapeutic activity in phlebology, in the treatment of chronic venous insufficiency (functional and organic) of the lower limbs.

15. PHARMACOKINETIC PROPERTIES

In humans, following oral administration of the medicinal product with carbon 14-labelled diosmin: Excretion is essentially faecal and urinary excretion is on average of 14% of the administered quantity.

The elimination half-life is 11 hours.

The product is highly metabolised, this metabolism is revealed by the presence of different phenol acids in the urine.

16. PACKAGING

Blister of 10 tablets. Box of 6 blisters.

17. STORAGE CONDITION, SHELF-LIFE

Do not store above 30°C.

The expiry date of this pack is printed on the box. Do not use this pack after this date.

18. NAME, ADDRESS OF MANUFACTURER

STELLAPHARM J.V. CO., LTD. – BRANCH 1

(A Joint Venture Company of Auxilto GmbH, Germany)

No. 40 Tu Do Avenue, Vietnam – Singapore Industrial Park,

An Phu Ward, Thuan An Town, Binh Duong Province, Vietnam

Tel: (+84 274) 3767 470 Fax: (+84 274) 3767 469



NIKKO PHARMA*

Under License of Nikko Pharma Japan Co., Ltd
日光ファーマ株式会社

PI20250716