

Leaflet - Etopine Eye Drop 5ml

Size: 110(W) x 160(H)mm

Color: Black

Font Size: 5pt

Important information. Please read carefully.

Etopine

Eye Drops 0.01% w/v

COMPOSITION

Each 5ml contains Atropine Sulphate 0.01% w/v.
Benzalkonium Chloride 0.0075% w/v as preservative.

PHARMACODYNAMICS

Antimuscarinic drugs such as atropine are competitive inhibitors of the actions of acetylcholine at the muscarinic receptors of autonomic effector sites innervated by parasympathetic (cholinergic postganglionic) nerves; they are also inhibitors of the action of acetylcholine on smooth muscle lacking cholinergic innervation.

PHARMACOKINETICS

Atropine is readily absorbed from the gastrointestinal tract. It is also absorbed from mucous membranes, the eye, and to some extent through intact skin. It is rapidly cleared from the blood and is distributed throughout the body. It crosses the blood-brain barrier. It is incompletely metabolised in the liver and is excreted in the urine as unchanged drug and metabolites. A half-life of about 4 hours has been reported. Atropine crosses the placenta and traces appear in breast milk.

INDICATIONS

Control progression of myopia

DOSEAGE AND ADMINISTRATION

One drop topically in eye(s) once at night or as prescribed by your doctor.

CONTRAINDICATIONS

Persons with glaucoma or a tendency towards glaucoma, and with hypersensitivity to belladonna alkaloids.

WARNINGS AND PRECAUTIONS

For external use only. Excessive use may produce symptoms of systemic atropine poisoning.

DRUG INTERACTIONS

The effects of atropine and other antimuscarinics may be enhanced by use with other drugs having antimuscarinic properties, such as amantadine, some antihistamines, phenothiazine antipsychotics, and tricyclic antidepressants. Inhibition of drug-metabolising enzymes by MAOIs may possibly enhance the effects of antimuscarinics. The reduction in gastric motility caused by antimuscarinics may affect the absorption of other drugs. Antimuscarinics may also antagonise the gastrointestinal effects of cisapride, domperidone, and metoclopramide. Antimuscarinics and parasympathomimetics may counteract each other's effects

PREGNANCY AND LACTATION

This product should be used with caution during pregnancy or lactation unless potential benefits outweigh the potential risks.

Pregnancy

Atropine rapidly crosses the placenta.

Lactation

No adverse effects have been observed in breast-feeding infants whose mothers were receiving atropine.

SIDE EFFECTS

May be associated with stinging sensation on instillation.

May cause allergy (itch, irritation, and redness), blurring of near vision and glare in some children.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Excessive use or hyper-susceptibility to belladonna alkaloids may produce symptoms of atropine poisoning (palpitation, hot dry skin, dizziness, tremor, unsteady gait and change in level of consciousness). You should consult a physician if your child demonstrates these symptoms.

SHELF-LIFE

The expiry date is indicated on the packaging.

STORAGE CONDITION

Store below 30°C. Discard four weeks after opening.

PRODUCT DESCRIPTION

Clear, colourless, sterile solution.

Available in 5ml plastic dropper bottle.

KEEP OUT OF REACH OF CHILDREN

For further information, please consult your pharmacist or physician.

Revision Date: 01-November-2024

Manufactured by

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia

BRAND OWNER

Eagle Eye Centre Pte Ltd
820, Thomson Road,
Mount Alvernia Hospital
Medical Centre D, #06-57 to 62
Singapore 574623