

## PACKAGE INSERT



This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

### 1 NAME OF THE MEDICINAL PRODUCT

Versavo 25mg/ml Concentrate for Solution for Infusion (Bevacizumab)

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml of concentrate contains 25 mg of bevacizumab\*.

Each 4 ml vial contains 100 mg of bevacizumab.

Each 16 ml vial contains 400 mg of bevacizumab.

For dilution and other handling recommendations, see section 6.6.

\*Bevacizumab is a recombinant humanised monoclonal antibody produced by DNA technology in Chinese Hamster Ovary cells.

For the full list of excipients, see section 6.1.

### 3 PHARMACEUTICAL FORM

Concentrate for solution for infusion.

Clear to slightly opalescent, colourless to pale brown liquid.

### 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

##### **Metastatic Colorectal Cancer(mCRC)**

Versavo (bevacizumab) in combination with fluoropyrimidine-based chemotherapy is indicated for treatment of patients with metastatic carcinoma of the colon or rectum.

##### **Metastatic Breast Cancer(mBC)**

Versavo in combination with paclitaxel is indicated for first-line treatment of patients with metastatic breast cancer.

##### **Advanced, metastatic or recurrent Non-Small Cell Lung Cancer (NSCLC)**

Versavo, in addition to platinum-based chemotherapy, is indicated for first-line treatment of patients with unresectable advanced, metastatic or recurrent non-squamous non-small cell lung cancer.

##### **Advanced and/or metastatic Renal Cell Cancer(mRCC)**

Versavo in combination with interferon alfa-2a is indicated for first-line treatment of patients with advanced and/or metastatic renal cell cancer.

##### **Glioblastoma**

Versavo is indicated for the treatment of glioblastoma with progressive disease following prior therapy as a single agent. The effectiveness of Versavo in glioblastoma is based on an improvement in objective response rate. There are no data demonstrating an improvement in disease-related symptoms or increased survival with Versavo.

##### **Epithelial Ovarian, Fallopian Tube and Primary Peritoneal Cancer**

Versavo, in combination with carboplatin and paclitaxel is indicated for the front-line treatment of advanced (FIGO\*stages III B, III C, and IV) epithelial ovarian, fallopian tube, or primary peritoneal cancer.

Versavo, in combination with carboplatin and gemcitabine or in combination with carboplatin and paclitaxel is indicated for the treatment of patients with recurrent, platinum-sensitive, epithelial ovarian, fallopian tube, or primary peritoneal cancer who have not received prior bevacizumab or other VEGF\*-targeted angiogenesis inhibitors.

Versavo in combination with paclitaxel, topotecan, or pegylated liposomal doxorubicin is indicated for the treatment of adult patients with platinum-resistant recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who received no more than two prior chemotherapy regimens and who have not received prior therapy with bevacizumab or other VEGF inhibitors or VEGF receptor - targeted agents.

\*FIGO: International Federation of Gynaecology and Obstetrics

\*VEGF: Vascular Endothelial Growth Factor

### **Cervical Cancer**

Versavo in combination with paclitaxel and cisplatin or paclitaxel and topotecan is indicated for the treatment of persistent, recurrent, or metastatic carcinoma of the cervix.

## **4.2 Posology and method of administration**

### ***General***

Versavo should be prepared by a healthcare professional using aseptic technique (see section 4.3 Special Instructions for Use, Handling and Disposal).

The initial Versavo dose should be delivered over 90 minutes as an intravenous infusion. If the first infusion is well tolerated, the second infusion may be administered over 60 minutes. If the 60-minute infusion is well tolerated, all subsequent infusions may be administered over 30 minutes.

Dose reduction of Versavo for adverse events is not recommended. If indicated, Versavo should either be permanently discontinued or temporarily suspended as described in section 4.4 *Warnings and Precautions*.

Versavo is not formulated for intravitreal use. (*see section 4.4 Warnings and Precautions*).

### ***Metastatic Colorectal Cancer (mCRC)***

The recommended dose of Versavo, administered as an infusion, is either 5 mg/kg or 10 mg/kg of body weight given once every 2 weeks or 7.5 mg/kg or 15 mg/kg of body weight given once every 3 weeks.

It is recommended that treatment be continued until progression of the underlying disease or until unacceptable toxicity.

### ***Metastatic Breast Cancer (mBC)***

The recommended dose of Versavo is 10mg/kg of body weight given once every 2 weeks or 15 mg/kg of body weight given once every 3 weeks as an intravenous infusion.

It is recommended that Versavo treatment be continued until progression of the underlying disease.

### ***Advanced, metastatic or recurrent Non-Small Cell Lung Cancer (NSCLC)***

Versavo is administered in addition to platinum-based chemotherapy for up to 6 cycles of treatment followed by Versavo as a single agent until disease progression.

The recommended dose of Versavo when used in addition to cisplatin-based chemotherapy is 7.5 mg/kg of body weight given once every 3 weeks as an intravenous infusion.

The recommended dose of Versavo when used in addition to carboplatin-based chemotherapy is 15 mg/kg of body weight given once every 3 weeks as an intravenous infusion.

### ***Advanced and/or metastatic Renal Cell Cancer (mRCC)***

The recommended dose of Versavo is 10 mg/kg of body weight given once every 2 weeks as an intravenous infusion. It is recommended that Versavo treatment be continued until progression of the underlying disease.

### ***Glioblastoma***

The recommended dose of Versavo, administered as an infusion is 10 mg/kg of body weight given once every 2 weeks.

### ***Epithelial Ovarian, Fallopian Tube and Primary Peritoneal Cancer***

The recommended dose of Versavo administered as an intravenous infusion is as follows.

*Front-line treatment:* 15 mg/kg of body weight given once every 3 weeks when administered in addition to carboplatin and paclitaxel for up to six cycles of treatment followed by continued use of Versavo as single agent for 15 months or until disease progression, whichever occurs earlier.

*Treatment of recurrent disease:*

*Platinum-sensitive:* 15 mg/kg of body weight given once every 3 weeks when administered in combination with carboplatin and paclitaxel for 6 cycles and up to 8 cycles followed by continued use of Versavo as a single agent until disease progression.

Alternatively, 15 mg/kg every 3 weeks when administered in combination with carboplatin and gemcitabine for 6 cycles and up to 10 cycles followed by continued use of Versavo as single agent until disease progression.

*Platinum-resistant:* 10 mg/kg body weight given once every 2 weeks when administered in combination with one of the following agents – paclitaxel, topotecan (given weekly) or pegylated liposomal doxorubicin (see section 5.1, study MO22224 for chemotherapy regimens).

Alternatively, 15 mg/kg every 3 weeks when administered in combination with topotecan given on days 1-5, every 3 weeks (see section 5.1, study MO22224 for chemotherapy regimen).

It is recommended that treatment be continued until disease progression.

### ***Cervical Cancer***

Versavo is administered in combination with one of the following chemotherapy regimens: paclitaxel and cisplatin or paclitaxel and topotecan (see section 3.1.2 study GOG-0240 for further details on the chemotherapy regimens).

The recommended dose of Versavo is 15 mg/kg of body weight given once every 3 weeks as an intravenous infusion. It is recommended that Versavo treatment be continued until progression of the underlying disease.

## **4.2.1 Special Dosage Instructions**

***Pediatric use:*** The safety and efficacy of bevacizumab in children and adolescents (<18 years) have not been established (see section 4.8 *Pediatric Use*).

***Geriatric use:*** No dose adjustment is required in patients  $\geq 65$  years of age.

***Renal impairment:*** The safety and efficacy of bevacizumab have not been studied in patients with renal impairment.

***Hepatic impairment:*** The safety and efficacy of bevacizumab have not been studied in patients with hepatic impairment.

## **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Hypersensitivity to Chinese Hamster Ovary (CHO) cell products or other recombinant human or humanised antibodies.

- Pregnancy (see section 4.6).

## 4.4 Special warnings and precautions for use

### *Traceability*

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

### *Gastrointestinal (GI) perforations and Fistulae* (see section 4.8)

Patients may be at an increased risk for the development of gastrointestinal perforation and gall bladder perforation when treated with bevacizumab. Intra-abdominal inflammatory process may be a risk factor for gastrointestinal perforations in patients with metastatic carcinoma of the colon or rectum, therefore, caution should be exercised when treating these patients. Prior radiation is a risk factor for GI perforation in patients treated for persistent, recurrent or metastatic cervical cancer with bevacizumab and all patients with GI perforation had a history of prior radiation. Therapy should be permanently discontinued in patients who develop gastrointestinal perforation.

### *GI-vaginal Fistulae in study GOG-0240*

Patients treated for persistent, recurrent, or metastatic cervical cancer with bevacizumab are at increased risk of fistulae between the vagina and any part of the GI tract (Gastrointestinal-vaginal fistulae). Prior radiation is a major risk factor for the development of GI-vaginal fistulae and all patients with GI-vaginal fistulae had a history of prior radiation. Recurrence of cancer within the field of prior radiation is an additional important risk factor for the development of GI-vaginal fistulae.

### *Non-GI Fistulae* (see section 4.8)

Patients may be at increased risk for the development of fistulae when treated with bevacizumab.

Permanently discontinue Versavo in patients with tracheoesophageal (TE) fistula or any Grade 4 fistula [US National Cancer Institute-Common Terminology Criteria for Adverse Events (NCI-CTCAE v.3)]. Limited information is available on the continued use of bevacizumab in patients with other fistulae.

In cases of internal fistula not arising in the gastrointestinal tract, discontinuation of Versavo should be considered.

### *Wound healing complications* (see section 4.8)

Bevacizumab may adversely affect the wound healing process. Serious wound healing complications, including anastomotic complications, with a fatal outcome have been reported. Therapy should not be initiated for at least 28 days following major surgery or until the surgical wound is fully healed. In patients who experienced wound healing complications during therapy, treatment should be withheld until the wound is fully healed. Therapy should be withheld for elective surgery.

Necrotising fasciitis, including fatal cases, has rarely been reported in patients treated with bevacizumab. This condition is usually secondary to wound healing complications, gastrointestinal perforation or fistula formation. Versavo therapy should be discontinued in patients who develop necrotising fasciitis, and appropriate treatment should be promptly initiated.

### *Hypertension* (see section 4.8)

An increased incidence of hypertension was observed in bevacizumab -treated patients. Clinical safety data suggest that the incidence of hypertension is likely to be dose-dependent. Pre-existing hypertension should be adequately controlled before starting Versavo treatment. There is no information on the effect of bevacizumab in patients with uncontrolled hypertension at the time of initiating therapy. Monitoring of blood pressure is generally recommended during therapy.

In most cases hypertension was controlled adequately using standard antihypertensive treatment appropriate for the individual situation of the affected patient. The use of diuretics to manage hypertension is not advised in patients who receive a cisplatin-based chemotherapy regimen. Versavo should be permanently discontinued if medically significant hypertension cannot be adequately controlled with antihypertensive therapy, or if the patient develops hypertensive crisis or hypertensive encephalopathy.

*Posterior Reversible Encephalopathy Syndrome (PRES)* (see section 4.8)

There have been rare reports of bevacizumab -treated patients developing signs and symptoms that are consistent with PRES, a rare neurologic disorder, which can present with the following signs and symptoms among others: seizures, headache, altered mental status, visual disturbance, or cortical blindness, with or without associated hypertension. A diagnosis of PRES requires confirmation by brain imaging, preferably magnetic resonance imaging (MRI). In patients developing PRES, treatment of specific symptoms including control of hypertension is recommended along with discontinuation of Versavo. The safety of reinitiating bevacizumab therapy in patients previously experiencing PRES is not known.

*Proteinuria* (see section 4.8)

Patients with a history of hypertension may be at increased risk for the development of proteinuria when treated with bevacizumab. There is evidence suggesting that all Grade (US National Cancer Institute-Common Terminology Criteria for Adverse Events [NCI-CTCAE v.3]) proteinuria may be related to the dose. Monitoring of proteinuria by dipstick urinalysis is recommended prior to starting and during therapy. Grade 4 proteinuria (nephrotic syndrome) was seen in up to 1.4% of patients treated with bevacizumab. Therapy should be permanently discontinued in patients who develop nephrotic syndrome (NCI-CTCAE v.3).

*Arterial thromboembolism* (see section 4.8)

In clinical trials, the incidence of arterial thromboembolic reactions including cerebrovascular accidents (CVAs), transient ischaemic attacks (TIAs) and myocardial infarctions (MIs) was higher in patients receiving bevacizumab in combination with chemotherapy compared to those who received chemotherapy alone. Patients receiving bevacizumab plus chemotherapy, with a history of arterial thromboembolism, diabetes or age greater than 65 years have an increased risk of developing arterial thromboembolic reactions during therapy. Caution should be taken when treating these patients with Versavo.

Therapy should be permanently discontinued in patients who develop arterial thromboembolic reactions.

*Venous thromboembolism* (see section 4.8)

Patients may be at risk of developing venous thromboembolic reactions, including pulmonary embolism under bevacizumab treatment.

Patients treated for persistent, recurrent, or metastatic cervical cancer with bevacizumab in combination with paclitaxel and cisplatin may be at increased risk of venous thromboembolic events.

Versavo should be discontinued in patients with life-threatening (Grade 4) thromboembolic reactions, including pulmonary embolism (NCI-CTCAE v.3). Patients with thromboembolic reactions  $\leq$  Grade 3 need to be closely monitored (NCI-CTCAE v.3).

*Haemorrhage*

Patients treated with bevacizumab have an increased risk of haemorrhage, especially tumour-associated haemorrhage. Versavo should be discontinued permanently in patients who experience Grade 3 or 4 bleeding during Versavo therapy (NCI-CTCAE v.3) (see section 4.8).

Patients with untreated CNS metastases were routinely excluded from clinical trials with bevacizumab, based on imaging procedures or signs and symptoms. Therefore, the risk of CNS haemorrhage in such patients has not been prospectively evaluated in randomised clinical trials (see section 4.8). Patients should be monitored for signs and symptoms of CNS bleeding, and Versavo treatment discontinued in cases of intracranial bleeding.

There is no information on the safety profile of bevacizumab in patients with congenital bleeding diathesis, acquired coagulopathy or in patients receiving full dose of anticoagulants for the treatment of thromboembolism prior to starting bevacizumab treatment, as such patients were excluded from clinical trials. Therefore, caution should be exercised before initiating therapy in these patients. However, patients who developed venous thrombosis while receiving therapy did not appear to have an increased rate of Grade 3 or above bleeding when treated with a full dose of warfarin and bevacizumab concomitantly (NCI-CTCAE v.3).

*Pulmonary haemorrhage/haemoptysis*

Patients with non-small cell lung cancer treated with Versavo may be at risk of serious, and in some cases fatal, pulmonary haemorrhage/haemoptysis. Patients with recent pulmonary haemorrhage/ haemoptysis ( $> 2.5$  ml of red blood) should not be treated with Versavo.

#### *Aneurysms and artery dissections*

The use of VEGF pathway inhibitors in patients with or without hypertension may promote the formation of aneurysms and/or artery dissections. Before initiating Versavo, this risk should be carefully considered in patients with risk factors such as hypertension or history of aneurysm.

#### *Congestive heart failure (CHF) (see section 4.8)*

Reactions consistent with CHF were reported in clinical trials. The findings ranged from asymptomatic declines in left ventricular ejection fraction to symptomatic CHF, requiring treatment or hospitalisation. Caution should be exercised when treating patients with clinically significant cardiovascular disease such as pre-existing coronary artery disease, or congestive heart failure with Versavo.

Most of the patients who experienced CHF had metastatic breast cancer and had received previous treatment with anthracyclines, prior radiotherapy to the left chest wall or other risk factors for CHF were present.

In patients in AVF3694g who received treatment with anthracyclines and who had not received anthracyclines before, no increased incidence of all Grade CHF was observed in the anthracycline + bevacizumab group compared to the treatment with anthracyclines only. CHF Grade 3 or higher reactions were somewhat more frequent among patients receiving bevacizumab in combination with chemotherapy than in patients receiving chemotherapy alone. This is consistent with results in patients in other studies of metastatic breast cancer who did not receive concurrent anthracycline treatment (NCI-CTCAE v.3) (see section 4.8).

#### *Neutropenia and infections (see section 4.8)*

Increased rates of severe neutropenia, febrile neutropenia, or infection with or without severe neutropenia (including some fatalities) have been observed in patients treated with some myelotoxic chemotherapy regimens plus bevacizumab in comparison to chemotherapy alone. This has mainly been seen in combination with platinum- or taxane-based therapies in the treatment of NSCLC, mBC, and in combination with paclitaxel and topotecan in persistent, recurrent, or metastatic cervical cancer.

#### *Hypersensitivity reactions (including anaphylactic shock)/infusion reactions (see section 4.8)*

Patients may be at risk of developing infusion/hypersensitivity reactions (including anaphylactic shock). Close observation of the patient during and following the administration of bevacizumab is recommended as expected for any infusion of a therapeutic humanised monoclonal antibody. If a reaction occurs, the infusion should be discontinued and appropriate medical therapies should be administered. A systematic premedication is not warranted.

#### *Osteonecrosis of the jaw (ONJ) (see section 4.8)*

Cases of ONJ have been reported in cancer patients treated with bevacizumab, the majority of whom had received prior or concomitant treatment with intravenous bisphosphonates, for which ONJ is an identified risk. Caution should be exercised when Versavo and intravenous bisphosphonates are administered simultaneously or sequentially.

Invasive dental procedures are also an identified risk factor. A dental examination and appropriate preventive dentistry should be considered prior to starting the treatment with Versavo. In patients who have previously received or are receiving intravenous bisphosphonates invasive dental procedures should be avoided, if possible.

#### *Intravitreal use*

Versavo is not formulated for intravitreal use.

#### *Eye disorders*

Individual cases and clusters of serious ocular adverse reactions have been reported following unapproved intravitreal use of bevacizumab compounded from vials approved for intravenous administration in cancer patients. These reactions included infectious endophthalmitis, intraocular inflammation such as sterile endophthalmitis, uveitis and vitritis, retinal detachment, retinal pigment epithelial tear, intraocular pressure increased, intraocular haemorrhage such as vitreous haemorrhage or retinal haemorrhage and conjunctival haemorrhage. Some of these reactions have resulted in various degrees of visual loss, including permanent blindness.

#### *Systemic effects following intravitreal use*

A reduction of circulating VEGF concentration has been demonstrated following intravitreal anti-VEGF therapy. Systemic adverse reactions including non-ocular haemorrhages and arterial thromboembolic reactions have been reported following intravitreal injection of VEGF inhibitors.

#### *Ovarian failure/fertility*

Bevacizumab may impair female fertility (see sections 4.6 and 4.8). Therefore, fertility preservation strategies should be discussed with women of child-bearing potential prior to starting treatment with Versavo.

#### **Important information about some of the ingredients of Versavo**

This medicine contains less than 1 mmol sodium (23mg) per vial, that is to say essentially 'sodium free'.

### **4.5 Interaction with other medicinal products and other forms of interaction**

#### *Effect of antineoplastic agents on bevacizumab pharmacokinetics*

No clinically relevant interaction of co-administered chemotherapy on bevacizumab pharmacokinetics was observed based on the results of population pharmacokinetic analyses. There were neither statistically significant nor clinically relevant differences in bevacizumab clearance in patients receiving bevacizumab monotherapy compared to patients receiving bevacizumab in combination with interferon alfa-2a, erlotinib or chemotherapies (IFL, 5-FU/LV, carboplatin/paclitaxel, capecitabine, doxorubicin or cisplatin/gemcitabine).

#### *Effect of bevacizumab on the pharmacokinetics of other antineoplastic agents*

No clinically relevant interaction of bevacizumab was observed on the pharmacokinetics of co-administered interferon alfa 2a, erlotinib (and its active metabolite OSI-420), or the chemotherapies irinotecan (and its active metabolite SN38), capecitabine, oxaliplatin (as determined by measurement of free and total platinum), and cisplatin. Conclusions on the impact of bevacizumab on gemcitabine pharmacokinetics cannot be drawn.

#### *Combination of bevacizumab and sunitinib malate*

In two clinical trials of metastatic renal cell carcinoma, microangiopathic haemolytic anaemia (MAHA) was reported in 7 of 19 patients treated with bevacizumab (10 mg/kg every two weeks) and sunitinib malate (50 mg daily) combination.

MAHA is a haemolytic disorder which can present with red cell fragmentation, anaemia, and thrombocytopenia. In addition, hypertension (including hypertensive crisis), elevated creatinine, and neurological symptoms were observed in some of these patients. All of these findings were reversible upon discontinuation of bevacizumab and sunitinib malate (see *Hypertension, Proteinuria, PRES* in section 4.4).

#### *Combination with platinum- or taxane-based therapies* (see sections 4.4 and 4.8)

Increased rates of severe neutropenia, febrile neutropenia, or infection with or without severe neutropenia (including some fatalities) have been observed mainly in patients treated with platinum- or taxane-based therapies in the treatment of NSCLC and mBC.

#### *Radiotherapy*

The safety and efficacy of concomitant administration of radiotherapy and bevacizumab has not been established.

#### *EGFR monoclonal antibodies in combination with bevacizumab chemotherapy regimens*

No interaction studies have been performed. EGFR monoclonal antibodies should not be administered for the treatment of mCRC in combination with bevacizumab-containing chemotherapy. Results from the randomised phase III studies, PACCE and CAIRO-2, in patients with mCRC suggest that the use of anti-EGFR monoclonal antibodies panitumumab and cetuximab, respectively, in combination with bevacizumab plus chemotherapy, is associated with decreased PFS and/or OS, and with increased toxicity compared with bevacizumab plus chemotherapy alone.

## 4.6 Fertility, pregnancy and lactation

### *Women of childbearing potential*

Women of childbearing potential have to use effective contraception during (and up to 6 months after) treatment.

### *Pregnancy*

There are no clinical trial data on the use of bevacizumab in pregnant women. Studies in animals have shown reproductive toxicity including malformations (see section 5.3). IgGs are known to cross the placenta, and bevacizumab is anticipated to inhibit angiogenesis in the foetus, and thus is suspected to cause serious birth defects when administered during pregnancy. In the post-marketing setting, cases of foetal abnormalities in women treated with bevacizumab alone or in combination with known embryotoxic chemotherapeutics have been observed (see section 4.8). Versavo is contraindicated in pregnancy (see section 4.3).

### *Breast-feeding*

It is not known whether bevacizumab is excreted in human milk. As maternal IgG is excreted in milk and bevacizumab could harm infant growth and development (see section 5.3), women must discontinue breast-feeding during therapy and not breast-feed for at least six months following the last dose of Versavo.

### *Fertility*

Repeat dose toxicity studies in animals have shown that bevacizumab may have an adverse effect on female fertility (see section 5.3). In a phase III trial in the adjuvant treatment of patients with colon cancer, a substudy with premenopausal women has shown a higher incidence of new cases of ovarian failure in the bevacizumab group compared to the control group. After discontinuation of bevacizumab treatment, ovarian function recovered in the majority of patients. Long term effects of the treatment with bevacizumab on fertility are unknown.

## 4.7 Effects on ability to drive and use machines

Bevacizumab has no or negligible influence on the ability to drive and use machines. However, somnolence and syncope have been reported with bevacizumab use (see table 1 in section 4.8). If patients are experiencing symptoms that affect their vision or concentration, or their ability to react, they should be advised not to drive and use machines until symptoms abate.

## 4.8 Undesirable effects

### Summary of the safety profile

The overall safety profile of bevacizumab is based on data from over 5,700 patients with various malignancies, predominantly treated with bevacizumab in combination with chemotherapy in clinical trials.

The most serious adverse reactions were:

- Gastrointestinal perforations (see section 4.4).
- Haemorrhage, including pulmonary haemorrhage/haemoptysis, which is more common in non-small cell lung cancer patients (see section 4.4).
- Arterial thromboembolism (see section 4.4).

The most frequently observed adverse reactions across clinical trials in patients receiving bevacizumab were hypertension, fatigue or asthenia, diarrhoea and abdominal pain.

Analyses of the clinical safety data suggest that the occurrence of hypertension and proteinuria with bevacizumab therapy are likely to be dose-dependent.

### Tabulated list of adverse reactions

The adverse reactions listed in this section fall into the following frequency categories: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data).

Tables 1 and 2 list adverse reactions associated with the use of bevacizumab in combination with different chemotherapy regimens in multiple indications, by MedDRA system organ class.



Table 1 provides all adverse reactions by frequency that were determined to have a causal relationship with bevacizumab through:

- comparative incidences noted between clinical trial treatment arms (with at least a 10% difference compared to the control arm for NCI-CTCAE Grade 1-5 reactions or at least a 2% difference compared to the control arm for NCI-CTCAE Grade 3-5 reactions,
- post-authorisation safety studies,
- spontaneous reporting,
- epidemiological studies\non-interventional or observational studies,
- or through an evaluation of individual case reports.

Table 2 provides the frequency of severe adverse reactions. Severe reactions are defined as adverse events with at least a 2% difference compared to the control arm in clinical studies for NCI-CTCAE Grade 3-5 reactions. Table 2 also includes adverse reactions which are considered by the MAH to be clinically significant or severe.

Post-marketing adverse reactions are included in both Tables 1 and 2, where applicable. Detailed information about these post-marketing reactions are provided in Table 3.

Adverse reactions are added to the appropriate frequency category in the tables below according to the highest incidence seen in any indication.

Within each frequency category, adverse reactions are presented in the order of decreasing seriousness.

Some of the adverse reactions are reactions commonly seen with chemotherapy; however, bevacizumab may exacerbate these reactions when combined with chemotherapeutic agents. Examples include palmar-plantar erythrodysesthesia syndrome with pegylated liposomal doxorubicin or capecitabine, peripheral sensory neuropathy with paclitaxel or oxaliplatin, nail disorders or alopecia with paclitaxel, and paronychia with erlotinib.

**Table 1: Adverse Reactions by Frequency**

| class                                | Very Common                                                                  | Common                                                                          | Uncommon | Rare                                                           | Very Rare                                | Frequency Not Known |
|--------------------------------------|------------------------------------------------------------------------------|---------------------------------------------------------------------------------|----------|----------------------------------------------------------------|------------------------------------------|---------------------|
| Infections and infestations          |                                                                              | Sepsis, Abscess <sup>b,d</sup> , Cellulitis, Infection, Urinary tract infection |          | Necrotising fasciitis <sup>a</sup>                             |                                          |                     |
| Blood and lymphatic system disorders | Febrile neutropenia, Leucopenia, Neutropenia <sup>b</sup> , Thrombocytopenia | Anaemia, Lymphopenia                                                            |          |                                                                |                                          |                     |
| Immune system disorders              |                                                                              | Hypersensitivity, infusion reactions <sup>a,b,d</sup>                           |          | Anaphylactic shock                                             |                                          |                     |
| Metabolism and nutrition disorders   | Anorexia, Hypomagnesaemia, Hyponatraemia                                     | Dehydration                                                                     |          |                                                                |                                          |                     |
| Nervous system disorders             | Peripheral sensory neuropathy <sup>b</sup> , Dysarthria, Headache, Dysgeusia | Cerebrovascular accident, Syncope, Somnolence                                   |          | Posterior reversible encephalopathy syndrome <sup>a, b,d</sup> | Hypertensive encephalopathy <sup>a</sup> |                     |

| class                                           | Very Common                                                                                       | Common                                                                                                                                                                             | Uncommon | Rare | Very Rare | Frequency Not Known                                                                   |
|-------------------------------------------------|---------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|------|-----------|---------------------------------------------------------------------------------------|
| Eye disorders                                   | Eye disorder, Lacrimation increased                                                               |                                                                                                                                                                                    |          |      |           |                                                                                       |
| Cardiac disorders                               |                                                                                                   | Congestive heart failure <sup>b,d</sup> , Supraventricular tachycardia                                                                                                             |          |      |           |                                                                                       |
| Vascular disorders                              | Hypertension <sup>b,d</sup> , Thromboembolism (venous) <sup>b,d</sup>                             | Thrombo-embolism (arterial) <sup>b,d</sup> , Haemorrhage <sup>b,d</sup> , Deep vein thrombosis                                                                                     |          |      |           | Renal thrombotic microangiopathy <sup>a,b</sup> , Aneurysms and artery dissections    |
| Respiratory, thoracic and mediastinal disorders | Dyspnoea, Rhinitis, Epistaxis, Cough                                                              | Pulmonary haemorrhage/ Haemoptysis <sup>b,d</sup> , Pulmonary embolism, Hypoxia, Dysphonia <sup>a</sup>                                                                            |          |      |           | Pulmonary hypertension <sup>a</sup> , Nasal septum perforation <sup>a</sup>           |
| Gastrointestinal disorders                      | Rectal haemorrhage, Stomatitis, Constipation, Diarrhoea, Nausea, Vomiting, Abdominal pain         | Gastrointestinal perforation <sup>b,d</sup> , Intestinal perforation, Ileus, Intestinal obstruction, Recto-vaginal fistulae <sup>d,e</sup> , Gastrointestinal Disorder, Proctalgia |          |      |           | Gastrointestinal ulcer <sup>a</sup>                                                   |
| Hepatobiliary disorders                         |                                                                                                   |                                                                                                                                                                                    |          |      |           | Gallbladder perforation <sup>a,b</sup>                                                |
| Skin and subcutaneous tissue disorders          | Wound healing complications <sup>b,d</sup> , Exfoliative dermatitis, Dry skin, Skin discoloration | Palmar-plantar erythro-dysaesthesia syndrome                                                                                                                                       |          |      |           |                                                                                       |
| Musculoskeletal and connective tissue disorders | Arthralgia, Myalgia                                                                               | Fistula <sup>b,d</sup> , Muscular weakness, Back pain                                                                                                                              |          |      |           | Osteonecrosis of the jaw <sup>a,b</sup> , Non-mandibular osteonecrosis <sup>a,f</sup> |
| Renal and urinary disorders                     | Proteinuria <sup>b,d</sup>                                                                        |                                                                                                                                                                                    |          |      |           |                                                                                       |
| Reproductive system and breast disorders        | Ovarian failure <sup>b,c,d</sup>                                                                  | Pelvic Pain                                                                                                                                                                        |          |      |           |                                                                                       |

| class                                                | Very Common                                            | Common   | Uncommon | Rare | Very Rare | Frequency Not Known                 |
|------------------------------------------------------|--------------------------------------------------------|----------|----------|------|-----------|-------------------------------------|
| Congenital, familial, and genetic disorder           |                                                        |          |          |      |           | Foetal abnormalities <sup>a,b</sup> |
| General disorders and administration site conditions | Asthenia, Fatigue, Pyrexia, Pain, Mucosal inflammation | Lethargy |          |      |           |                                     |
| Investigations                                       | Weight decreased                                       |          |          |      |           |                                     |

When events were noted as both all grade and grade 3-5 adverse drug reactions in clinical trials, the highest frequency observed in patients has been reported. Data are unadjusted for the differential time on treatment.

<sup>a</sup> For further information please refer to Table 3 'Adverse reactions reported in post-marketing setting.'

<sup>b</sup> Terms represent a group of events that describe a medical concept rather than a single condition or MedDRA (Medical Dictionary for Regulatory Activities) preferred term. This group of medical terms may involve the same underlying pathophysiology (e.g. arterial thromboembolic reactions include cerebrovascular accident, myocardial infarction, transient ischaemic attack and other arterial thromboembolic reactions).

<sup>c</sup> Based on a substudy from NSABP C-08 with 295 patients

<sup>d</sup> For additional information refer below within section "Further information on selected serious adverse reactions."

<sup>e</sup> Recto-vaginal fistulae are the most common fistulae in the GI-vaginal fistula category.

<sup>f</sup> Observed in pediatric population only

**Table 2: Severe Adverse Reactions by Frequency**

| System organ class                   | Very Common                                                                  | Common                                                                          | Uncommon | Rare               | Very Rare | Frequency Not Known                |
|--------------------------------------|------------------------------------------------------------------------------|---------------------------------------------------------------------------------|----------|--------------------|-----------|------------------------------------|
| Infections and infestations          |                                                                              | Sepsis, Cellulitis, Abscess <sup>a,b</sup> , Infection, Urinary tract infection |          |                    |           | Necrotising fasciitis <sup>c</sup> |
| Blood and lymphatic system disorders | Febrile neutropenia, Leucopenia, Neutropenia <sup>a</sup> , Thrombocytopenia | Anaemia, Lymphopenia                                                            |          |                    |           |                                    |
| Immune system disorders              |                                                                              | Hypersensitivity, infusion reactions <sup>a,b,c</sup>                           |          | Anaphylactic shock |           |                                    |
| Metabolism and nutrition disorders   |                                                                              | Dehydration Hyponatraemia                                                       |          |                    |           |                                    |

| System organ class                              | Very Common                                 | Common                                                                                                                                           | Uncommon | Rare | Very Rare | Frequency Not Known                                                                                      |
|-------------------------------------------------|---------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|----------|------|-----------|----------------------------------------------------------------------------------------------------------|
| Nervous system disorders                        | Peripheral sensory neuropathy <sup>a</sup>  | Cerebrovascular accident, Syncope, Somnolence, Headache                                                                                          |          |      |           | Posterior reversible encephalopathy syndrome <sup>a,b,c</sup> , Hypertensive encephalopathy <sup>c</sup> |
| Cardiac disorders                               |                                             | Congestive heart failure <sup>a,b</sup> , Supraventricular tachycardia                                                                           |          |      |           |                                                                                                          |
| Vascular disorders                              | Hypertension <sup>a,b</sup>                 | Thromboembolism arterial <sup>a,b</sup> , Haemorrhage <sup>a,b</sup> , Thromboembolism (venous) <sup>a,b</sup> , Deep vein thrombosis            |          |      |           | Renal thrombotic microangiopathy <sup>b,c</sup> , Aneurysms and artery dissections                       |
| Respiratory, thoracic and mediastinal disorders |                                             | Pulmonary haemorrhage/ Haemoptysis <sup>a,b</sup> , Pulmonary embolism, Epistaxis, Dyspnoea, Hypoxia                                             |          |      |           | Pulmonary hypertension <sup>c</sup> , Nasal septum perforation <sup>c</sup>                              |
| Gastrointestinal disorders                      | Diarrhoea, Nausea, Vomiting, Abdominal pain | Intestinal perforation, Ileus, Intestinal obstruction, Recto-vaginal fistulae <sup>c,d</sup> , Gastrointestinal disorder, Stomatitis, Proctalgia |          |      |           | Gastrointestinal perforation <sup>a,b</sup> , Gastrointestinal ulcer <sup>c</sup> , Rectal haemorrhage   |
| Hepatobiliary disorders                         |                                             |                                                                                                                                                  |          |      |           | Gallbladder perforation <sup>b,c</sup>                                                                   |
| Skin and subcutaneous tissue disorders          |                                             | Wound healing complications <sup>a,b</sup> , Palmar-plantar erythrodysesthesia syndrome                                                          |          |      |           |                                                                                                          |
| Musculoskeletal and connective tissue disorders |                                             | Fistula <sup>a,b</sup> , Myalgia, Arthralgia, Muscular weakness, Back Pain                                                                       |          |      |           | Osteonecrosis of the jaw <sup>b,c</sup>                                                                  |
| Renal and urinary disorders                     |                                             | Proteinuria <sup>a,b</sup>                                                                                                                       |          |      |           |                                                                                                          |
| Reproductive system and breast disorders        |                                             | Pelvic pain                                                                                                                                      |          |      |           | Ovarian failure <sup>a,b</sup>                                                                           |

| System organ class                                   | Very Common        | Common                               | Uncommon | Rare | Very Rare | Frequency Not Known                 |
|------------------------------------------------------|--------------------|--------------------------------------|----------|------|-----------|-------------------------------------|
| Congenital, familial, and genetic disorder           |                    |                                      |          |      |           | Foetal abnormalities <sup>a,c</sup> |
| General disorders and administration site conditions | Asthenia, Fatigue, | Pain, Lethargy, Mucosal Inflammation |          |      |           |                                     |

Table 2 provides the frequency of severe adverse reactions. Severe reactions are defined as adverse events with at least a 2% difference compared to the control arm in clinical studies for NCI-CTCAE Grade 3-5 reactions. Table 2 also includes adverse reactions which are considered by the MAH to be clinically significant or severe. These clinically significant adverse reactions were reported in clinical trials but the grade 3-5 reactions did not meet the threshold of at least a 2% difference compared to the control arm. Table 2 also includes clinically significant adverse reactions that were observed only in the postmarketing setting, therefore, the frequency and NCI-CTCAE grade is not known. These clinically significant reactions have therefore been included in Table 2 within the column entitled "Frequency Not Known."

<sup>a</sup> Terms represent a group of events that describe a medical concept rather than a single condition or MedDRA (Medical Dictionary for Regulatory Activities) preferred term. This group of medical terms may involve the same underlying pathophysiology (e.g. arterial thromboembolic reactions include cerebrovascular accident, myocardial infarction, transient ischaemic attack and other arterial thromboembolic reactions).

<sup>b</sup> For additional information refer below within section "Further information on selected serious adverse reactions"

<sup>c</sup> For further information please refer to Table 3 'Adverse reactions reported in post-marketing setting.'

<sup>d</sup> Recto-vaginal fistulae are the most common fistulae in the GI-vaginal fistula category

#### Description of selected serious adverse reactions

##### *Gastrointestinal (GI) perforations and Fistulae* (see section 4.4)

Bevacizumab has been associated with serious cases of gastrointestinal perforation.

Gastrointestinal perforations have been reported in clinical trials with an incidence of less than 1% in patients with non-squamous non-small cell lung cancer, up to 1.3% in patients with metastatic breast cancer, up to 2.0% in patients with metastatic renal cell cancer or in patients with ovarian cancer, and up to 2.7% (including gastrointestinal fistula and abscess) in patients with metastatic colorectal cancer. From a clinical trial in patients with persistent, recurrent, or metastatic cervical cancer (study GOG-0240), GI perforations (all grade) were reported in 3.2% of patients, all of whom had a history of prior pelvic radiation.

The occurrence of those events varied in type and severity, ranging from free air seen on the plain abdominal X-ray, which resolved without treatment, to intestinal perforation with abdominal abscess and fatal outcome. In some cases underlying intra-abdominal inflammation was present, either from gastric ulcer disease, tumour necrosis, diverticulitis, or chemotherapy-associated colitis.

Fatal outcome was reported in approximately a third of serious cases of gastrointestinal perforations, which represents between 0.2%-1% of all Versavo treated patients.

In bevacizumab clinical trials, gastrointestinal fistulae (all grade) have been reported with an incidence of up to 2% in patients with metastatic colorectal cancer and ovarian cancer, but were also reported less commonly in patients with other types of cancer.

##### *GI-vaginal Fistulae in study GOG-0240*

In a trial of patients with persistent, recurrent or metastatic cervical cancer, the incidence of GI-vaginal fistulae was 8.3% in bevacizumab-treated patients and 0.9% in control patients, all of whom had a history of prior pelvic radiation. The frequency of GI-vaginal fistulae in the group treated with bevacizumab + chemotherapy was higher in patients with recurrence within the field of prior radiation (16.7%) compared

with patients with no prior radiation and/ or no recurrence inside the field of prior radiation (3.6%). The corresponding frequencies in the control group receiving chemotherapy alone were 1.1% vs. 0.8%, respectively. Patients who develop GI-vaginal fistulae may also have bowel obstructions and require surgical intervention as well as diverting ostomies.

#### *Non-GI Fistulae* (see section 4.4)

Bevacizumab use has been associated with serious cases of fistulae including reactions resulting in death. From a clinical trial in patients with persistent, recurrent, or metastatic cervical cancer (GOG-240), 1.8% of bevacizumab-treated patients and 1.4% of control patients were reported to have had non-gastrointestinal vaginal, vesical, or female genital tract fistulae.

Uncommon ( $\geq 0.1\%$  to  $< 1\%$ ) reports of fistulae that involve areas of the body other than the gastrointestinal tract (e.g. bronchopleural and biliary fistulae) were observed across various indications. Fistulae have also been reported in post-marketing experience.

Reactions were reported at various time points during treatment ranging from one week to greater than 1 year from initiation of bevacizumab, with most reactions occurring within the first 6 months of therapy.

#### *Wound healing* (see section 4.4)

As bevacizumab may adversely impact wound healing, patients who had major surgery within the last 28 days were excluded from participation in phase III clinical trials.

In clinical trials of metastatic carcinoma of the colon or rectum, there was no increased risk of post-operative bleeding or wound healing complications observed in patients who underwent major surgery 28-60 days prior to starting bevacizumab. An increased incidence of post-operative bleeding or wound healing complication occurring within 60 days of major surgery was observed if the patient was being treated with bevacizumab at the time of surgery. The incidence varied between 10% (4/40) and 20% (3/15).

Serious wound healing complications, including anastomotic complications, have been reported, some of which had a fatal outcome.

In locally recurrent and metastatic breast cancer trials, Grade 3-5 wound healing complications were observed in up to 1.1% of patients receiving bevacizumab compared with up to 0.9% of patients in the control arms (NCI-CTCAE v.3).

In clinical trials of ovarian cancer, Grade 3-5 wound healing complications were observed in up to 1.8% of patients in the bevacizumab arm versus 0.1% in the control arm (NCI-CTCAE v.3)

#### *Hypertension* (see section 4.4)

In clinical trials, with the exception of study JO25567, the overall incidence of hypertension (all grades) ranged up to 42.1% in the bevacizumab containing arms compared with up to 14% in the control arms. The overall incidence of NCI-CTC Grade 3 and 4 hypertension in patients receiving bevacizumab ranged from 0.4% to 17.9%. Grade 4 hypertension (hypertensive crisis) occurred in up to 1.0% of patients treated with bevacizumab and chemotherapy compared to up to 0.2% of patients treated with the same chemotherapy alone.

In study JO25567, all grade hypertension was observed in 77.3% of the patients who received bevacizumab in combination with erlotinib as first-line treatment for non-squamous NSCLC with EGFR activating mutations, compared to 14.3% of patients treated with erlotinib alone. Grade 3 hypertension was 60.0% in patients treated with bevacizumab in combination with erlotinib compared to 11.7% in patients treated with erlotinib alone. There were no grade 4 or 5 hypertension events.

Hypertension was generally adequately controlled with oral anti-hypertensives such as angiotensin-converting enzyme inhibitors, diuretics and calcium-channel blockers. It rarely resulted in discontinuation of bevacizumab treatment or hospitalisation.

Very rare cases of hypertensive encephalopathy have been reported, some of which were fatal.

The risk of bevacizumab-associated hypertension did not correlate with the patients' baseline characteristics, underlying disease or concomitant therapy.

#### *Posterior Reversible Encephalopathy Syndrome* (see section 4.4)

There have been rare reports of bevacizumab -treated patients developing signs and symptoms that are consistent with PRES, a rare neurological disorder. Presentation may include seizures, headache, altered

mental status, visual disturbance, or cortical blindness, with or without associated hypertension. The clinical presentation of PRES is often nonspecific, and therefore the diagnosis of PRES requires confirmation by brain imaging, preferably MRI.

In patients developing PRES, early recognition of symptoms with prompt treatment of specific symptoms including control of hypertension (if associated with severe uncontrolled hypertension) is recommended in addition to discontinuation of bevacizumab therapy. Symptoms usually resolve or improve within days after treatment discontinuation, although some patients have experienced some neurologic sequelae. The safety of reinitiating Versavo therapy in patients previously experiencing PRES is not known.

Across clinical trials, 8 cases of PRES have been reported. Two of the eight cases did not have radiological confirmation via MRI.

#### *Proteinuria* (see section 4.4)

In clinical trials, proteinuria has been reported within the range of 0.7% to 54.7% of patients receiving bevacizumab.

Proteinuria ranged in severity from clinically asymptomatic, transient, trace proteinuria to nephrotic syndrome, with the great majority as Grade 1 proteinuria (NCI-CTCAE v.3). Grade 3 proteinuria was reported in up to 10.9% of treated patients. Grade 4 proteinuria (nephrotic syndrome) was seen in up to 1.4% of treated patients. Testing for proteinuria is recommended prior to start of Versavo therapy. In most clinical trials urine protein levels of  $\geq 2\text{g}/24\text{ hrs}$  led to the holding of bevacizumab until recovery to  $< 2\text{g}/24\text{ hrs}$ .

#### *Haemorrhage* (see section 4.4)

In clinical trials across all indications the overall incidence of NCI-CTCAE v.3 Grade 3-5 bleeding reactions ranged from 0.4% to 6.9% in bevacizumab treated patients, compared with up to 4.5% of patients in the chemotherapy control group.

From a clinical trial in patients with persistent, recurrent, or metastatic cervical cancer (study GOG-0240), grade 3-5 bleeding reactions have been reported in up to 8.3% of patients treated with bevacizumab in combination with paclitaxel and topotecan compared with up to 4.6% of patients treated with paclitaxel and topotecan.

The haemorrhagic reactions that have been observed in clinical trials were predominantly tumour-associated haemorrhage (see below) and minor mucocutaneous haemorrhage (e.g. epistaxis).

#### *Tumour-associated haemorrhage* (see section 4.4)

Major or massive pulmonary haemorrhage/haemoptysis has been observed primarily in trials in patients with non-small cell lung cancer (NSCLC). Possible risk factors include squamous cell histology, treatment with antirheumatic/anti-inflammatory substances, treatment with anticoagulants, prior radiotherapy, bevacizumab therapy, previous medical history of atherosclerosis, central tumour location and cavitation of tumours prior to or during therapy. The only variables that showed statistically significant correlations with bleeding were bevacizumab therapy and squamous cell histology. Patients with NSCLC of known squamous cell histology or mixed cell type with predominant squamous cell histology were excluded from subsequent phase III trials, while patients with unknown tumour histology were included.

In patients with NSCLC excluding predominant squamous histology, all Grade reactions were seen with a frequency of up to 9.3% when treated with bevacizumab plus chemotherapy compared with up to 5% in the patients treated with chemotherapy alone. Grade 3-5 reactions have been observed in up to 2.3% of patients treated with bevacizumab plus chemotherapy as compared with  $< 1\%$  with chemotherapy alone (NCI-CTCAE v.3). Major or massive pulmonary haemorrhage/haemoptysis can occur suddenly and up to two thirds of the serious pulmonary haemorrhages resulted in a fatal outcome.

Gastrointestinal haemorrhages, including rectal bleeding and melaena have been reported in colorectal cancer patients, and have been assessed as tumour-associated haemorrhages.

Tumour-associated haemorrhage was also seen rarely in other tumour types and locations, including cases of central nervous system (CNS) bleeding in patients with CNS metastases (see section 4.4).

The incidence of CNS bleeding in patients with untreated CNS metastases receiving bevacizumab has not been prospectively evaluated in randomised clinical trials. In an exploratory retrospective analysis of data from 13 completed randomised trials in patients with various tumour types, 3 patients out of 91 (3.3%) with brain metastases experienced CNS bleeding (all Grade 4) when treated with bevacizumab, compared to 1 case (Grade 5) out of 96 patients (1%) that were not exposed to bevacizumab. In two subsequent

studies in patients with treated brain metastases (which included around 800 patients), one case of Grade 2 CNS haemorrhage was reported in 83 subjects treated with bevacizumab (1.2%) at the time of interim safety analysis (NCI-CTCAE v.3).

Across all clinical trials, mucocutaneous haemorrhage has been seen in up to 50% of bevacizumab-treated patients. These were most commonly NCI-CTCAE v.3 Grade 1 epistaxis that lasted less than 5 minutes, resolved without medical intervention and did not require any changes in the bevacizumab treatment regimen. Clinical safety data suggest that the incidence of minor mucocutaneous haemorrhage (e.g. epistaxis) may be dose-dependent.

There have also been less common reactions of minor mucocutaneous haemorrhage in other locations, such as gingival bleeding or vaginal bleeding.

*Thromboembolism* (see section 4.4)

*Arterial thromboembolism:* An increased incidence of arterial thromboembolic reactions was observed in patients treated with bevacizumab across indications, including cerebrovascular accidents, myocardial infarction, transient ischaemic attacks, and other arterial thromboembolic reactions.

In clinical trials, the overall incidence of arterial thromboembolic reactions ranged up to 3.8% in the bevacizumab containing arms compared with up to 2.1% in the chemotherapy control arms. Fatal outcome was reported in 0.8% of patients receiving bevacizumab compared to 0.5% in patients receiving chemotherapy alone. Cerebrovascular accidents (including transient ischaemic attacks) were reported in up to 2.7% of patients treated with bevacizumab in combination with chemotherapy compared to up to 0.5% of patients treated with chemotherapy alone. Myocardial infarction was reported in up to 1.4% of patients treated with bevacizumab in combination with chemotherapy compared to up to 0.7% of patients treated with chemotherapy alone.

In one clinical trial evaluating bevacizumab in combination with 5-fluorouracil/folinic acid, AVF2192g, patients with metastatic colorectal cancer who were not candidates for treatment with irinotecan were included. In this trial arterial thromboembolic reactions were observed in 11% (11/100) of patients compared to 5.8% (6/104) in the chemotherapy control group.

*Venous thromboembolism:* The incidence of venous thromboembolic reactions in clinical trials was similar in patients receiving bevacizumab in combination with chemotherapy compared to those receiving the control chemotherapy alone. Venous thromboembolic reactions include deep venous thrombosis, pulmonary embolism and thrombophlebitis.

In clinical trials across indications, the overall incidence of venous thromboembolic reactions ranged from 2.8% to 17.3% of bevacizumab-treated patients compared with 3.2% to 15.6% in the control arms.

Grade 3-5 (NCI-CTCAE v.3) venous thromboembolic reactions have been reported in up to 7.8% of patients treated with chemotherapy plus bevacizumab compared with up to 4.9% in patients treated with chemotherapy alone (across indications, excluding persistent, recurrent, or metastatic cervical cancer).

From a clinical trial in patients with persistent, recurrent, or metastatic cervical cancer (study GOG-0240), grade 3-5 venous thromboembolic events have been reported in up to 15.6% of patients treated with bevacizumab in combination with paclitaxel and cisplatin compared with up to 7.0% of patients treated with paclitaxel and cisplatin.

Patients who have experienced a venous thromboembolic reaction may be at higher risk for a recurrence if they receive bevacizumab in combination with chemotherapy versus chemotherapy alone.

*Congestive heart failure (CHF)*

In clinical trials with bevacizumab, congestive heart failure (CHF) was observed in all cancer indications studied to date, but occurred predominantly in patients with metastatic breast cancer. In four phase III trials (AVF2119g, E2100, BO17708 and AVF3694g) in patients with metastatic breast cancer CHF Grade 3 (NCI-CTCAE v.3) or higher was reported in up to 3.5% of patients treated with bevacizumab in combination with chemotherapy compared with up to 0.9% in the control arms. For patients in study AVF3694g who received anthracyclines concomitantly with bevacizumab, the incidences of Grade 3 or higher CHF for the respective bevacizumab and control arms were similar to those in the other studies in metastatic breast cancer: 2.9% in the anthracycline + bevacizumab arm and 0% in the anthracycline + placebo arm. In addition, in study AVF3694g the incidences of all Grade CHF were similar between the anthracycline + bevacizumab (6.2%) and the anthracycline + placebo arms (6.0%).



Most patients who developed CHF during mBC trials showed improved symptoms and/or left ventricular function following appropriate medical therapy.

In most clinical trials of bevacizumab, patients with pre-existing CHF of NYHA (New York Heart Association) II-IV were excluded, therefore, no information is available on the risk of CHF in this population.

Prior anthracyclines exposure and/or prior radiation to the chest wall may be possible risk factors for the development of CHF.

An increased incidence of CHF has been observed in a clinical trial of patients with diffuse large B-cell lymphoma when receiving bevacizumab with a cumulative doxorubicin dose greater than 300 mg/m<sup>2</sup>. This phase III clinical trial compared rituximab/cyclophosphamide/doxorubicin/vincristine/prednisone (R-CHOP) plus bevacizumab to R-CHOP without bevacizumab. While the incidence of CHF was, in both arms, above that previously observed for doxorubicin therapy, the rate was higher in the R-CHOP plus bevacizumab arm. These results suggest that close clinical observation with appropriate cardiac assessments should be considered for patients exposed to cumulative doxorubicin doses greater than 300 mg/m<sup>2</sup> when combined with bevacizumab.

*Hypersensitivity reactions (including anaphylactic shock)/infusion reactions* (see section 4.4 and *Post-marketing experience* below)

In some clinical trials anaphylactic and anaphylactoid-type reactions were reported more frequently in patients receiving bevacizumab in combination with chemotherapy than with chemotherapy alone. The incidence of these reactions in some clinical trials of bevacizumab is common (up to 5% in bevacizumab-treated patients).

#### *Infections*

From a clinical trial in patients with persistent, recurrent, or metastatic cervical cancer (study GOG-0240), grade 3-5 infections have been reported in up to 24% of patients treated with bevacizumab in combination with paclitaxel and topotecan compared with up to 13% of patients treated with paclitaxel and topotecan.

#### *Ovarian failure/fertility* (see sections 4.4 and 4.6)

In NSABP C-08, a phase III trial of bevacizumab in adjuvant treatment of patients with colon cancer, the incidence of new cases of ovarian failure, defined as amenorrhoea lasting 3 or more months, FSH level  $\geq$  30 mIU/mL and a negative serum  $\beta$ -HCG pregnancy test, has been evaluated in 295 premenopausal women. New cases of ovarian failure were reported in 2.6% patients in the mFOLFOX-6 group compared to 39% in the mFOLFOX-6 + bevacizumab group. After discontinuation of bevacizumab treatment, ovarian function recovered in 86.2% of these evaluable women. Long term effects of the treatment with bevacizumab on fertility are unknown.

#### *Laboratory abnormalities*

Decreased neutrophil count, decreased white blood cell count and presence of urine protein may be associated with Versavo treatment.

Across clinical trials, the following Grade 3 and 4 (NCI-CTCAE v.3) laboratory abnormalities occurred in patients treated with bevacizumab with at least a 2% difference compared to the corresponding control groups: hyperglycaemia, decreased haemoglobin, hypokalaemia, hyponatraemia, decreased white blood cell count, increased international normalised ratio (INR).

Clinical trials have shown that transient increases in serum creatinine (ranging between 1.5-1.9 times baseline level), both with and without proteinuria, are associated with the use of bevacizumab. The observed increase in serum creatinine was not associated with a higher incidence of clinical manifestations of renal impairment in patients treated with bevacizumab.

#### Other special populations

##### *Elderly patients*

In randomised clinical trials, age > 65 years was associated with an increased risk of developing arterial thromboembolic reactions, including cerebrovascular accidents (CVAs), transient ischaemic attacks (TIAs) and myocardial infarctions (MIs). Other reactions with a higher frequency seen in patients over 65 were Grade 3-4 leucopenia and thrombocytopenia (NCI-CTCAE v.3); and all Grade neutropenia, diarrhoea, nausea, headache and fatigue as compared to those aged  $\leq$  65 years when treated with bevacizumab (see sections 4.4 and 4.8 under *Thromboembolism*). In one clinical trial, the incidence of

hypertension of grade  $\geq 3$  was two fold higher in patients aged  $> 65$  years than in the younger age group ( $<65$  years). In a study of platinum-resistant recurrent ovarian cancer patients, alopecia, mucosal inflammation, peripheral sensory neuropathy, proteinuria and hypertension were also reported and occurred at a rate at least 5% higher in the CT + BV arm for bevacizumab-treated patients  $\geq 65$  years of age compared with bevacizumab-treated patients aged  $< 65$  years.

No increase in the incidence of other reactions, including gastrointestinal perforation, wound healing complications, congestive heart failure, and haemorrhage was observed in elderly patients ( $> 65$  years) receiving bevacizumab as compared to those aged  $\leq 65$  years treated with bevacizumab.

#### *Paediatric population*

The safety and efficacy of Versavo in children less than 18 years old have not been established.

In study BO25041 of bevacizumab added to postoperative radiation therapy (RT) with concomitant and adjuvant temozolomide in paediatric patients with newly diagnosed supratentorial, infratentorial, cerebellar, or peduncular high-grade glioma, the safety profile was comparable with that observed in other tumour types in adults treated with bevacizumab.

In study BO20924 of bevacizumab with current standard of care in rhabdomyosarcoma and non-rhabdomyosarcoma soft tissue sarcoma, the safety profile of bevacizumab treated children was comparable with that observed in adults treated with bevacizumab.

Versavo is not approved for use in patients under the age of 18 years. In published literature reports, cases of non-mandibular osteonecrosis have been observed in patients under the age of 18 years treated with bevacizumab.

#### Post-marketing experience

**Table 3 Adverse reactions reported in post-marketing setting**

| <i>System organ class (SOC)</i>                        | <i>Reactions (frequency*)</i>                                                                                                                                                                                                                                                                                                                                                                       |
|--------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Infections and Infestations</i>                     | Necrotising fasciitis, usually secondary to wound healing complications, gastrointestinal perforation or fistula formation (rare) (see also section 4.4)                                                                                                                                                                                                                                            |
| <i>Immune system disorders</i>                         | Hypersensitivity reactions and infusion reactions (common); with the following possible co-manifestations: dyspnoea/difficulty breathing, flushing/redness/rash, hypotension or hypertension, oxygen desaturation, chest pain, rigors and nausea/vomiting (see also section 4.4 and <i>Hypersensitivity reactions/infusion reactions</i> above)<br>Anaphylactic shock (rare) (see also section 4.4) |
| <i>Nervous system disorders</i>                        | Hypertensive encephalopathy (very rare) (see also section 4.4 and <i>Hypertension</i> in section 4.8)<br>Posterior Reversible Encephalopathy Syndrome (PRES), (rare) (see also section 4.4)                                                                                                                                                                                                         |
| <i>Vascular disorders</i>                              | Renal thrombotic microangiopathy, which may be clinically manifested as proteinuria (not known) with or without concomitant sunitinib use. For further information on proteinuria see section 4.4 and <i>Proteinuria</i> in section 4.8.                                                                                                                                                            |
| <i>Respiratory, thoracic and mediastinal disorders</i> | Nasal septum perforation (not known)<br>Pulmonary hypertension (not known)<br>Dysphonia (common)                                                                                                                                                                                                                                                                                                    |
| <i>Gastrointestinal disorders</i>                      | Gastrointestinal ulcer (not known)                                                                                                                                                                                                                                                                                                                                                                  |
| <i>Hepatobiliary disorders</i>                         | Gall bladder perforation (not known)                                                                                                                                                                                                                                                                                                                                                                |
| <i>Musculoskeletal and connective tissue disorders</i> | Cases of Osteonecrosis of the Jaw (ONJ) have been reported in patients treated with bevacizumab, most of which occurred in patients who had identified risk factors for ONJ, in particular exposure to intravenous bisphosphonates and/or a history of dental                                                                                                                                       |

|                                                   |                                                                                                                                                                        |
|---------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                   | disease requiring invasive dental procedures (see also section 4.4)                                                                                                    |
|                                                   | Cases of non-mandibular osteonecrosis have been observed in bevacizumab treated paediatric patients (see section 4.8, Paediatric population).                          |
| <i>Congenital, familial, and genetic disorder</i> | Cases of foetal abnormalities in women treated with bevacizumab alone or in combination with known embryotoxic chemotherapeutics have been observed (see section 4.6 ) |

\* if specified, the frequency has been derived from clinical trial data

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions to National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers→Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]”

## 4.9 Overdose

The highest dose tested in humans (20 mg/kg of body weight, intravenous every 2 weeks) was associated with severe migraine in several patients.

# 5 PHARMACOLOGICAL PROPERTIES

## 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antineoplastic and immunomodulating agents, antineoplastic agents, other antineoplastic agents, monoclonal antibodies, ATC code: L01FG01

Versavo is a biosimilar medicinal product.

#### Mechanism of action

Bevacizumab binds to vascular endothelial growth factor (VEGF), the key driver of vasculogenesis and angiogenesis, and thereby inhibits the binding of VEGF to its receptors, Flt-1 (VEGFR-1) and KDR (VEGFR-2), on the surface of endothelial cells. Neutralising the biological activity of VEGF regresses the vascularisation of tumours, normalises remaining tumour vasculature, and inhibits the formation of new tumour vasculature, thereby inhibiting tumour growth.

#### Pharmacodynamic effects

Administration of bevacizumab or its parental murine antibody to xenotransplant models of cancer in nude mice resulted in extensive anti-tumour activity in human cancers, including colon, breast, pancreas and prostate. Metastatic disease progression was inhibited and microvascular permeability was reduced.

#### Clinical efficacy

##### Metastatic carcinoma of the colon or rectum (mCRC)

The safety and efficacy of the recommended dose (5 mg/kg of body weight every two weeks) in metastatic carcinoma of the colon or rectum were studied in three randomised, active-controlled clinical trials in combination with fluoropyrimidine-based first-line chemotherapy. Bevacizumab was combined with two chemotherapy regimens:

- AVF2107g: A weekly schedule of irinotecan/bolus 5-fluorouracil/folinic acid (IFL) for a total of 4 weeks of each 6 week-cycle (Saltz regimen).
- AVF0780g: In combination with bolus 5-fluorouracil/folinic acid (5-FU/FA) for a total of 6 weeks of each 8 week-cycle (Roswell Park regimen).
- AVF2192g: In combination with bolus 5-FU/FA for a total of 6 weeks of each 8 week-cycle (Roswell Park regimen) in patients who were not optimal candidates for first-line irinotecan treatment.

Three additional studies with bevacizumab have been conducted in mCRC patients: first-line (NO16966), second-line with no previous bevacizumab treatment (E3200), and second-line with previous bevacizumab treatment following disease progression in first-line (ML18147). In these studies, bevacizumab was administered at the following dosing regimens in combination with FOLFOX-4 (5-FU/LV/oxaliplatin), XELOX (capecitabine/oxaliplatin), and fluoropyrimidine/irinotecan and fluoropyrimidine/oxaliplatin:

- NO16966: Bevacizumab 7.5 mg/kg of body weight every 3 weeks in combination with oral capecitabine and intravenous oxaliplatin (XELOX) or bevacizumab 5 mg/kg every 2 weeks in combination with leucovorin plus 5-fluorouracil bolus, followed by 5-fluorouracil infusion, with intravenous oxaliplatin (FOLFOX-4).
- E3200: Bevacizumab 10 mg/kg of body weight every 2 weeks in combination with leucovorin and 5-fluorouracil bolus, followed by 5-fluorouracil infusion, with intravenous oxaliplatin (FOLFOX-4) in bevacizumab-naïve patients.
- ML18147: Bevacizumab 5.0 mg/kg of body weight every 2 weeks or bevacizumab 7.5 mg/kg of body weight every 3 weeks in combination with fluoropyrimidine/irinotecan or fluoropyrimidine/oxaliplatin in patients with disease progression following first-line treatment with bevacizumab. Use of irinotecan- or oxaliplatin-containing regimen was switched depending on first-line usage of either oxaliplatin or irinotecan.

#### AVF2107g

This was a phase III randomised, double-blind, active-controlled clinical trial evaluating bevacizumab in combination with IFL as first-line treatment for metastatic carcinoma of the colon or rectum. Eight hundred and thirteen patients were randomised to receive IFL + placebo (Arm 1) or IFL + bevacizumab (5 mg/kg every 2 weeks, Arm 2). A third group of 110 patients received bolus 5-FU/FA + bevacizumab (Arm 3). Enrolment in Arm 3 was discontinued, as pre-specified, once safety of bevacizumab with the IFL regimen was established and considered acceptable. All treatments were continued until disease progression. The overall mean age was 59.4 years; 56.6% of patients had an ECOG performance status of 0, 43% had a value of 1 and 0.4% had a value of 2. 15.5% had received prior radiotherapy and 28.4% prior chemotherapy.

The primary efficacy variable of the trial was overall survival. The addition of bevacizumab to IFL resulted in statistically significant increases in overall survival, progression-free survival and overall response rate (see Table 4). The clinical benefit, as measured by overall survival, was seen in all pre-specified patient subgroups, including those defined by age, sex, performance status, location of primary tumour, number of organs involved and duration of metastatic disease.

The efficacy results of bevacizumab in combination with IFL-chemotherapy are displayed in Table 4.

**Table 4 Efficacy results for trial AVF2107g**

|                           | AVF2107g                     |                                         |
|---------------------------|------------------------------|-----------------------------------------|
|                           | Arm 1<br>IFL + placebo       | Arm 2<br>IFL + bevacizumab <sup>a</sup> |
| Number of patients        | 411                          | 402                                     |
| Overall survival          |                              |                                         |
| Median time (months)      | 15.6                         | 20.3                                    |
| 95% CI                    | 14.29 – 16.99                | 18.46 – 24.18                           |
| Hazard ratio <sup>b</sup> | 0.660<br>(p-value = 0.00004) |                                         |
| Progression-free survival |                              |                                         |
| Median time (months)      | 6.2                          | 10.6                                    |
| Hazard ratio              | 0.54<br>(p-value < 0.0001)   |                                         |
| Overall response rate     |                              |                                         |
| Rate (%)                  | 34.8                         | 44.8                                    |
|                           | (p-value = 0.0036)           |                                         |

<sup>a</sup> 5 mg/kg every 2 weeks.

<sup>b</sup> Relative to control arm.

Among the 110 patients randomised to Arm 3 (5-FU/FA + bevacizumab) prior to discontinuation of this arm, the median overall survival was 18.3 months and the median progression free survival was 8.8 months.

#### *AVF2192g*

This was a phase II randomised, double-blind, active-controlled clinical trial evaluating the efficacy and safety of bevacizumab in combination with 5-FU/FA as first-line treatment for metastatic colorectal cancer in patients who were not optimal candidates for first-line irinotecan treatment. One hundred and five patients were randomised to 5-FU/FA + placebo arm and 104 patients to 5-FU/FA + bevacizumab (5 mg/kg every 2 weeks) arm. All treatments were continued until disease progression. The addition of bevacizumab 5 mg/kg every two weeks to 5-FU/FA resulted in higher objective response rates, significantly longer progression-free survival, and a trend in longer survival as compared to 5-FU/FA chemotherapy alone.

#### *AVF0780g*

This was a phase II randomised, active-controlled, open-labelled clinical trial investigating bevacizumab in combination with 5-FU/FA as first-line treatment of metastatic colorectal cancer. The median age was 64 years. 19% of the patients had received prior chemotherapy and 14% prior radiotherapy. Seventy-one patients were randomised to receive bolus 5-FU/FA or 5-FU/FA + bevacizumab (5 mg/kg every 2 weeks). A third group of 33 patients received bolus 5-FU/FA + bevacizumab (10 mg/kg every 2 weeks). Patients were treated until disease progression. The primary endpoints of the trial were objective response rate and progression-free survival. The addition of bevacizumab 5 mg/kg every two weeks to 5-FU/FA resulted in higher objective response rates, longer progression-free survival, and a trend in longer survival, compared with 5-FU/FA chemotherapy alone (see Table 5). These efficacy data are consistent with the results from trial AVF2107g.

The efficacy data from trials AVF0780g and AVF2192g investigating bevacizumab in combination with 5-FU/FA-chemotherapy are summarised in Table 5.

**Table 5 Efficacy results for trials AVF0780g and AVF2192g**

|                           | AVF0780g   |                                    |                                    | AVF2192g          |                       |
|---------------------------|------------|------------------------------------|------------------------------------|-------------------|-----------------------|
|                           | 5-FU/FA    | 5-FU/FA + bevacizumab <sup>a</sup> | 5-FU/FA + bevacizumab <sup>b</sup> | 5-FU/FA + placebo | 5-FU/FA + bevacizumab |
| Number of patients        | 36         | 35                                 | 33                                 | 105               | 104                   |
| Overall survival          |            |                                    |                                    |                   |                       |
| Median time (months)      | 13.6       | 17.7                               | 15.2                               | 12.9              | 16.6                  |
| 95% CI                    |            |                                    |                                    | 10.35 - 16.95     | 13.63 - 19.32         |
| Hazard ratio <sup>c</sup> | -          | 0.52                               | 1.01                               |                   | 0.79                  |
| p-value                   |            | 0.073                              | 0.978                              |                   | 0.16                  |
| Progression-free survival |            |                                    |                                    |                   |                       |
| Median time (months)      | 5.2        | 9.0                                | 7.2                                | 5.5               | 9.2                   |
| Hazard ratio              |            | 0.44                               | 0.69                               |                   | 0.5                   |
| p-value                   | -          | 0.0049                             | 0.217                              |                   | 0.0002                |
| Overall response rate     |            |                                    |                                    |                   |                       |
| Rate (percent)            | 16.7       | 40.0                               | 24.2                               | 15.2              | 26                    |
| 95% CI                    | 7.0 - 33.5 | 24.4 - 57.8                        | 11.7 - 42.6                        | 9.2 - 23.9        | 18.1 - 35.6           |
| p-value                   |            | 0.029                              | 0.43                               |                   | 0.055                 |
| Duration of response      |            |                                    |                                    |                   |                       |
| Median time (months)      | NR         | 9.3                                | 5.0                                | 6.8               | 9.2                   |

|                           |          |          |           |             |              |
|---------------------------|----------|----------|-----------|-------------|--------------|
| 25–75 percentile (months) | 5.5 - NR | 6.1 - NR | 3.8 – 7.8 | 5.59 - 9.17 | 5.88 - 13.01 |
|---------------------------|----------|----------|-----------|-------------|--------------|

<sup>a</sup> 5 mg/kg every 2 weeks.

<sup>b</sup> 10 mg/kg every 2 weeks.

<sup>c</sup> Relative to control arm.

NR = not reached.

#### NO16966

This was a phase III randomised, double-blind (for bevacizumab), clinical trial investigating bevacizumab 7.5 mg/kg in combination with oral capecitabine and IV oxaliplatin (XELOX), administered on a 3-weekly schedule; or bevacizumab 5 mg/kg in combination with leucovorin with 5-fluorouracil bolus, followed by 5-fluorouracil infusion, with IV oxaliplatin (FOLFOX-4), administered on a 2-weekly schedule. The trial contained two parts: an initial unblinded 2-arm part (Part I) in which patients were randomised to two different treatment groups (XELOX and FOLFOX-4) and a subsequent 2 x 2 factorial 4-arm part (Part II) in which patients were randomised to four treatment groups (XELOX + placebo, FOLFOX-4 + placebo, XELOX + bevacizumab, FOLFOX-4 + bevacizumab). In Part II, treatment assignment was double-blind with respect to bevacizumab.

Approximately 350 patients were randomised into each of the 4 trial arms in the Part II of the trial.

**Table 6 Treatment regimens in trial NO16966 (mCRC)**

|                                                                 | Treatment                                     | Starting dose                                                                                                                   | Schedule                                                                                                                      |
|-----------------------------------------------------------------|-----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------|
| FOLFOX-4<br>or<br>FOLFOX-4<br>+ bevacizumab                     | Oxaliplatin<br>Leucovorin<br>+ 5-Fluorouracil | 85 mg/m <sup>2</sup> IV 2 h<br>200 mg/m <sup>2</sup> IV 2 h<br>400 mg/m <sup>2</sup> IV bolus, 600 mg/m <sup>2</sup> IV<br>22 h | Oxaliplatin on<br>day 1<br>Leucovorin on<br>day 1 and 2<br>5-fluorouracil<br>IV<br>bolus/infusion,<br>each on days 1<br>and 2 |
|                                                                 | Placebo or bevacizumab                        | 5 mg/kg IV 30-90 min                                                                                                            | Day 1, prior to<br>FOLFOX-4,<br>every 2 weeks                                                                                 |
| XELOX<br>or<br>XELOX<br>+ bevacizumab                           | Oxaliplatin<br>Capecitabine                   | 130 mg/m <sup>2</sup> IV 2 h<br>1000 mg/m <sup>2</sup> oral bid                                                                 | Oxaliplatin on<br>day 1<br>Capecitabine<br>oral bid for 2<br>weeks<br>(followed by 1<br>week off<br>treatment)                |
|                                                                 | Placebo or bevacizumab                        | 7.5 mg/kg IV 30-90 min                                                                                                          | Day 1, prior to<br>XELOX, q 3<br>weeks                                                                                        |
| 5-Fluorouracil: IV bolus injection immediately after leucovorin |                                               |                                                                                                                                 |                                                                                                                               |

The primary efficacy parameter of the trial was the duration of progression-free survival. In this trial, there were two primary objectives: to show that XELOX was non-inferior to FOLFOX-4 and to show that bevacizumab in combination with FOLFOX-4 or XELOX chemotherapy was superior to chemotherapy alone. Both co-primary objectives were met:

- Non-inferiority of the XELOX-containing arms compared with the FOLFOX-4-containing arms in the overall comparison was demonstrated in terms of progression-free survival and overall survival in the eligible per-protocol population.
- Superiority of the bevacizumab-containing arms versus the chemotherapy alone arms in the overall comparison was demonstrated in terms of progression-free survival in the ITT population (Table 7).

Secondary PFS analyses, based on 'on-treatment'-based response assessments, confirmed the significantly superior clinical benefit for patients treated with bevacizumab (analyses shown in Table 7), consistent with the statistically significant benefit observed in the pooled analysis.

**Table 7 Key efficacy results for the superiority analysis (ITT population, trial NO16966)**

| Endpoint (months)                               | FOLFOX-4<br>XELOX + placebo<br>(n=701) | or<br>FOLFOX-4<br>XELOX<br>bevacizumab<br>(n=699) | or<br>+ | P-value  |
|-------------------------------------------------|----------------------------------------|---------------------------------------------------|---------|----------|
| Primary endpoint                                |                                        |                                                   |         |          |
| Median PFS**                                    | 8.0                                    | 9.4                                               |         | 0.0023   |
| Hazard ratio (97.5% CI) <sup>a</sup>            | 0.83 (0.72–0.95)                       |                                                   |         |          |
| Secondary endpoints                             |                                        |                                                   |         |          |
| Median PFS (on treatment)**                     | 7.9                                    | 10.4                                              |         | < 0.0001 |
| Hazard ratio (97.5% CI)                         | 0.63 (0.52-0.75)                       |                                                   |         |          |
| Overall response rate<br>(invest. assessment)** | 49.2 %                                 | 46.5 %                                            |         |          |
| Median overall survival*                        | 19.9                                   | 21.2                                              |         | 0.0769   |
| Hazard ratio (97.5% CI)                         | 0.89 (0.76-1.03)                       |                                                   |         |          |

\* Overall survival analysis at clinical cut-off 31 January 2007

\*\* Primary analysis at clinical cut-off 31 January 2006

<sup>a</sup> relative to control arm

In the FOLFOX treatment subgroup, the median PFS was 8.6 months in placebo and 9.4 months in bevacizumab treated patients, HR = 0.89, 97.5% CI = [0.73; 1.08]; p-value = 0.1871, the corresponding results in the XELOX treatment subgroup being 7.4 vs. 9.3 months, HR = 0.77, 97.5% CI = [0.63; 0.94]; p-value = 0.0026.

The median overall survival was 20.3 months in placebo and 21.2 months in bevacizumab treated patients in the FOLFOX treatment subgroup, HR=0.94, 97.5% CI = [0.75; 1.16]; p-value = 0.4937, the corresponding results in the XELOX, treatment subgroup being 19.2 vs. 21.4 months, HR = 0.84, 97.5% CI = [0.68; 1.04]; p-value = 0.0698.

#### *ECOG E3200*

This was a phase III randomised, active-controlled, open-label trial investigating bevacizumab 10 mg/kg in combination with leucovorin with 5-fluorouracil bolus and then 5-fluorouracil infusional, with IV oxaliplatin (FOLFOX-4), administered on a 2-weekly schedule in previously-treated patients (second line) with advanced colorectal cancer. In the chemotherapy arms, the FOLFOX-4 regimen used the same doses and schedule as shown in Table 6 for trial NO16966.

The primary efficacy parameter of the trial was overall survival, defined as the time from randomisation to death from any cause. Eight hundred and twenty-nine patients were randomised (292 FOLFOX-4, 293 bevacizumab + FOLFOX-4 and 244 bevacizumab monotherapy). The addition of bevacizumab to FOLFOX-4 resulted in a statistically significant prolongation of survival. Statistically significant improvements in progression-free survival and objective response rate were also observed (see Table 8).

**Table 8 Efficacy results for trial E3200**

|                    | E3200    |                                        |
|--------------------|----------|----------------------------------------|
|                    | FOLFOX-4 | FOLFOX-4<br>bevacizumab <sup>a</sup> + |
| Number of patients | 292      | 293                                    |
| Overall survival   |          |                                        |
| Median (months)    | 10.8     | 13.0                                   |

|                           |                             |               |
|---------------------------|-----------------------------|---------------|
| 95% CI                    | 10.12 – 11.86               | 12.09 – 14.03 |
| Hazard ratio <sup>b</sup> | 0.751<br>(p-value = 0.0012) |               |
| Progression-free survival |                             |               |
| Median (months)           | 4.5                         | 7.5           |
| Hazard ratio              | 0.518<br>(p-value < 0.0001) |               |
| Objective response rate   |                             |               |
| Rate                      | 8.6%                        | 22.2%         |
|                           | (p-value < 0.0001)          |               |

<sup>a</sup> 10 mg/kg every 2 weeks

<sup>b</sup> Relative to control arm

No significant difference was observed in the duration of overall survival between patients who received bevacizumab monotherapy compared to patients treated with FOLFOX-4. Progression-free survival and objective response rate were inferior in the bevacizumab monotherapy arm compared to the FOLFOX-4 arm.

#### ML18147

This was a Phase III randomised, controlled, open-label trial investigating bevacizumab 5.0 mg/kg every 2 weeks or 7.5 mg/kg every 3 weeks in combination with fluoropyrimidine-based chemotherapy versus fluoropyrimidine-based chemotherapy alone in patients with mCRC who have progressed on a first-line bevacizumab-containing regimen.

Patients with histologically confirmed mCRC and disease progression were randomised 1:1 within 3 months after discontinuation of bevacizumab first-line therapy to receive fluoropyrimidine/oxaliplatin- or fluoropyrimidine/irinotecan-based chemotherapy (chemotherapy switched depending on first-line chemotherapy) with or without bevacizumab. Treatment was given until progressive disease or unacceptable toxicity. The primary outcome measure was overall survival defined as the time from randomisation until death from any cause.

A total of 820 patients were randomised. The addition of bevacizumab to fluoropyrimidine-based chemotherapy resulted in a statistically significant prolongation of survival in patients with mCRC who have progressed on a first-line bevacizumab-containing regimen (ITT = 819) (see Table 9).

**Table 9 Efficacy Results for Study ML18147 (ITT population)**

|                                         |                                                                                         |                                                                                                                       |
|-----------------------------------------|-----------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------|
|                                         | ML18147                                                                                 |                                                                                                                       |
|                                         | fluoropyrimidine/irinotecan<br>or<br>fluoropyrimidine/oxaliplatin<br>based chemotherapy | fluoropyrimidine/irinotecan<br>or<br>fluoropyrimidine/oxaliplatin<br>based chemotherapy +<br>bevacizumab <sup>a</sup> |
| Number of Patients                      | 410                                                                                     | 409                                                                                                                   |
| <b><u>Overall Survival</u></b>          |                                                                                         |                                                                                                                       |
| Median (months)                         | 9.8                                                                                     | 11.2                                                                                                                  |
| Hazard ratio (95% confidence interval)  | 0.81 (0.69, 0.94)<br>(p-value = 0.0062)                                                 |                                                                                                                       |
| <b><u>Progression-Free Survival</u></b> |                                                                                         |                                                                                                                       |
| Median (months)                         | 4.1                                                                                     | 5.7                                                                                                                   |



|                                             |                                         |      |
|---------------------------------------------|-----------------------------------------|------|
| Hazard ratio (95% confidence interval)      | 0.68 (0.59, 0.78)<br>(p-value < 0.0001) |      |
| <b><u>Objective Response Rate (ORR)</u></b> |                                         |      |
| Patients included in analysis               | 406                                     | 404  |
| Rate                                        | 3.9%                                    | 5.4% |
|                                             | (p-value = 0.3113)                      |      |

<sup>a</sup> 5.0 mg/kg every 2 weeks or 7.5 mg/kg every 3 weeks

Statistically significant improvements in progression-free survival were also observed. Objective response rate was low in both treatment arms and the difference was not significant.

Study E3200 used a 5 mg/kg/week equivalent dose of bevacizumab in bevacizumab-naïve patients, while study ML18147 used a 2.5 mg/kg/week equivalent dose of bevacizumab in bevacizumab-pretreated patients. A cross-trial comparison of the efficacy and safety data is limited by differences between these studies, most notably in patient populations, previous bevacizumab exposure and chemotherapy regimens. Both the 5 mg/kg/week and 2.5 mg/kg/week equivalent doses of bevacizumab provided a statistically significant benefit with regards to OS (HR 0.751 in study E3200; HR 0.81 in study ML18147) and PFS (HR 0.518 in study E3200; HR 0.68 in study ML18147). In terms of safety, there was a higher overall incidence of Grade 3-5 AEs in study E3200 relative to study ML18147.

#### Metastatic breast cancer (mBC)

Two large Phase III trials were designed to investigate the treatment effect of bevacizumab in combination with two individual chemotherapy agents, as measured by the primary endpoint of PFS. A clinically meaningful and statistically significant improvement in PFS was observed in both trials.

Summarised below are PFS results for the individual chemotherapy agents included in the indication:

- Study E2100 (paclitaxel)
- Median PFS increase 5.6 months, HR 0.421 (p < 0.0001, 95% CI 0.343; 0.516)
- Study AVF3694g (capecitabine)
- Median PFS increase 2.9 months, HR 0.69 (p = 0.0002, 95% CI 0.56; 0.84)

Further details of each study and the results are provided below.

#### ECOG E2100

Trial E2100 was an open-label, randomised, active controlled, multicentre clinical trial evaluating bevacizumab in combination with paclitaxel for locally recurrent or metastatic breast cancer in patients who had not previously received chemotherapy for locally recurrent and metastatic disease. Patients were randomised to paclitaxel alone (90 mg/m<sup>2</sup> IV over 1 hour once weekly for three out of four weeks) or in combination with bevacizumab (10 mg/kg IV infusion every two weeks). Prior hormonal therapy for the treatment of metastatic disease was allowed. Adjuvant taxane therapy was allowed only if it was completed at least 12 months prior to trial entry. Of the 722 patients in the trial, the majority of patients had HER2-negative disease (90%), with a small number of patients with unknown (8%) or confirmed HER2-positive status (2%), who had previously been treated with or were considered unsuitable for trastuzumab therapy. Furthermore, 65% of patients had received adjuvant chemotherapy including 19% prior taxanes and 49% prior anthracyclines. Patients with central nervous system metastases, including previously treated or resected brain lesions, were excluded.

In trial E2100, patients were treated until disease progression. In situations where early discontinuation of chemotherapy was required, treatment with bevacizumab as a single agent continued until disease progression. The patient characteristics were similar across the trial arms. The primary endpoint of this trial was progression free survival (PFS), based on trial investigators' assessment of disease progression. In addition, an independent review of the primary endpoint was also conducted. The results of this trial are presented in Table 10.

**Table 10 Trial E2100 efficacy results**

| Progression-free survival                             |                          |                                       |                         |                                       |
|-------------------------------------------------------|--------------------------|---------------------------------------|-------------------------|---------------------------------------|
|                                                       | Investigator assessment* |                                       | IRF assessment          |                                       |
|                                                       | Paclitaxel<br>(n=354)    | Paclitaxel/<br>bevacizumab<br>(n=368) | Paclitaxel<br>(n=354)   | Paclitaxel/<br>bevacizumab<br>(n=368) |
| Median PFS (months)                                   | 5.8                      | 11.4                                  | 5.8                     | 11.3                                  |
| HR<br>(95% CI)                                        | 0.421<br>(0.343; 0.516)  |                                       | 0.483<br>(0.385; 0.607) |                                       |
| p-value                                               | < 0.0001                 |                                       | < 0.0001                |                                       |
| Response rates (for patients with measurable disease) |                          |                                       |                         |                                       |
|                                                       | Investigator assessment  |                                       | IRF assessment          |                                       |
|                                                       | Paclitaxel<br>(n=273)    | Paclitaxel/<br>bevacizumab<br>(n=252) | Paclitaxel<br>(n=243)   | Paclitaxel/<br>bevacizumab<br>(n=229) |
| % pts with objective<br>response                      | 23.4                     | 48.0                                  | 22.2                    | 49.8                                  |
| p-value                                               | < 0.0001                 |                                       | < 0.0001                |                                       |

\* primary analysis

| Overall survival   |                         |                                    |
|--------------------|-------------------------|------------------------------------|
|                    | Paclitaxel<br>(n=354)   | Paclitaxel/ bevacizumab<br>(n=368) |
| Median OS (months) | 24.8                    | 26.5                               |
| HR<br>(95% CI)     | 0.869<br>(0.722; 1.046) |                                    |
| p-value            | 0.1374                  |                                    |

The clinical benefit of bevacizumab as measured by PFS was seen in all pre-specified subgroups tested (including disease-free interval, number of metastatic sites, prior receipt of adjuvant chemotherapy and oestrogen receptor (ER) status).

#### *AVF3694g*

Study AVF3694g was a Phase III, multicentre, randomised, placebo-controlled trial designed to evaluate the efficacy and safety of bevacizumab in combination with chemotherapy compared to chemotherapy plus placebo as first-line treatment for patients with HER2-negative metastatic or locally recurrent breast cancer.

Chemotherapy was chosen at the investigator's discretion prior to randomisation in a 2:1 ratio to receive either chemotherapy plus bevacizumab or chemotherapy plus placebo. The choices of chemotherapy included capecitabine, taxane (protein-bound paclitaxel, docetaxel), and anthracycline-based agents (doxorubicin/ cyclophosphamide, epirubicin/ cyclophosphamide, 5-fluorouracil/ doxorubicin/ cyclophosphamide, 5-fluorouracil/epirubicin/cyclophosphamide) given every three weeks (q3w). Bevacizumab or placebo was administered at a dose of 15 mg/kg q3w.

This study included a blinded treatment phase, an optional open-label post-progression phase, and a survival follow-up phase. During the blinded treatment phase, patients received chemotherapy and medicinal product (bevacizumab or placebo) every 3 weeks until disease progression, treatment-limiting toxicity, or death. On documented disease progression, patients who entered the optional open-label phase could receive open-label bevacizumab together with a wide-range of second line therapies.

Statistical analyses were performed independently for 1) patients who received capecitabine in combination with bevacizumab or placebo; 2) patients who received taxane-based or anthracycline-based chemotherapy in combination with bevacizumab or placebo. The primary endpoint of the study was PFS by investigator assessment. In addition, the primary endpoint was also assessed by an independent review committee (IRC).

The results of this study from the final protocol defined analyses for progression free survival and response rates for the independently powered capecitabine cohort of Study AVF3694g are presented in Table 11 Results from an exploratory overall survival analysis which include an additional 7 months of follow-up (approximately 46% of patients had died) are also presented. The percentage of patients who received bevacizumab in the open-label phase was 62.1% in the capecitabine + placebo arm and 49.9% in the capecitabine + bevacizumab arm.

**Table 11 Efficacy results for study AVF3694g: – Capecitabine<sup>a</sup> and bevacizumab/Placebo (Cap + bevacizumab/PI)**

| Progression-free survival <sup>b</sup>                            |                         |                               |                           |                              |
|-------------------------------------------------------------------|-------------------------|-------------------------------|---------------------------|------------------------------|
|                                                                   | Investigator Assessment |                               | IRC Assessment            |                              |
|                                                                   | Cap + Pl<br>(n=206)     | Cap<br>bevacizumab<br>(n=409) | + Cap + Pl<br>(n=206)     | Cap + bevacizumab<br>(n=409) |
| Median PFS (months)                                               | 5.7                     | 8.6                           | 6.2                       | 9.8                          |
| Hazard ratio vs placebo<br>arm (95% CI)                           | 0.69 (0.56; 0.84)       |                               | 0.68 (0.54; 0.86)         |                              |
| p-value                                                           | 0.0002                  |                               | 0.0011                    |                              |
| Response rate (for patients with measurable disease) <sup>b</sup> |                         |                               |                           |                              |
|                                                                   | Cap + Pl (n=161)        |                               | Cap + bevacizumab (n=325) |                              |
| % pts with objective<br>response                                  | 23.6                    |                               | 35.4                      |                              |
| p-value                                                           | 0.0097                  |                               |                           |                              |
| Overall survival <sup>b</sup>                                     |                         |                               |                           |                              |
| HR<br>(95% CI)                                                    | 0.88 (0.69; 1.13)       |                               |                           |                              |
| p-value (exploratory)                                             | 0.33                    |                               |                           |                              |

<sup>a</sup>1000 mg/m<sup>2</sup> oral twice daily for 14 days administered every 3 weeks

<sup>b</sup>Stratified analysis included all progression and death events except those where non-protocol therapy (NPT) was initiated prior to documented progression; data from those patients were censored at the last tumour assessment prior to starting NPT.

An unstratified analysis of PFS (investigator assessed) was performed that did not censor for non-protocol therapy prior to disease progression. The results of these analyses were very similar to the primary PFS results.

#### Non-small cell lung cancer (NSCLC)

##### *First-line treatment of non-squamous NSCLC in combination with platinum-based chemotherapy*

The safety and efficacy of bevacizumab, in addition to platinum-based chemotherapy, in the first-line treatment of patients with non-squamous non-small cell lung cancer (NSCLC), was investigated in trials E4599 and BO17704. An overall survival benefit has been demonstrated in trial E4599 with a 15 mg/kg/q3wk dose of bevacizumab. Trial BO17704 has demonstrated that both 7.5 mg/kg/q3wk and 15 mg/kg/q3wk bevacizumab doses increase progression free survival and response rate.

##### *E4599*

E4599 was an open-label, randomised, active-controlled, multicentre clinical trial evaluating bevacizumab as first-line treatment of patients with locally advanced (stage IIIb with malignant pleural effusion), metastatic or recurrent NSCLC other than predominantly squamous cell histology.

Patients were randomised to platinum-based chemotherapy (paclitaxel 200 mg/m<sup>2</sup>) and carboplatin AUC = 6.0, both by IV infusion (PC) on day 1 of every 3-week cycle for up to 6 cycles or PC in combination with bevacizumab at a dose of 15 mg/kg IV infusion day 1 of every 3-week cycle. After completion of six cycles of carboplatin-paclitaxel chemotherapy or upon premature discontinuation of chemotherapy, patients on the bevacizumab + carboplatin–paclitaxel arm continued to receive bevacizumab as a single agent every 3 weeks until disease progression. 878 patients were randomised to the two arms.

During the trial, of the patients who received trial treatment, 32.2% (136/422) of patients received 7-12 administrations of bevacizumab and 21.1% (89/422) of patients received 13 or more administrations of bevacizumab.

The primary endpoint was duration of survival. Results are presented in Table 12.

**Table 12 Efficacy results for trial E4599**

|                           | Arm 1<br>Carboplatin/ Paclitaxel         | Arm 2<br>Carboplatin/ Paclitaxel +<br>bevacizumab 15 mg/kg q 3<br>weeks |
|---------------------------|------------------------------------------|-------------------------------------------------------------------------|
| Number of patients        | 444                                      | 434                                                                     |
| Overall survival          |                                          |                                                                         |
| Median (months)           | 10.3                                     | 12.3                                                                    |
| Hazard ratio              | 0.80 (p=0.003)<br>95% CI (0.69; 0.93)    |                                                                         |
| Progression-free survival |                                          |                                                                         |
| Median (months)           | 4.8                                      | 6.4                                                                     |
| Hazard ratio              | 0.65 (p < 0.0001)<br>95% CI (0.56; 0.76) |                                                                         |
| Overall response rate     |                                          |                                                                         |
| Rate (percent)            | 12.9                                     | 29.0 (p < 0.0001)                                                       |

In an exploratory analysis, the extent of bevacizumab benefit on overall survival was less pronounced in the subgroup of patients who did not have adenocarcinoma histology.

#### *BO17704*

Trial BO17704 was a randomised, double-blind phase III trial of bevacizumab in addition to cisplatin and gemcitabine versus placebo, cisplatin and gemcitabine in patients with locally advanced (stage IIIb with supraclavicular lymph node metastases or with malignant pleural or pericardial effusion), metastatic or recurrent non-squamous NSCLC, who had not received prior chemotherapy. The primary endpoint was progression free survival, secondary endpoints for the trial included the duration of overall survival.

Patients were randomised to platinum-based chemotherapy, cisplatin 80 mg/m<sup>2</sup> intravenous infusion on day 1 and gemcitabine 1250 mg/m<sup>2</sup> intravenous infusion on days 1 and 8 of every 3-week cycle for up to 6 cycles (CG) with placebo or CG with bevacizumab at a dose of 7.5 or 15 mg/kg IV infusion day 1 of every 3-week cycle. In the bevacizumab-containing arms, patients could receive bevacizumab as a single-agent every 3 weeks until disease progression or unacceptable toxicity. Trial results show that 94% (277 / 296) of eligible patients went on to receive single agent bevacizumab at cycle 7. A high proportion of patients (approximately 62%) went on to receive a variety of non-protocol specified anti-cancer therapies, which may have impacted the analysis of overall survival.

The efficacy results are presented in Table 13.

**Table 13 Efficacy results for trial BO17704**

|                                            | Cisplatin/Gemcitabine +<br>placebo | Cisplatin/Gemcitabine +<br>bevacizumab 7.5 mg/kg<br>q 3 weeks | Cisplatin/Gemcitabine<br>+ bevacizumab 15<br>mg/kg q 3 weeks |
|--------------------------------------------|------------------------------------|---------------------------------------------------------------|--------------------------------------------------------------|
| Number of patients                         | 347                                | 345                                                           | 351                                                          |
| Progression-free<br>survival               |                                    |                                                               |                                                              |
| Median (months)                            | 6.1                                | 6.7<br>(p=0.0026)                                             | 6.5<br>(p=0.0301)                                            |
| Hazard ratio                               |                                    | 0.75<br>[0.62; 0.91]                                          | 0.82<br>[0.68; 0.98]                                         |
| Best overall response<br>rate <sup>a</sup> | 20.1%                              | 34.1%<br>(p < 0.0001)                                         | 30.4%<br>(p=0.0023)                                          |

\*patients with measurable disease at baseline

|                  |      |                      |                      |
|------------------|------|----------------------|----------------------|
| Overall survival |      |                      |                      |
| Median (months)  | 13.1 | 13.6<br>(p=0.4203)   | 13.4<br>(p=0.7613)   |
| Hazard ratio     |      | 0.93<br>[0.78; 1.11] | 1.03<br>[0.86, 1.23] |

*First-line treatment of non-squamous NSCLC with EGFR activating mutations in combination with erlotinib*

#### JO25567

Study JO25567 was a randomized, open-label, multi-center Phase II study conducted in Japan to evaluate the efficacy and safety of bevacizumab used in addition to erlotinib in patients with non-squamous NSCLC with EGFR activating mutations (exon 19 deletion or exon 21 L858R mutation) who had not received prior systemic therapy for Stage IIIB/IV or recurrent disease.

The primary endpoint was progression-free survival (PFS) based on independent review assessment. Secondary endpoints included overall survival, response rate, disease control rate, duration of response, and safety.

EGFR mutation status was determined for each patient prior to patient screening and 154 patients were randomised to receive either erlotinib + bevacizumab (erlotinib 150 mg oral daily + bevacizumab [15 mg/kg IV every 3 weeks]) or erlotinib monotherapy (150 mg oral daily) until disease progression (PD) or unacceptable toxicity. In the absence of PD, discontinuation of one component of study treatment in the erlotinib + bevacizumab arm did not lead to discontinuation of the other component of study treatment as specified in the study protocol.

The efficacy results of the study are presented in Table 14.

**Table 14 Efficacy results for study JO25567**

|                                              | <b>Erlotinib</b><br>N = 77 <sup>#</sup> | <b>Erlotinib +<br/>Bevacizumab</b><br>N = 75 <sup>#</sup> |
|----------------------------------------------|-----------------------------------------|-----------------------------------------------------------|
| <b>PFS<sup>^</sup></b> (months)              |                                         |                                                           |
| Median                                       | 9.7                                     | 16.0                                                      |
| HR (95% CI)                                  | 0.54 (0.36; 0.79)                       |                                                           |
| p-value                                      | 0.0015                                  |                                                           |
| <b>Overall Response Rate</b>                 |                                         |                                                           |
| Rate (n)                                     | 63.6% (49)                              | 69.3% (52)                                                |
| p-value                                      | 0.4951                                  |                                                           |
| <b>Overall Survival<sup>*</sup></b> (months) |                                         |                                                           |
| Median                                       | 47.4                                    | 47.0                                                      |
| HR (95% CI)                                  | 0.81 (0.53; 1.23)                       |                                                           |
| p-value                                      | 0.3267                                  |                                                           |

<sup>#</sup> A total of 154 patients (ECOG Performance Status 0 or 1) were randomized. However, two of the randomized patients discontinued the study before receiving any study treatment

<sup>^</sup> Blinded independent review (protocol-defined primary analysis)

<sup>\*</sup> Exploratory analysis: final OS analysis at clinical cut off on 31 October 2017, approx. 59% of patients had died.

CI, confidence interval; HR, Hazard ratio from unstratified Cox regression analysis; NR, not reached.

Advanced and/or metastatic renal cell cancer (mRCC)

*Bevacizumab in combination with interferon alfa-2a for the first-line treatment of advanced and/ or metastatic renal cell cancer (BO17705)*

This was a phase III randomised double-blind trial conducted to evaluate the efficacy and safety of bevacizumab in combination with interferon (IFN) alfa-2a versus IFN alfa-2a alone as first-line treatment in mRCC. The 649 randomised patients (641 treated) had Karnofsky Performance Status (KPS) of  $\geq 70\%$ , no CNS metastases and adequate organ function. Patients were nephrectomised for primary renal cell carcinoma. Bevacizumab 10 mg/kg was given every 2 weeks until disease progression. IFN alfa-2a was given up to 52 weeks or until disease progression at a recommended starting dose of 9 MIU three times a week, allowing a dose reduction to 3 MIU three times a week in 2 steps. Patients were stratified according to country and Motzer score and the treatment arms were shown to be well balanced for the prognostic factors.

The primary endpoint was overall survival, with secondary endpoints for the trial including progression-free survival. The addition of bevacizumab to IFN- $\alpha$ -2a significantly increased PFS and objective tumour response rate. These results have been confirmed through an independent radiological review. However, the increase in the primary endpoint of overall survival by 2 months was not significant (HR=0.91). A high proportion of patients (approximately 63% IFN/placebo; 55% bevacizumab/IFN) received a variety of non-specified post-trial anti-cancer therapies, including antineoplastic agents, which may have impacted the analysis of overall survival.

The efficacy results are presented in Table 15

**Table 15 Efficacy results for trial BO17705**

|                                                                         | BO17705                                  |                                    |
|-------------------------------------------------------------------------|------------------------------------------|------------------------------------|
|                                                                         | Placebo + IFN <sup>a</sup>               | Bv <sup>b</sup> + IFN <sup>a</sup> |
| Number of patients                                                      | 322                                      | 327                                |
| Progression-free survival                                               |                                          |                                    |
| Median (months)                                                         | 5.4                                      | 10.2                               |
| Hazard ratio<br>95% CI                                                  | 0.63<br>0.52, 0.75<br>(p-value < 0.0001) |                                    |
| Objective response rate (%) in Patients with<br>measurable disease<br>N | 289                                      | 306                                |
| Response rate                                                           | 12.8%                                    | 31.4%                              |
|                                                                         | (p-value < 0.0001)                       |                                    |

<sup>a</sup> Interferon alfa-2a 9 MIU 3x/week

<sup>b</sup> Bevacizumab 10 mg/kg q 2 wk

|                        |                                        |      |
|------------------------|----------------------------------------|------|
| Overall survival       |                                        |      |
| Median (months)        | 21.3                                   | 23.3 |
| Hazard ratio<br>95% CI | 0.91<br>0.76, 1.10<br>(p-value 0.3360) |      |

An exploratory multivariate Cox regression model using backward selection indicated that the following baseline prognostic factors were strongly associated with survival independent of treatment: gender, white blood cell count, platelets, body weight loss in the 6 months prior to trial entry, number of metastatic sites, sum of longest diameter of target lesions, Motzer score. Adjustment for these baseline factors resulted in a treatment hazard ratio of 0.78 (95% CI [0.63; 0.96], p=0.0219), indicating a 22% reduction in the risk of death for patients in the bevacizumab + IFN alfa-2a arm compared to IFN alfa-2a arm.

Ninety seven (97) patients in the IFN alfa-2a arm and 131 patients in the bevacizumab arm reduced the dose of IFN alfa-2a from 9 MIU to either 6 or 3 MIU three times a week as pre-specified in the protocol. Dose-reduction of IFN alfa-2a did not appear to affect the efficacy of the combination of bevacizumab and IFN alfa-2a based on PFS event free rates over time, as shown by a sub-group analysis. The 131

patients in the bevacizumab + IFN alfa-2a arm who reduced and maintained the IFN alfa-2a dose at 6 or 3 MIU during the trial, exhibited at 6, 12 and 18 months PFS event free rates of 73, 52 and 21% respectively, as compared to 61, 43 and 17% in the total population of patients receiving bevacizumab + IFN alfa-2a.

#### *AVF2938*

This was a randomised, double-blind, phase II clinical trial investigating bevacizumab 10 mg/kg in a 2 weekly schedule with the same dose of bevacizumab in combination with 150 mg daily erlotinib, in patients with metastatic clear cell RCC. A total of 104 patients were randomised to treatment in this trial, 53 to bevacizumab 10 mg/kg every 2 weeks plus placebo and 51 to bevacizumab 10 mg/kg every 2 weeks plus erlotinib 150 mg daily. The analysis of the primary endpoint showed no difference between the bevacizumab + Placebo arm and the bevacizumab + Erlotinib arm (median PFS 8.5 versus 9.9 months). Seven patients in each arm had an objective response. The addition of erlotinib to bevacizumab did not result in an improvement in OS (HR = 1.764; p=0.1789), duration of objective response (6.7 vs 9.1 months) or time to symptom progression (HR = 1.172; p=0.5076).

#### *AVF0890*

This was a randomised phase II trial conducted to compare the efficacy and safety of bevacizumab versus placebo. A total of 116 patients were randomised to receive bevacizumab 3 mg/kg every 2 weeks (n=39), 10 mg/kg every 2 weeks; (n=37), or placebo (n=40). An interim analysis showed there was a significant prolongation of the time to progression of disease in the 10 mg/kg group as compared with the placebo group (hazard ratio, 2.55; p < 0.001). There was a small difference, of borderline significance, between the time to progression of disease in the 3 mg/kg group and that in the placebo group (hazard ratio, 1.26; p=0.053). Four patients had objective (partial) response, and all of these had received the 10 mg/kg dose bevacizumab; the ORR for the 10 mg/kg dose was 10%.

#### *Epithelial ovarian, fallopian tube and primary peritoneal cancer*

##### *Front-line treatment of ovarian cancer*

The safety and efficacy of bevacizumab in the front-line treatment of patients with epithelial ovarian, fallopian tube or primary peritoneal cancer were studied in two phase III trials (GOG-0218 and BO17707) that evaluated the effect of the addition of bevacizumab to carboplatin and paclitaxel compared to the chemotherapy regimen alone.

#### *GOG-0218*

The GOG-0218 study was a phase III multicentre, randomised, double-blind, placebo-controlled, three arm study evaluating the effect of adding bevacizumab to an approved chemotherapy regimen (carboplatin and paclitaxel) in patients with advanced ( Stages IIIB, IIIC and IV according to FIGO staging version dated 1988) epithelial ovarian, fallopian tube or primary peritoneal cancer.

Patients who had received prior therapy with bevacizumab or prior systemic anticancer therapy for ovarian cancer (e.g. chemotherapy, monoclonal antibody therapy, tyrosine kinase inhibitor therapy, or hormonal therapy) or previous radiotherapy to the abdomen or pelvis were excluded from the study.

A total of 1873 patients were randomised in equal proportions to the following three arms:

CPP arm: Five cycles of placebo (started cycle 2) in combination with carboplatin (AUC 6) and paclitaxel (175 mg/m<sup>2</sup>) for 6 cycles followed by placebo alone, for a total of up to 15 months of therapy

CPB15 arm: Five cycles of bevacizumab (15 mg/kg q3w started cycle 2) in combination with carboplatin (AUC 6) and paclitaxel (175 mg/m<sup>2</sup>) for 6 cycles followed by placebo alone, for a total of up to 15 months of therapy

CPB15+ arm: Five cycles of bevacizumab (15 mg/kg q3w started cycle 2) in combination with carboplatin (AUC 6) and paclitaxel (175 mg/m<sup>2</sup>) for 6 cycles followed by continued use of bevacizumab (15 mg/kg q3w) as single agent for a total of up to 15 months of therapy.

The majority of patients included in the study were White (87% in all three arms); the median age was 60 years in CPP and CPB15 arms and 59 years in CPB15+ arm; and 29% of patients in CPP or CPB15 and 26% in CPB15+ were over 65 years of age. Overall approximately 50% of patients had a GOG PS of 0 at baseline, 43% a GOG PS score of 1, and 7% a GOG PS score of 2. Most patients had EOC (82% in CPP and CPB15, 85% in CPB15+) followed by PPC (16% in CPP, 15% in CPB15, 13% in CPB15+) and FTC (1% in CPP, 3% in CPB15, 2% in CPB15+). The majority of patients had serous adenocarcinoma

histologic type (85% in CPP and CPB15, 86% in CPB15+). Overall approximately 34% of patients were FIGO Stage III optimally debulked with gross residual disease, 40% Stage III sub-optimally debulked, and 26% were Stage IV patients.

The primary endpoint was PFS based on investigator's assessment of disease progression based on radiological scans or CA 125 levels, or symptomatic deterioration per protocol. In addition, a prespecified analysis of the data censoring for CA-125 progression events was conducted, as well as an independent review of PFS as determined by radiological scans.

The trial met its primary objective of PFS improvement. Compared to patients treated with chemotherapy (carboplatin and paclitaxel) alone in the front-line setting, patients who received bevacizumab at a dose of 15 mg/kg q3w in combination with chemotherapy and continued to receive bevacizumab alone (CPB15+), had a clinically meaningful and statistically significant improvement in PFS.

In patients who only received bevacizumab in combination with chemotherapy and did not continue to receive bevacizumab alone (CPB15), no clinically meaningful benefit in PFS was observed.

The results of this study are summarised in Table 16.

**Table 16 Efficacy results from study GOG-0218**

| Progression-free survival <sup>1</sup> |                  |                      |                      |
|----------------------------------------|------------------|----------------------|----------------------|
|                                        | CPP<br>(n = 625) | CPB15<br>(n = 625)   | CPB15+<br>(n = 623)  |
| Median PFS (months)                    | 10.6             | 11.6                 | 14.7                 |
| Hazard Ratio (95% CI) <sup>2</sup>     |                  | 0.89<br>(0.78, 1.02) | 0.70<br>(0.61, 0.81) |
| p-value <sup>3, 4</sup>                |                  | 0.0437               | < 0.0001             |
| Objective response Rate <sup>5</sup>   |                  |                      |                      |
|                                        | CPP<br>(n = 396) | CPB15<br>(n = 393)   | CPB15 +<br>(n = 403) |
| % pts with objective response          | 63.4             | 66.2                 | 66.0                 |
| p-value                                |                  | 0.2341               | 0.2041               |
| Overall survival <sup>6</sup>          |                  |                      |                      |
|                                        | CPP<br>(n = 625) | CPB15<br>(n = 625)   | CPB15 +<br>(n = 623) |
| Median OS (months)                     | 40.6             | 38.8                 | 43.8                 |
| Hazard Ratio (95% CI) <sup>2</sup>     |                  | 1.07 (0.91, 1.25)    | 0.88 (0.75, 1.04)    |
| p-value <sup>3</sup>                   |                  | 0.2197               | 0.0641               |

<sup>1</sup> Investigator assessed GOG protocol-specified PFS analysis (neither censored for CA-125 progressions nor censored for NPT prior to disease progression) with data cut-off date of 25 February, 2010.

<sup>2</sup> Relative to the control arm; stratified hazard ratio.

<sup>3</sup> One-sided log-rank p-value

<sup>4</sup> Subject to a p-value boundary of 0.0116.

<sup>5</sup> Patients with measurable disease at baseline.

<sup>6</sup> Final overall survival analysis performed when 46.9% of the patients had died.

Prespecified PFS analyses were conducted, all with a cut-off date of 29 September 2009. The results of these prespecified analyses are as follows:

The protocol specified analysis of investigator-assessed PFS (without censoring for CA-125 progression or non-protocol therapy [NPT]) shows a stratified hazard ratio of 0.71 (95% CI: 0.61-0.83, 1-sided log-rank p-value < 0.0001) when CPB15+ is compared with CPP, with a median PFS of 10.4 months in the CPP arm and 14.1 months in the CPB15+ arm.

The primary analysis of investigator-assessed PFS (censoring for CA-125 progressions and NPT) shows a stratified hazard ratio of 0.62 (95% CI: 0.52-0.75, 1-sided log-rank p-value < 0.0001) when CPB15+ is



compared with CPP, with a median PFS of 12.0 months in the CPP arm and 18.2 months in the CPB15+ arm.

The analysis of PFS as determined by the independent review committee (censoring for NPT) shows a stratified hazard ratio of 0.62 (95% CI: 0.50-0.77, 1-sided log-rank p-value < 0.0001) when CPB15+ is compared with CPP, with a median PFS of 13.1 in the CPP arm and 19.1 months in the CPB15+ arm.

PFS subgroup analyses by disease stage and debulking status are summarised in Table 17. These results demonstrate robustness of the analysis of PFS as shown in Table 16.

**Table 17 PFS<sup>1</sup> results by disease stage and debulking status from study GOG-0218**

| Randomised patients stage III optimally debulked disease <sup>2,3</sup>       |                  |                      |                      |
|-------------------------------------------------------------------------------|------------------|----------------------|----------------------|
|                                                                               | CPP<br>(n = 219) | CPB15<br>(n = 204)   | CPB15+<br>(n = 216)  |
| Median PFS (months)                                                           | 12.4             | 14.3                 | 17.5                 |
| Hazard ratio (95% CI) <sup>4</sup>                                            |                  | 0.81<br>(0.62, 1.05) | 0.66<br>(0.50, 0.86) |
| Randomised patients with stage III suboptimally debulked disease <sup>3</sup> |                  |                      |                      |
|                                                                               | CPP<br>(n = 253) | CPB15<br>(n = 256)   | CPB15+<br>(n = 242)  |
| Median PFS (months)                                                           | 10.1             | 10.9                 | 13.9                 |
| Hazard ratio (95% CI) <sup>4</sup>                                            |                  | 0.93<br>(0.77, 1.14) | 0.78<br>(0.63, 0.96) |
| Randomised patients with stage IV disease                                     |                  |                      |                      |
|                                                                               | CPP<br>(n = 153) | CPB15<br>(n = 165)   | CPB15+<br>(n = 165)  |
| Median PFS (months)                                                           | 9.5              | 10.4                 | 12.8                 |
| Hazard Ratio (95% CI) <sup>4</sup>                                            |                  | 0.90<br>(0.70, 1.16) | 0.64<br>(0.49, 0.82) |

<sup>1</sup> Investigator assessed GOG protocol-specified PFS analysis (neither censored for CA-125 progressions nor censored for NPT prior to disease progression) with data cut-off date of 25 February, 2010

<sup>2</sup> With gross residual disease.

<sup>3</sup> 3.7% of the overall randomised patient population had Stage IIIB disease.

<sup>4</sup> Relative to the control arm.

#### *BO17707 (ICON7)*

BO17707 was a Phase III, two arm, multicentre, randomised, controlled, open-label study comparing the effect of adding bevacizumab to carboplatin plus paclitaxel in patients with FIGO stage I or IIA (Grade 3 or clear cell histology only; n = 142), or FIGO stage IIB - IV (all Grades and all histological types, n = 1386) epithelial ovarian, fallopian tube or primary peritoneal cancer following surgery (NCI-CTCAE v.3). FIGO staging version dated 1988 was used in this trial.

Patients who had received prior therapy with bevacizumab or prior systemic anticancer therapy for ovarian cancer (e.g. chemotherapy, monoclonal antibody therapy, tyrosine kinase inhibitor therapy, or hormonal therapy) or previous radiotherapy to the abdomen or pelvis were excluded from the study.

A total of 1528 patients were randomised in equal proportions to the following two arms:

CP arm: Carboplatin (AUC 6) and paclitaxel (175 mg/m<sup>2</sup>) for 6 cycles of 3 weeks duration

CPB7.5+ arm: Carboplatin (AUC 6) and paclitaxel (175 mg/m<sup>2</sup>) for 6 cycles of 3 weeks plus bevacizumab (7.5 mg/kg q3w) for up to 12 months (bevacizumab was started at cycle 2 of chemotherapy if treatment was initiated within 4 weeks of surgery or at cycle 1 if treatment was initiated more than 4 weeks after surgery).

The majority of patients included in the study were White (96%), the median age was 57 years in both treatment arms, 25% of patients in each treatment arm were 65 years of age or over, and approximately 50% of patients had an ECOG PS of 1; 7% of patients in each treatment arm had an ECOG PS of 2. The

majority of patients had EOC (87.7%) followed by PPC (6.9%) and FTC (3.7%) or a mixture of the three origins (1.7%). Most patients were FIGO Stage III (both 68%) followed by FIGO Stage IV (13% and 14%), FIGO Stage II (10% and 11%) and FIGO Stage I (9% and 7%). The majority of the patients in each treatment arm (74% and 71%) had poorly differentiated (Grade 3) primary tumours at baseline. The incidence of each histologic sub-type of EOC was similar between the treatment arms; 69% of patients in each treatment arm had serous adenocarcinoma histologic type.

The primary endpoint was PFS as assessed by the investigator using RECIST.

The trial met its primary objective of PFS improvement. Compared to patients treated with chemotherapy (carboplatin and paclitaxel) alone in the front-line setting, patients who received bevacizumab at a dose of 7.5 mg/kg q3w in combination with chemotherapy and continued to receive bevacizumab for up to 18 cycles had a statistically significant improvement in PFS.

The results of this study are summarised in Table 18.

**Table 18 Efficacy results from study BO17707 (ICON7)**

| Progression-free survival            |                                            |                      |
|--------------------------------------|--------------------------------------------|----------------------|
|                                      | CP<br>(n = 764)                            | CPB7.5+<br>(n =764)  |
| Median PFS (months) <sup>2</sup>     | 16.9                                       | 19.3                 |
| Hazard ratio [95% CI] <sup>2</sup>   | 0.86 [0.75; 0.98]<br>(p-value = 0.0185)    |                      |
| Objective Response Rate <sup>1</sup> |                                            |                      |
|                                      | CP<br>(n = 277)                            | CPB7.5+<br>(n = 272) |
| Response rate                        | 54.9%                                      | 64.7%                |
|                                      | (p-value = 0.0188)                         |                      |
| Overall Survival <sup>3</sup>        |                                            |                      |
|                                      | CP<br>(n = 764)                            | CPB7.5+<br>(n = 764) |
| Median (months)                      | 58.0                                       | 57.4                 |
| Hazard ratio [95% CI]                | 0.99 [0. 85; 1. 15]<br>(p-value = 0. 8910) |                      |

<sup>1</sup> In patients with measurable disease at baseline.

<sup>2</sup> Investigator assessed PFS analysis with data cut-off date of 30 November 2010.

<sup>3</sup> Final overall survival analysis performed when 46.7% of the patients had died with data cut-off date of 31 March 2013.

The primary analysis of investigator-assessed PFS with a data cut-off date of 28 February 2010 shows an unstratified hazard ratio of 0.79 (95% CI: 0.68-0.91, 2-sided log-rank p-value 0.0010) with a median PFS of 16.0 months in the CP arm and 18.3 months in the CPB7.5+ arm.

PFS subgroup analyses by disease stage and debulking status are summarised in Table 19. These results demonstrate robustness of the primary analysis of PFS as shown in Table 18.

**Table 19 PFS<sup>1</sup> results by disease stage and debulking status from study BO17707 (ICON7)**

| Randomised patients stage III optimally debulked disease <sup>2,3</sup>       |                 |                      |
|-------------------------------------------------------------------------------|-----------------|----------------------|
|                                                                               | CP<br>(n = 368) | CPB7.5+<br>(n = 383) |
| Median PFS (months)                                                           | 17.7            | 19.3                 |
| Hazard ratio (95% CI) <sup>4</sup>                                            |                 | 0.89<br>(0.74, 1.07) |
| Randomised patients with stage III suboptimally debulked disease <sup>3</sup> |                 |                      |
|                                                                               | CP<br>(n = 154) | CPB7.5+<br>(n = 140) |

|                                           |                |                      |
|-------------------------------------------|----------------|----------------------|
| Median PFS (months)                       | 10.1           | 16.9                 |
| Hazard ratio (95% CI) <sup>4</sup>        |                | 0.67<br>(0.52, 0.87) |
| Randomised patients with stage IV disease |                |                      |
|                                           | CP<br>(n = 97) | CPB7.5+<br>(n = 104) |
| Median PFS (months)                       | 10.1           | 13.5                 |
| Hazard Ratio (95% CI) <sup>4</sup>        |                | 0.74<br>(0.55, 1.01) |

<sup>1</sup> Investigator assessed PFS analysis with data cut-off date of 30 November 2010.

<sup>2</sup> With or without gross residual disease.

<sup>3</sup> 5.8% of the overall randomised patient population had Stage IIIB disease.

<sup>4</sup> Relative to the control arm.

#### *Recurrent ovarian cancer*

The safety and efficacy of bevacizumab in the treatment of recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer was studied in three phase III trials (AVF4095g, MO22224 and GOG-0213) with different patient populations and chemotherapy regimens.

- AVF4095g evaluated the efficacy and safety of bevacizumab in combination with carboplatin and gemcitabine, followed by bevacizumab as a single agent in patients with platinum-sensitive recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer.
- GOG-0213 evaluated the efficacy and safety of bevacizumab in combination with carboplatin and paclitaxel, followed by bevacizumab as a single agent in patients with platinum-sensitive recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer.
- MO22224 evaluated the efficacy and safety of bevacizumab in combination with paclitaxel, topotecan, or pegylated liposomal doxorubicin in patients with platinum-resistant recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer.

#### *AVF4095g*

The safety and efficacy of bevacizumab in the treatment of patients with platinum-sensitive, recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, who have not received prior chemotherapy in the recurrent setting or prior bevacizumab treatment, was studied in a phase III randomised, double-blind, placebo-controlled trial (AVF4095g). The study compared the effect of adding bevacizumab to carboplatin and gemcitabine chemotherapy and continuing bevacizumab as a single agent to progression, to carboplatin and gemcitabine alone.

Only patients with histologically documented ovarian, primary peritoneal, or fallopian tube carcinoma that had recurred > 6 months after platinum-based chemotherapy and who had not received chemotherapy in the recurrent setting and who have not received prior therapy with bevacizumab or other VEGF inhibitors or VEGF receptor-targeted agents were included in the study.

A total of 484 patients with measurable disease were randomised 1:1 to either:

- Carboplatin (AUC4, Day 1) and gemcitabine (1000 mg/m<sup>2</sup> on Days 1 and 8) and concurrent placebo every 3 weeks for 6 and up to 10 cycles followed by placebo (every 3 weeks) alone until disease progression or unacceptable toxicity
- Carboplatin (AUC4, Day 1) and gemcitabine (1000 mg/m<sup>2</sup> on Days 1 and 8) and concurrent bevacizumab (15 mg/kg Day 1) every 3 weeks for 6 and up to 10 cycles followed by bevacizumab (15 mg/kg every 3 weeks) alone until disease progression or unacceptable toxicity

The primary endpoint was progression-free survival based on investigator assessment using modified RECIST 1.0. Additional endpoints included objective response, duration of response, overall survival and safety. An independent review of the primary endpoint was also conducted.

The results of this study are summarised in Table 20.

**Table 20 Efficacy results from study AVF4095g**

| Progression-free survival        |                           |                                |                                |                                |
|----------------------------------|---------------------------|--------------------------------|--------------------------------|--------------------------------|
|                                  | Investigator Assessment   |                                | IRC Assessment                 |                                |
|                                  | Placebo+ C/G<br>(n=242)   | bevacizumab + C/G<br>(n=242)   | Placebo+ C/G<br>(n=242)        | bevacizumab + C/G<br>(n=242)   |
| <i>Not censored for NPT</i>      |                           |                                |                                |                                |
| Median PFS (months)              | 8.4                       | 12.4                           | 8.6                            | 12.3                           |
| Hazard ratio<br>(95% CI)         | 0.524 [0.425, 0.645]      |                                | 0.480 [0.377, 0.613]           |                                |
| p –value                         | <0.0001                   |                                | <0.0001                        |                                |
| <i>Censored for NPT</i>          |                           |                                |                                |                                |
| Median PFS (months)              | 8.4                       | 12.4                           | 8.6                            | 12.3                           |
| Hazard ratio<br>(95% CI)         | 0.484 [0.388, 0.605]      |                                | 0.451 [0.351, 0.580]           |                                |
| p –value                         | < 0.0001                  |                                | <0.0001                        |                                |
| Objective response rate          |                           |                                |                                |                                |
|                                  | Investigator Assessment   |                                | IRC Assessment                 |                                |
|                                  | Placebo+ C/G<br>(n = 242) | bevacizumab + C/G<br>(n = 242) | Placebo+ C/G<br>(n = 242)      | bevacizumab + C/G<br>(n = 242) |
| % pts with objective<br>response | 57.4%                     | 78.5%                          | 53.7%                          | 74.8%                          |
| p –value                         | < 0.0001                  |                                | < 0.0001                       |                                |
| Overall survival                 |                           |                                |                                |                                |
|                                  | Placebo+ C/G<br>(n = 242) |                                | bevacizumab + C/G<br>(n = 242) |                                |
| Median OS (months)               | 32.9                      |                                | 33.6                           |                                |
| Hazard Ratio<br>(95% CI)         | 0.952 [0.771, 1.176]      |                                |                                |                                |
| p-value                          | 0.6479                    |                                |                                |                                |

PFS subgroup analyses depending on recurrence since last platinum therapy are summarised in Table 21.

**Table 21 Progression-free survival by time from last platinum therapy to recurrence**

|                                               | Investigator Assessment    |                                |
|-----------------------------------------------|----------------------------|--------------------------------|
| Time from last platinum therapy to recurrence | Placebo + C/G<br>(n = 242) | bevacizumab + C/G<br>(n = 242) |
| <b>6 - 12 months (n=202)</b>                  |                            |                                |
| <b>Median</b>                                 | 8.0                        | 11.9                           |
| Hazard ratio (95% CI)                         | 0.41 (0.29 - 0.58)         |                                |
| <b>&gt; 12 months (n=282)</b>                 |                            |                                |
| <b>Median</b>                                 | 9.7                        | 12.4                           |
| Hazard ratio (95% CI)                         | 0.55 (0.41 – 0.73)         |                                |

### GOG-0213

GOG-0213, a phase III randomized controlled open label trial, studied the safety and efficacy of bevacizumab in the treatment of patients with platinum-sensitive, recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, who have not received prior chemotherapy in the recurrent setting. There was no exclusion criterion for prior anti-angiogenic therapy. The study evaluated the effect of adding bevacizumab to carboplatin+paclitaxel and continuing bevacizumab as a single agent until disease progression or unacceptable toxicity compared to carboplatin+paclitaxel alone.

A total of 673 patients were randomized in equal proportions to the following two treatment arms:

- CP arm: Carboplatin (AUC5) and paclitaxel (175 mg/m<sup>2</sup> IV) every 3 weeks for 6 and up to 8 cycles.
- CPB arm: Carboplatin (AUC5) and paclitaxel (175 mg/m<sup>2</sup> IV) and concurrent bevacizumab (15 mg/kg) every 3 weeks for 6 and up to 8 cycles, followed by bevacizumab (15 mg/kg every 3 weeks) alone until disease progression or unacceptable toxicity.

Most patients in both the CP arm (80.4%) and the CPB arm (78.9%) were White. The median age was 60.0 years in the CP arm and 59.0 years in the CPB arm. The majority of patients (CP: 64.6%; CPB: 68.8%) were in the age category < 65 years. At baseline, most patients in both treatment arms had a GOG PS of 0 (CP: 82.4%; CPB: 80.7%) or 1 (CP: 16.7%; CPB: 18.1%). A GOG PS of 2 at baseline was reported in 0.9% of patients in the CP arm and in 1.2% of patients in the CPB arm.

The primary efficacy endpoint was overall survival (OS). The main secondary efficacy endpoint was progression-free survival (PFS). Results are presented in Table 22.

**Table 22 Efficacy results<sup>1,2</sup> from study GOG-0213**

| Primary Endpoint                                       |                          |                |
|--------------------------------------------------------|--------------------------|----------------|
| Overall Survival (OS)                                  | CP<br>(n=336)            | CPB<br>(n=337) |
| Median OS (months)                                     | 37.3                     | 42.6           |
| Hazard ratio (95% CI) (eCRF) <sup>a</sup>              | 0.823 [CI: 0.680, 0.996] |                |
| p-Value                                                | 0.0447                   |                |
| Hazard ratio (95% CI) (registration form) <sup>b</sup> | 0.838 [CI: 0.693, 1.014] |                |
| p-Value                                                | 0.0683                   |                |
| Secondary Endpoint                                     |                          |                |
| Progression-free survival (PFS)                        | CP<br>(n=336)            | CPB<br>(n=337) |
| Median PFS (months)                                    | 10.2                     | 13.8           |
| Hazard ratio (95% CI)                                  | 0.613 [CI: 0.521, 0.721] |                |
| p-value                                                | <0.0001                  |                |

<sup>1</sup> Final Analysis <sup>2</sup> Tumour assessments and response evaluations were determined by the investigators using the GOG RECIST criteria (Revised RECIST guideline (version 1.1). Eur J Cancer. 2009;45:228Y247).

<sup>a</sup> Hazard ratio was estimated from Cox proportional hazards models stratified by the duration of platinum free-interval prior to enrolling onto this study per eCRF (electronic case report form) and secondary surgical debulking status Yes/No (Yes=randomized to undergo cytoreduction or randomized to not undergo cytoreduction; No= not a candidate or did not consent to cytoreduction). <sup>b</sup> stratified by the duration of treatment free-interval prior to enrolling onto this study per the registration form, and secondary surgical debulking status Yes/No.

The trial met its primary objective of OS improvement. Treatment with bevacizumab at 15 mg/kg every 3 weeks in combination with chemotherapy (carboplatin and paclitaxel) for 6 and up to 8 cycles, followed by bevacizumab until disease progression or unacceptable toxicity resulted, when data were derived from eCRF, in a clinically meaningful and statistically significant improvement in OS compared to treatment with carboplatin and paclitaxel alone.

#### MO22224

Study MO22224 evaluated the efficacy and safety of bevacizumab in combination with chemotherapy for platinum-resistant recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer. This study was designed as an open-label, randomized, two-arm Phase III evaluation of bevacizumab plus chemotherapy (CT+BV) versus chemotherapy alone (CT).

A total of 361 patients were enrolled into this study and administered either chemotherapy (paclitaxel, topotecan, or pegylated liposomal doxorubicin (PLD) alone or in combination with bevacizumab:

- CT Arm (chemotherapy alone):
  - Paclitaxel 80 mg/m<sup>2</sup> as a 1-hour IV infusion on Days 1, 8, 15, and 22 every 4 weeks.
  - Topotecan 4 mg/m<sup>2</sup> as a 30-minute IV infusion on Days 1, 8, and 15 every 4 weeks. Alternatively, a 1.25 mg/m<sup>2</sup> dose could be administered over 30 minutes on Days 1–5 every 3 weeks.
  - PLD 40 mg/m<sup>2</sup> as a 1 mg/min IV infusion on Day 1 only every 4 weeks. After Cycle 1, the drug could be delivered as a 1-hour infusion.
- CT+BV Arm (chemotherapy plus bevacizumab):
  - The chosen chemotherapy was combined with bevacizumab 10 mg/kg IV every 2 weeks (or bevacizumab 15 mg/kg every 3 weeks if used in combination with topotecan 1.25 mg/m<sup>2</sup> on Days 1–5 every 3 weeks)

Eligible patients had epithelial ovarian, fallopian tube or primary peritoneal cancer that progressed within <6 months of previous platinum therapy consisting of a minimum of 4 platinum therapy cycles. Patients should have had a life expectancy of ≥ 12 weeks and no prior radiotherapy to the pelvis or abdomen. Most patients were FIGO Stage IIIC or Stage IV. The majority of patients in both arms had an ECOG Performance Status (PS) of 0 (CT: 56.4% vs. CT + BV: 61.2%). The percentage of patients with an ECOG PS of 1 or ≥ 2 was 38.7% and 5.0% in the CT arm, and 29.8% and 9.0% in the CT + BV arm. Information on race exists for 29.3% of patients and nearly all patients were white. The median age of patients was 61.0 (range: 25-84) years. A total of 16 patients (4.4%) were > 75 years old. The overall rates of discontinuation due to adverse events were 8.8% in the CT arm and 43.6% in the CT + BV arm (mostly due to Grade 2-3 adverse events) and the median time to discontinuation in the CT + BV arm was 5.2 months compared with 2.4 months in the CT arm. The rates of discontinuation due to adverse events in the subgroup of patients > 65 years old were 8.8% in the CT arm and 50.0% in the CT + BV arm. The HR for PFS was 0.47 (95% CI: 0.35, 0.62) and 0.45 (95% CI: 0.31, 0.67) for the < 65 and ≥ 65 subgroups, respectively.

The primary endpoint was progression-free-survival, with secondary endpoints including objective response rate and overall survival. Results are presented in Table 23.

**Table 23 Efficacy Results from Study MO22224**

| <u>Primary Endpoint</u>              |                      |                  |
|--------------------------------------|----------------------|------------------|
| Progression-Free Survival*           |                      |                  |
|                                      | CT<br>(n=182)        | CT+BV<br>(n=179) |
| Median (months)                      | 3.4                  | 6.7              |
| Hazard ratio<br>(95% CI)             | 0.379 [0.296, 0.485] |                  |
| p-value                              | <0.0001              |                  |
| <u>Secondary Endpoints</u>           |                      |                  |
| Objective Response Rate**            |                      |                  |
|                                      | CT<br>(n=144)        | CT+BV<br>(n=142) |
| % patients with objective response   | 18 (12.5%)           | 40 (28.2%)       |
| p –value                             | 0.0007               |                  |
| Overall Survival (final analysis)*** |                      |                  |
|                                      | CT<br>(n=182)        | CT+BV<br>(n=179) |
| Median OS (months)                   | 13.3                 | 16.6             |
| Hazard Ratio<br>(95% CI)             | 0.870 [0.678, 1.116] |                  |

|         |        |
|---------|--------|
| p-value | 0.2711 |
|---------|--------|

All analyses presented in this table are stratified analyses.

\* Primary analysis was performed with a data cut-off date of 14 November 2011.

\*\*Randomized Patients with Measurable Disease at Baseline.

\*\*\*The final analysis of overall survival was performed when 266 deaths, which account for 73.7 % of enrolled patients, were observed.

The trial met its primary objective of PFS improvement. Compared to patients treated with chemotherapy (paclitaxel, topotecan or PLD) alone in the recurrent platinum-resistant setting, patients who received bevacizumab at a dose of 10 mg/kg every 2 weeks (or 15 mg/kg every 3 weeks if used in combination with 1.25 mg/m<sup>2</sup> topotecan on Days 1–5 every 3 weeks) in combination with chemotherapy and continued to receive bevacizumab until disease progression or unacceptable toxicity, had a statistically significant improvement in PFS. The exploratory PFS and OS analyses by chemotherapy cohort (paclitaxel, topotecan and PLD) are summarized in Table 24.

**Table 24: Exploratory PFS and OS analyses by chemotherapy cohort**

|                       | CT                | CT+BV |
|-----------------------|-------------------|-------|
| <b>Paclitaxel</b>     | n=115             |       |
| Median PFS (months)   | 3.9               | 9.2   |
| Hazard ratio (95% CI) | 0.47 [0.31, 0.72] |       |
| Median OS (months)    | 13.2              | 22.4  |
| Hazard ratio (95% CI) | 0.64 [0.41, 0.99] |       |
| <b>Topotecan</b>      | n=120             |       |
| Median PFS (months)   | 2.1               | 6.2   |
| Hazard ratio (95% CI) | 0.28 [0.18, 0.44] |       |
| Median OS (months)    | 13.3              | 13.8  |
| Hazard ratio (95% CI) | 1.07 [0.70, 1.63] |       |
| <b>PLD</b>            | n=126             |       |
| Median PFS (months)   | 3.5               | 5.1   |
| Hazard ratio (95% CI) | 0.53 [0.36, 0.77] |       |
| Median OS (months)    | 14.1              | 13.7  |
| Hazard ratio (95% CI) | 0.91 [0.61, 1.35] |       |

### Cervical Cancer

#### *GOG-0240*

The efficacy and safety of bevacizumab in combination with chemotherapy (paclitaxel and

cisplatin or paclitaxel and topotecan) in the treatment for patients with persistent, recurrent or metastatic carcinoma of the cervix was evaluated in study GOG-0240, a randomised, four-arm, open label, multi-centre phase III trial.

A total of 452 patients were randomised to receive either

- Paclitaxel 135 mg/m<sup>2</sup> IV over 24 hours on Day 1 and cisplatin 50 mg/m<sup>2</sup> IV on Day 2, every 3 weeks (q3w); or  
Paclitaxel 175 mg/m<sup>2</sup> IV over 3 hours on Day 1 and cisplatin 50 mg/m<sup>2</sup> IV on Day 2 (q3w); or  
Paclitaxel 175 mg/m<sup>2</sup> IV over 3 hours on Day 1 and cisplatin 50 mg/m<sup>2</sup> IV on Day 1 (q3w)
- Paclitaxel 135 mg/m<sup>2</sup> IV over 24 hours on Day 1 and cisplatin 50 mg/m<sup>2</sup> IV on Day 2 plus bevacizumab 15 mg/kg IV on Day 2 (q3w); or

Paclitaxel 175 mg/m<sup>2</sup> IV over 3 hours on Day 1 and cisplatin 50 mg/m<sup>2</sup> IV on Day 2 z plus bevacizumab 15 mg/kg IV on Day 2 (q3w); or

Paclitaxel 175 mg/m<sup>2</sup> IV over 3 hours on Day 1 and cisplatin 50 mg/m<sup>2</sup> IV on Day 1 plus bevacizumab 15 mg/kg IV on Day 1 (q3w)

- Paclitaxel 175 mg/m<sup>2</sup> IV over 3 hours on Day 1 and topotecan 0.75 mg/m<sup>2</sup> IV over 30 minutes on days 1-3 (q3w)
- Paclitaxel 175 mg/m<sup>2</sup> IV over 3 hours on Day 1 and topotecan 0.75 mg/m<sup>2</sup> IV over 30 minutes on Days 1-3 plus bevacizumab 15 mg/kg IV on Day 1 (q3w)

Eligible patients had persistent, recurrent or metastatic squamous cell carcinoma, adenosquamous carcinoma, or adenocarcinoma of the cervix which was not amenable to curative treatment with surgery and/or radiation therapy and who have not received prior therapy with bevacizumab or other VEGF inhibitors or VEGF receptor–targeted agents.

The median age was 46.0 years (range: 20-83) in the Chemo alone group and 48.0 (range: 22-85) in the Chemo+bevacizumab group; with 9.3% of patients in the Chemo alone group and 7.5% of patients in the Chemo+bevacizumab group over the age of 65 years.

Of the 452 patients randomized at baseline, the majority of patients were white (80.0% in the Chemo alone group and 75.3% in the Chemo+ bevacizumab group), had squamous cell carcinoma (67.1% in the Chemo alone group and 69.6% in the Chemo+bevacizumab group), had persistent/recurrent disease (83.6% in the Chemo alone group and 82.8% in the Chemo+bevacizumab group), had 1-2 metastatic sites (72.0% in the Chemo alone group and 76.2% in the Chemo+bevacizumab group), had lymph node involvement (50.2% in the Chemo alone group and 56.4% in the Chemo+bevacizumab group), and had a platinum free interval  $\geq$  6 months (72.5% in the Chemo alone group and 64.4% in the Chemo+bevacizumab group).

The primary efficacy endpoint was overall survival. Secondary efficacy endpoints included progression-free survival and objective response rate. Results from the primary analysis and the follow-up analysis are presented by bevacizumab Treatment and by Trial Treatment in Table 25 and Table 26, respectively.

**Table 25 Efficacy results from study GOG-0240 by bevacizumab Treatment**

|                                                                 | Chemotherapy<br>(n=225)                                | Chemotherapy<br>+ bevacizumab<br>(n=227) |
|-----------------------------------------------------------------|--------------------------------------------------------|------------------------------------------|
| <b><u>Primary Endpoint</u></b>                                  |                                                        |                                          |
| <b>Overall Survival – Primary analysis<sup>6</sup></b>          |                                                        |                                          |
| Median (months) <sup>1</sup>                                    | 12.9                                                   | 16.8                                     |
| Hazard ratio [95% CI]                                           | 0.74 [0.58, 0.94]<br>(p-value <sup>5</sup> = 0.0132)   |                                          |
| <b>Overall Survival – Follow-up analysis<sup>7</sup></b>        |                                                        |                                          |
| Median (months) <sup>1</sup>                                    | 13.3                                                   | 16.8                                     |
| Hazard ratio [95% CI]                                           | 0.76 [0.62, 0.94]<br>(p-value <sup>5,8</sup> = 0.0126) |                                          |
| <b><u>Secondary Endpoints</u></b>                               |                                                        |                                          |
| <b>Progression-free survival – Primary analysis<sup>6</sup></b> |                                                        |                                          |
| Median PFS (months) <sup>1</sup>                                | 6.0                                                    | 8.3                                      |
| Hazard ratio [95% CI]                                           | 0.66 [0.54, 0.81]<br>(p-value <sup>5</sup> <0.0001)    |                                          |
| <b>Best Overall Response – Primary analysis<sup>6</sup></b>     |                                                        |                                          |
| Responders (Response rate <sup>2</sup> )                        | 76 (33.8 %)                                            | 103 (45.4 %)                             |
| 95% CI for Response Rates <sup>3</sup>                          | [27.6%, 40.4%]                                         | [38.8%, 52.1%]                           |
| Difference in Response Rates                                    | 11.60%                                                 |                                          |
| 95% CI for Difference in Response Rates <sup>4</sup>            | [2.4%, 20.8%]                                          |                                          |



|                            |        |
|----------------------------|--------|
| p-value (Chi-squared Test) | 0.0117 |
|----------------------------|--------|

<sup>1</sup> Kaplan-Meier estimates

<sup>2</sup> Patients and percentage of patients with best overall response of confirmed CR or PR; percentage calculated on patients with measurable disease at baseline

<sup>3</sup> 95% CI for one sample binomial using Pearson-Clopper method

<sup>4</sup> Approximate 95% CI for difference of two rates using Hauck-Anderson method

<sup>5</sup> log-rank test (stratified)

<sup>6</sup> Primary analysis was performed with a data cut-off date of 12 December 2012 and is considered the final analysis

<sup>7</sup> Follow-up analysis was performed with a data cut-off date of 07 March 2014

<sup>8</sup> p-value displayed for descriptive purpose only

**Table 26 Overall survival results from study GOG-0240 by Trial Treatment**

| Treatment Comparison                            | Other Factor          | Overall survival – Primary analysis <sup>1</sup><br>Hazard Ratio (95% CI) | Overall survival - Follow-up analysis <sup>2</sup><br>Hazard Ratio (95% CI) |
|-------------------------------------------------|-----------------------|---------------------------------------------------------------------------|-----------------------------------------------------------------------------|
| bevacizumab vs. No bevacizumab                  | Cisplatin+ Paclitaxel | 0.72 (0.51, 1.02)<br>(17.5 vs.14.3 months; p = 0.0609)                    | 0.75 (0.55, 1.01)<br>(17.5 vs.15.0 months; p = 0.0584)                      |
|                                                 | Topotecan+ Paclitaxel | 0.76 (0.55, 1.06)<br>(14.9 vs. 11.9 months; p = 0.1061)                   | 0.79 (0.59, 1.07)<br>(16.2 vs. 12.0 months; p = 0.1342)                     |
| Topotecan+ Paclitaxel vs. Cisplatin+ Paclitaxel | bevacizumab           | 1.15 (0.82, 1.61)<br>(14.9 vs. 17.5 months; p = 0.4146)                   | 1.15 (0.85, 1.56)<br>(16.2 vs 17.5 months; p = 0.3769)                      |
|                                                 | No bevacizumab        | 1.13 (0.81, 1.57)<br>(11.9 vs.14.3 months; p = 0.4825)                    | 1.08 (0.80, 1.45)<br>(12.0 vs 15.0 months; p = 0.6267)                      |

<sup>1</sup> Primary analysis was performed with a data cut-off date of 12 December 2012 and is considered the final analysis

<sup>2</sup> Follow-up analysis was performed with a data cut-off date of 07 March 2014; all p-values are displayed for descriptive purpose only

#### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies, in all subsets of the paediatric population, in breast carcinoma, adenocarcinoma of the colon and rectum, lung carcinoma (small cell and non-small cell carcinoma), kidney and renal pelvis carcinoma (excluding nephroblastoma, nephroblastomatosis, clear cell sarcoma, mesoblastic nephroma, renal medullary carcinoma and rhabdoid tumour of the kidney), ovarian carcinoma (excluding rhabdomyosarcoma and germ cell tumours), fallopian tube carcinoma (excluding rhabdomyosarcoma and germ cell tumours), peritoneal carcinoma (excluding blastomas and sarcomas) and cervix and corpus uteri carcinoma.

#### *High-grade glioma*

Anti-tumour activity was not observed in two earlier studies among a total of 30 children aged > 3 years old with relapsed or progressive high-grade glioma when treated with bevacizumab and irinotecan (CPT-11). There is insufficient information to determine the safety and efficacy of bevacizumab in children with newly-diagnosed high-grade glioma.

In a single-arm study (PBTC-022), 18 children with recurrent or progressive non-pontine high-grade glioma (including 8 with glioblastoma [WHO Grade IV], 9 with anaplastic astrocytoma [Grade III] and 1 with anaplastic oligodendroglioma [Grade III]) were treated with bevacizumab (10 mg/kg) two weeks apart and then with bevacizumab in combination with CPT-11 (125-350 mg/m<sup>2</sup>) once every two weeks until progression. There were no objective (partial or complete) radiological responses (MacDonald

criteria). Toxicity and adverse reactions included arterial hypertension and fatigue as well as CNS ischaemia with acute neurological deficit.

In a retrospective single institution series, 12 consecutive (2005 to 2008) children with relapsed or progressive high-grade glioma (3 with WHO Grade IV, 9 with Grade III) were treated with bevacizumab (10 mg/kg) and irinotecan (125 mg/m<sup>2</sup>) every 2 weeks. There were no complete responses and 2 partial responses (MacDonald criteria).

In a randomized phase II study (BO25041) a total of 121 patients aged  $\geq 3$  years to  $<18$  years with newly diagnosed supratentorial or infratentorial cerebellar or peduncular high-grade glioma (HGG) were treated with post operative radiation therapy (RT) and adjuvant temozolomide (T) with and without bevacizumab: 10 mg/kg every 2 weeks IV.

The study did not meet its primary endpoint of demonstrating a significant improvement of EFS (Central Radiology Review Committee (CRRC)-assessed) when bevacizumab was added to the RT/T arm compared with RT/T alone (HR = 1.44; 95% CI: 0.90, 2.30). These results were consistent with those from various sensitivity analyses and in clinically relevant subgroups. The results for all secondary endpoints (investigator assessed EFS, and ORR and OS) were consistent in showing no improvement associated with the addition of bevacizumab to the RT/T arm compared with the RT/T arm alone.

Addition of bevacizumab to RT/T did not demonstrate clinical benefit in study BO25041 in 60 evaluable children patients with newly diagnosed supratentorial or infratentorial cerebellar or peduncular high-grade glioma (HGG) (See section 4.2 for information on paediatric use).

#### *Soft tissue sarcoma*

In a randomized phase II study (BO20924) a total of 154 patients aged  $\geq 6$  months to  $<18$  years with newly diagnosed metastatic rhabdomyosarcoma and non-rhabdomyosarcoma soft tissue sarcoma were treated with standard of care (Induction IVADO/IVA+/- local therapy followed by Maintenance Vinorelbine and cyclophosphamide) with or without bevacizumab (2.5 mg/kg/week) for a total duration of treatment of approximately 18 months. At the time of the final primary analysis, the primary endpoint of EFS by independent central review did not show a statistically significant difference between the two treatment arms, with HR of 0.93 (95% CI: 0.61, 1.41; p-value = 0.72). The difference in ORR per independent central review was 18% (CI: 0.6%, 35.3%) between the two treatment arms in the few patients who had evaluable tumor at baseline and had a confirmed response prior to receiving any local therapy : 27/75 patients (36.0%, 95% CI: 25.2%, 47.9%) in the Chemo arm and 34/63 patients (54.0%, 95% CI: 40.9%, 66.6%) in the Bv + Chemo arm. The final Overall Survival (OS) analyses showed no significant clinical benefit from addition of bevacizumab to chemotherapy in this patient population.

Addition of bevacizumab to standard of care did not demonstrate clinical benefit in clinical trial BO20924, in 71 evaluable children (from age 6 months to less than 18 years old) patients with metastatic Rhabdomyosarcoma and non-Rhabdomyosarcoma Soft Tissue Sarcoma.

(See section 4.2 for information on paediatric use).

The incidence of AEs, including Grade  $\geq 3$  AEs and SAEs, was similar between the two treatment arms. No AEs leading to death occurred in either treatment arm; all deaths were attributed to disease progression. Bevacizumab addition to multimodal standard of care treatment seemed to be tolerated in this paediatric population.

### **Clinical studies with Dr Reddy's bevacizumab biosimilar**

**BZ-01-001 study** was a single Dose, double-Blind, parallel arm, comparative pharmacokinetic study (Phase I) of three bevacizumab preparations, i.e. DRL\_BZ, reference medicinal product -RMP (Avastin®, EU approved) and reference product - RP (Avastin®, US licensed) which were administered by the intravenous route to normal healthy male volunteers. The study was conducted at 2 study centers in New Zealand. In this double-blind, parallel-group, Phase 1 study (BZ-01-001), men aged 20–45 years were randomized 1:1:1 to receive a single intravenous infusion of 1 mg kg<sup>-1</sup> of bevacizumab as DRL\_BZ, RMP or RP. A total of 149 subjects were randomized (DRL\_BZ, 50; RMP, 50; RP, 49). Primary endpoints included maximum observed serum concentration (C<sub>max</sub>), area under the concentration–time curve from time zero (pre-dose) extrapolated to infinity (AUC(0– $\infty$ )), and area under the concentration–time curve from time zero (pre-dose) to last

quantifiable concentration (AUC(0–t)). Secondary objectives were to compare the safety and immunogenicity of DRL\_BZ with those of the reference products.

For the primary parameters AUC(0–∞), AUC(0–t), and C<sub>max</sub>, a 3-way statistical comparison was conducted to test between DRL\_BZ the RP and the RMP, as well as between the RP and RMP. Based on the predefined acceptance criteria, PK similarity was concluded for all 3 comparisons DRL\_BZ versus RP [US-licensed Avastin] and RMP [EU-approved Avastin] for all primary PK parameters (AUC(0–∞), AUC(0–t), and C<sub>max</sub>) since the 90% CIs for all comparisons were contained within the predefined bioequivalence limits of 80.00% to 125.00% bioequivalence limits acceptance range prospectively defined in the Study Protocol

The observed GMRs (90% CI) were: for the DRL\_BZ vs. RMP comparison 98.40% (92.48–104.70) for C<sub>max</sub>, 94.69% (89.56–100.10) for AUC(0–t) and 94.44% (89.24–99.94) for AUC(0–∞); for the DRL\_BZ vs. RP comparison 99.57% (94.25–105.20) for C<sub>max</sub>, 97.68% (92.42–103.23) for AUC(0–t) and 97.76% (92.47–103.35) for AUC(0–∞). For the reference product comparisons (RMP/RP), the observed GMRs (90% CI) were 101.19% (95.19–107.57) for C<sub>max</sub>, 103.16% (97.92–108.68%) for AUC(0–t) and 103.52% (98.13–109.20%) for AUC(0–∞). Hence, the 90% CIs for the GMRs of the PK primary endpoints (AUC(0–∞), AUC(0–t), and C<sub>max</sub>) were within the prespecified acceptance margins (80–125%) for PK equivalence for all three comparisons: DRL\_BZ vs. RMP, DRL\_BZ vs. RP and RMP vs. RP

**Table 27: Statistical comparison of key pharmacokinetic parameters (pharmacokinetic population)**

| Parameter<br>(Units)              | Treatment | n  | Geometric<br>Least-<br>squares<br>Means | 95%<br>Confidence<br>Interval | Pairwise Comparison |              |                               |
|-----------------------------------|-----------|----|-----------------------------------------|-------------------------------|---------------------|--------------|-------------------------------|
|                                   |           |    |                                         |                               | Pair                | Ratio<br>(%) | 90%<br>Confidence<br>Interval |
| C <sub>Max</sub>                  | DRL_BZ    | 50 | 20.99                                   | 20.01, 22.02                  | DRL_BZ/RMP          | 98.40        | 92.48, 104.70                 |
|                                   | RP        | 47 | 21.08                                   | 20.13, 22.08                  | DRL_BZ/RP           | 99.57        | 94.25, 105.20                 |
|                                   | RMP       | 50 | 21.33                                   | 20.14, 22.60                  | RMP/RP              | 101.19       | 95.19, 107.57                 |
| AUC <sub>(0-t)</sub><br>(h*µg/ml) | DRL_BZ    | 50 | 6186                                    | 5883,6504                     | DRL_BZ/RMP          | 94.69        | 89.56,100.10                  |
|                                   | RP        | 43 | 6333                                    | 6058, 6621                    | DRL_BZ/RP           | 97.68        | 92.42,103.23                  |
|                                   | RMP       | 48 | 6533                                    | 6246,6834                     | RMP/RP              | 103.16       | 97.92,108.68                  |
| AUC <sub>(0-∞)</sub><br>(h*µg/ml) | DRL_BZ    | 50 | 6328                                    | 6019,6654                     | DRL_BZ/RMP          | 94.44        | 89.24, 99.94                  |
|                                   | RP        | 43 | 6473                                    | 6189, 6771                    | DRL_BZ/RP           | 97.76        | 92.47, 103.35                 |
|                                   | RMP       | 48 | 6701                                    | 6395, 7021                    | RMP/RP              | 103.52       | 98.13, 109.20                 |

Results based on analysis of variance model with treatment as a fixed effect on the log-transformed values of C<sub>max</sub>, AUC(0–t) and AUC(0–∞). AUC(0–∞), area under the concentration–time curve from time zero (pre-dose) extrapolated to infinity; AUC(0–t), area under the concentration–time curve from time zero (pre-dose) to last quantifiable concentration; C<sub>max</sub>, maximum observed serum concentration over the entire blood sampling interval; CI, confidence interval; DRL\_BZ, Dr. Reddy's Laboratories bevacizumab; RMP, reference medicinal product (EU-approved Avastin®); RP, reference product (US-licensed Avastin®)

The most frequently reported TEAE by preferred term (PT) was headache (10 TEAEs for 9 subjects [18.0%] on DRL\_BZ, 14 TEAEs for 12 subjects [24.5%] on RP, and 24 TEAEs for 20 subjects [40.0%] on RMP). Overall, the majority of subjects had TEAEs that were considered Grade 1 in severity (81subjects [54.4%] with 248 TEAEs). On DRL\_BZ, 26 subjects (52.0%) had 76 TEAEs Grade 1 in severity and 11 subjects (22.0%) had 13 TEAEs considered Grade 2, on RP, 28 subjects (57.1%) had 79 TEAEs considered Grade 1 and 10 subjects (20.4%) had 10 TEAEs considered Grade 2 in severity, and on RMP, 27 subjects (54.0%) had 93 TEAEs considered Grade 1 and 7 subjects (14.0%) had 11 TEAEs considered Grade 2 in severity. One Grade 3 severity TEAE of abdominal pain (considered not to be related to the study drug) was reported in the RMP arm. No Grade 3 severity TEAEs were reported on either DRL\_BZ or RP. Of the 37 subjects (74.0%) with 89 TEAEs on DRL\_BZ, 5 subjects (10.0%) had 7 TEAEs considered related to the study drug; of the 38 subjects (77.6%) with 89 TEAEs on RP, 9 subjects (18.4%) had 13 TEAEs considered related to the study drug; and of the 35 subjects (70.0%) with 105

TEAEs on RMP, 11 subjects (22.0%) had 20 TEAEs considered related to the study drug. Only one subject was confirmed positive for ADA at the Day 85 visit and was thus followed up for up to 12 months. In this subject, ADA results were positive at both 6- and 9-months visits, and the 12-months sample was negative. The subject's nAb results were negative at the Day 85 visit, positive at the 6 months visit (ADA follow-up 1), and negative on subsequent visits.

**BZ-01-002 study:** This is a randomized, multicenter, double-blind study conducted as parts A and B (Phase III). The part A study was planned in patients with mCRC and part B in patients with NSCLC. The study was approved by an independent ethics committee or an institutional review board at different study centers.

The objective of this study was to compare the efficacy, safety, pharmacokinetics, and immunogenicity of a proposed bevacizumab biosimilar (DRL\_BZ) with the innovator Avastin (reference medicinal product [RMP]) in patients with nonresectable metastatic colorectal cancer (mCRC) over a period of 9 months and advanced nonsquamous non-small cell lung cancer (NSCLC) over 6 months. In part A, a total of 117 mCRC patients were intended to receive 5 mg/kg of bevacizumab every 2 weeks along with mFOLFOX6 chemotherapy for a maximum of 18 cycles. In part B, 60 NSCLC patients were to receive 15 mg/kg of bevacizumab every 3 weeks along with pemetrexed and carboplatin for the initial four cycles, followed by pemetrexed for another four cycles.

The primary endpoint was the progression-free survival rate at 6 months (PFS6) in both subparts. The anticipated sample size was 106 evaluable mCRC patients to achieve 85% statistical power for concluding non-inferiority with a margin of half the difference (18.8%) between DRL\_BZ and Avastin, along with a pilot study involving 60 evaluable NSCLC patients. Safety comparison included assessing adverse events (AEs), infusion reactions, and lab abnormalities. Immunogenicity comparison involved the incidence of antidrug antibodies (ADAs) and neutralizing antibodies (NAbs). Pharmacokinetic comparison was planned after the first and fourth dosing cycles of treatment in 24 NSCLC patients.

The PFS6 for mCRC patients treated with DRL\_BZ and RMP was 57.8% and 50% respectively, with a difference in efficacy of 7.8 (–8.7, 23.7), which met the non-inferiority margin defined in the protocol [two-sided 90% CI for the difference in proportion of PFS6 responders in DRL\_BZ minus proportion of PFS6 responders in the Avastin® has lower limit  $\geq -18.8\%$ ], proving the non-inferiority of DRL\_BZ efficacy as compared with Avastin®. The PFS9 was 31.1% and 22.9%, with a difference of 8.2% (–6.9%, 22.9%). The objective response rate (ORR) for DRL\_BZ and RMP was 28.8% and 22.4%, while the disease control rate (DCR) was 44.2% and 37.9% respectively. Overall survival at the EOS was 92.3% (two sided 95% CIs 81.8, 97.0) in the DRL\_BZ arm and 93.1% (two sided 95% CIs 83.6, 97.3) in the Avastin® arm. The difference in percentage between the arms was -0.8% with a two sided 95% CI of -12.1 to 9.8. For NSCLC patients the PFS6 was 44.0% (two sided 90% CI 29.1, 60.1) in the DRL\_BZ arm and 45.0% (two sided 90% CI 28.4, 62.8) in the Avastin® arm, for the mITT population. The ORR was 41.4% and 48.1%, and the DCR was 62.1% and 63%, respectively.

The mean ( $\pm$ SD)  $C_{max}$  for Cycle 1 in the DRL\_BZ arm was 365.45 ( $\pm$ 92.11)  $\mu$ g/mL and in the Avastin® arm was 336.60 ( $\pm$ 118.79)  $\mu$ g/mL. The mean ( $\pm$ SD)  $C_{max}$  for Cycle 4 in the DRL\_BZ arm was 464.40 ( $\pm$ 57.03)  $\mu$ g/mL and in the Avastin® arm was 406.00 ( $\pm$ 68.50)  $\mu$ g/mL. The mean ( $\pm$ SD)  $AUC_{0-t}$  for Cycle 1 in the DRL\_BZ arm was 55518.47 ( $\pm$ 18363.73)  $\mu$ g\*h/mL and in the Avastin® arm was 41374.00 ( $\pm$ 18848.72)  $\mu$ g\*h/mL. The mean ( $\pm$ SD)  $AUC_{0-t}$  for Cycle 4 in the DRL\_BZ arm was 88560.13 ( $\pm$ 10127.80)  $\mu$ g\*h/mL and in the Avastin® arm was 74293.18 ( $\pm$ 39516.68)  $\mu$ g\*h/mL. The mean ( $\pm$ SD)  $AUC_{0-\tau}$  for Cycle 1 in the DRL\_BZ arm was 55518.47 ( $\pm$ 18363.73)  $\mu$ g\*h/mL and in the Avastin® arm was 41374.00 ( $\pm$ 18848.72)  $\mu$ g\*h/mL. The mean ( $\pm$ SD)  $AUC_{0-\tau}$  for Cycle 4 in the DRL\_BZ arm was 88560.13 ( $\pm$ 10127.80)  $\mu$ g\*h/mL and in the Avastin® arm was 74293.18 ( $\pm$ 39516.68)  $\mu$ g\*h/mL.

In mCRC patients none of the patients tested positive for antidrug antibodies (ADAs) at baseline and post bevacizumab administration at weeks 4, 8, 17 and 37) in the DRL\_BZ arm. In the Avastin® arm, overall four patients tested positive for ADAs, of these two patients tested negative and two patients tested positive for neutralising antibodies (nAbs). These two patients turned negative for ADAs in subsequent cycle evaluation i.e. week 17). In NSCLC patients none of the patients tested positive for anti-drug antibodies at baseline and post bevacizumab administration.

The safety analysis population in mCRC group consisted of 117 patients (58 DRL\_BZ arm and 59 Avastin® arm) who received at least one dose of DRL\_BZ or Avastin®. The total dose, treatment duration,

and duration of observation were comparable for both arms. A total of 1246 (572 in the DRL\_BZ arm and 674 in the Avastin® arm) TEAEs were reported in 113 (96.6%) patients [57 (98.3) patients in the DRL\_BZ arm 56 (94.9) patients in the Avastin® arm]. The incidence of TEAEs ((98.3% in the DRL\_BZ arm and 94.9% in the Avastin® arm) was comparable in both arms. The majority of the reported TEAEs were mild [DRL\_BZ arm: 315 events in 52 (89.7%) patients, Avastin® arm: 357 events in 49 (83.1%) patients], followed by moderate TEAEs [DRL\_BZ arm: 150 events in 44 (75.9%) patients, Avastin® arm: 216 events in 47 (79.7%) patients], severe TEAEs [DRL\_BZ arm: 86 events in 36 (62.1%) patients, Avastin® arm: 78 events in 30 (50.8%) patients], life threatening TEAEs [DRL\_BZ arm: 17 events in 10 (17.2%) patients, Avastin® arm: 19 events in 11 (18.6%) patients] and death TEAEs (4 events in each arm).

The most frequently reported TEAEs (>10%) in both arms were, thrombocytopenia [DRL\_BZ arm: 51 events in 19 (32.8%) patients, Avastin® arm: 53 events in 19 (32.2%) patients, anaemia [DRL\_BZ arm: 41 events in 19 (32.8%) patients, Avastin® arm: 31 events in 21 (35.6%) patients and diarrhoea [DRL\_BZ arm: 34 events in 21 (36.2%) patients, Avastin® arm: 39 events in 21 (35.6%) patients] ]. The number of infusion related reactions reported in the study has been low; a total of 9 infusion related reactions (4 events in the DRL\_BZ arm and 5 events in the Avastin® arm) were reported. Majority of the reported infusion related reactions were mild to moderate in intensity. Overall 67 SAEs (27 in the DRL\_BZ arm, 40 in the Avastin® arm) were reported in 37 (31.6%) patients. of these, (17 (29.3%) patients were in the DRL\_BZ arm and 20 (33.9%) patients in the Avastin® arm. A total of 8 (6.8%) patients (4 patients in each arm) reported TEAEs resulting in death, none of them were related to study drug. A total of 30 (25.6%) patients were discontinued from the study due to TEAEs, of which 16 (27.6%) were in the DRL\_BZ arm and 14 (23.7%) in the Avastin® arm. Overall the safety profile of the patients in DRL\_BZ test arm was also in line the known safety profile of bevacizumab.

In NSCLC group, the incidence of TEAEs in the DRL\_BZ arm was 93.3% (28 patients) and in the Avastin® arm was 76.7% (23 patients). Eight (26.7%) patients in the DRL\_BZ arm as compared to nine (30.0%) patients in the Avastin® arm had at least one related TEAE. There were 17 patients with SAEs; 12 (40.0%) in the DRL\_BZ arm and 5 (16.7%) in the Avastin® arm. The most common TEAEs in both arms were **anaemia** [DRL\_BZ arm: 29 events in 15 (50%) patients, Avastin® arm: 26 events in 14 (46.7%) patients] and **asthenia** [12 events in 11 (36.7%) patients in each arm]. The following TEAEs occurred in >10% of patients in both treatment arms: thrombocytopenia, constipation, diarrhoea, nausea, stomatitis, pyrexia, decreased appetite and alopecia.

## 5.2 Pharmacokinetic properties

The pharmacokinetic data for bevacizumab are available from ten clinical trials in patients with solid tumours. In all clinical trials, bevacizumab was administered as an IV infusion. The rate of infusion was based on tolerability, with an initial infusion duration of 90 minutes. The pharmacokinetics of bevacizumab was linear at doses ranging from 1 to 10 mg/kg.

### Distribution

The typical value for central volume ( $V_c$ ) was 2.73 L and 3.28 L for female and male patients respectively, which is in the range that has been described for IgGs and other monoclonal antibodies. The typical value for peripheral volume ( $V_p$ ) was 1.69 L and 2.35 L for female and male patients respectively, when bevacizumab is co-administered with anti-neoplastic agents. After correcting for body weight, male patients had a larger  $V_c$  (+ 20%) than female patients.

### Biotransformation

Assessment of bevacizumab metabolism in rabbits following a single IV dose of  $^{125}$ I-bevacizumab indicated that its metabolic profile was similar to that expected for a native IgG molecule which does not bind VEGF. The metabolism and elimination of bevacizumab is similar to endogenous IgG i.e. primarily via proteolytic catabolism throughout the body, including endothelial cells, and does not rely primarily on elimination through the kidneys and liver. Binding of the IgG to the FcRn receptor results in protection from cellular metabolism and the long terminal half-life.

### Elimination

The value for clearance is, on average, equal to 0.188 and 0.220 L/day for female and male patients, respectively. After correcting for body weight, male patients had a higher bevacizumab clearance (+

17%) than females. According to the two-compartmental model, the elimination half-life is 18 days for a typical female patient and 20 days for a typical male patient.

Low albumin and high tumour burden are generally indicative of disease severity. Bevacizumab clearance was approximately 30% faster in patients with low levels of serum albumin and 7% faster in subjects with higher tumour burden when compared with a typical patient with median values of albumin and tumour burden.

#### Pharmacokinetics in special populations

The population pharmacokinetics were analysed in adult and pediatric patients to evaluate the effects of demographic characteristics. In adults, the results showed no significant difference in the pharmacokinetics of bevacizumab in relation to age.

##### *Renal impairment*

No trials have been conducted to investigate the pharmacokinetics of bevacizumab in renally impaired patients since the kidneys are not a major organ for bevacizumab metabolism or excretion.

##### *Hepatic impairment*

No trials have been conducted to investigate the pharmacokinetics of bevacizumab in patients with hepatic impairment since the liver is not a major organ for bevacizumab metabolism or excretion.

##### *Paediatric population*

The pharmacokinetics of bevacizumab were evaluated in 152 children, adolescents and young adults (7 months to 21 years, 5.9 to 125 kg) across 4 clinical studies using a population pharmacokinetic model. The pharmacokinetic results show that the clearance and volume of distribution of bevacizumab were comparable between paediatric and young adult patients when normalised by body weight, with exposure trending lower as body weight decreased. Age was not associated with the pharmacokinetics of bevacizumab when body weight was taken into account.

The pharmacokinetics of bevacizumab was well characterized by the paediatric population PK model for 70 patients in Study BO20924 (1.4 to 17.6 years; 11.6 to 77.5 kg) and 59 patients in Study BO25041 (1 to 17 years; 11.2 to 82.3 kg). In Study BO20924, bevacizumab exposure was generally lower compared to a typical adult patient at the same dose. In Study BO25041, bevacizumab exposure was similar compared to a typical adult at the same dose. In both studies, bevacizumab exposure trended lower as body weight decreased.

### **5.3 Preclinical safety data**

In studies of up to 26 weeks duration in cynomolgus monkeys, physeal dysplasia was observed in young animals with open growth plates, at bevacizumab average serum concentrations below the expected human therapeutic average serum concentrations. In rabbits, bevacizumab was shown to inhibit wound healing at doses below the proposed clinical dose. Effects on wound healing were shown to be fully reversible.

Studies to evaluate the mutagenic and carcinogenic potential of bevacizumab have not been performed.

No specific studies in animals have been conducted to evaluate the effect on fertility. An adverse effect on female fertility can however be expected as repeat dose toxicity studies in animals have shown inhibition of the maturation of ovarian follicles and a decrease/absence of corpora lutea and associated decrease in ovarian and uterus weight as well as a decrease in the number of menstrual cycles.

Bevacizumab has been shown to be embryotoxic and teratogenic when administered to rabbits. Observed effects included decreases in maternal and foetal body weights, an increased number of foetal resorptions and an increased incidence of specific gross and skeletal foetal malformations. Adverse foetal outcomes were observed at all tested doses, of which the lowest dose resulted in average serum concentrations approximately 3 times larger than in humans receiving 5 mg/kg every 2 weeks. Information on foetal malformations observed in the post marketing setting are provided in section 4.6 Fertility, Pregnancy and Lactation and 4.8 Undesirable Effects.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

$\alpha,\alpha$ -Trehalose dihydrate

Sodium Phosphate (monobasic, monohydrate)

Sodium Phosphate (dibasic, anhydrous)

Polysorbate 20

Water for injection

### **6.2 Incompatibilities**

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

A concentration dependent degradation profile of bevacizumab was observed when diluted with glucose solutions (5%).

### **6.3 Shelf life**

Vial (unopened)

3 years.

Diluted medicinal product

Chemical and physical in-use stability has been demonstrated at  $5 \pm 3$  °C or  $25 \pm 2$  °C up to 60 days. From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions are the responsibility of the user and would normally not be longer than 48 hours at  $5 \pm 3$  °C or  $25 \pm 2$  °C, unless dilution has taken place in controlled and validated aseptic conditions.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C-8°C).

Do not freeze.

Keep the vial in the outer carton in order to protect from light.

For storage conditions after dilution of the medicinal product, see section 6.3.

### **6.5 Nature and contents of container**

4 ml solution in a vial (Type I glass) with a stopper (chlorobutyl rubber) sealed with aluminum flip off seal containing 100 mg of bevacizumab

16 ml solution in a vial (Type I glass) with a stopper (chlorobutyl rubber) sealed with aluminum flip off seal containing 400 mg of bevacizumab.

Pack of 1 vial.

### **6.6 Special precautions for Use, Handling and Disposal**

Versavo should be prepared by a healthcare professional using aseptic technique to ensure the sterility of the prepared solution. A sterile needle and syringe should be used to prepare Versavo.

The necessary amount of bevacizumab should be withdrawn and diluted to the required administration volume with sodium chloride 9 mg/ml (0.9%) solution for injection. The concentration of the final bevacizumab solution should be kept within the range of 1.4 mg/ml to 16.5 mg/ml.

Parenteral medicinal products should be inspected visually for particulate matter and discolouration prior to administration.

No incompatibilities between Versavo and polyvinyl chloride or polyolefine bags or infusion sets have been observed.

Versavo is for single-use only, as the product contains no preservatives. Any unused medicinal product or waste material should be disposed in accordance with local requirements.

## **7. PRODUCT REGISTRATION HOLDER**

DR. REDDY'S LABORATORIES MALAYSIA SDN BHD, UNIT NO. SO-29-07 AND SO-29-08, MENARA 1, STRATA OFFICE, NO. 3, JALAN BANGSAR, KL ECO CITY, 59200 KUALA LUMPUR MALAYSIA

## **8 MANUFACTURER**

Dr. Reddy's Laboratories Ltd., BIOLOGICS., SURVEY NO.47 & 44 (PART), BACHUPALLY VILLAGE, BACHUPALLY MANDAL, MEDCHAL - MALKAJGIRI DISTRICT, PINCODE 500090, TELANGANA STATE, INDIA

## **9 DATE OF REVISION OF THE TEXT**

April 2025