



PENTONIL 200MG/ML INJECTION

Description and Composition

PENTONIL 200MG/ML INJECTION is a pink clear solution. Each ml contains 200mg Pentobarbital Sodium as its active ingredient and 10.4mg Benzyl Alcohol as its preservative.

Pharmacodynamics

Pentobarbital is a barbiturate. Barbiturates act by depressing the central nervous system, inhibiting accumulation of calcium in nervous tissue, which leads to liberation of norepinephrine, acetylcholine, glutamate and gamma-aminobutyric acid. This leads to sedation and coma.

Barbiturates also act on cardiac and respiratory systems bringing on apnoea and cardiac arrest.

Pharmacokinetics

After intravenous administration fast distribution over the tissues will occur.

Pentobarbital is mainly eliminated through the liver by biotransformation, particularly by the Cytochrome P₄₅₀ system, as well as by excretion in the kidneys and redistribution. In pigs redistribution in fatty tissue might cause reduced plasma concentrations and prolonged action. Barbiturates may diffuse through the placenta in foetal tissue, and traces of barbiturates may be present in the breast milk.

Indication

For euthanasia of small animals (mainly dogs and cats) and cattle.

Recommended Dose

0.7ml per kg body weight by rapid intravenous injection. The product solution is not intended for dilution with water or any other fluid.

Route of Administration

To be given by intravenous injection.

Contraindication

Not to be used in animals intended for animal consumption. Do not use for anaesthesia.

Warning and Precautions

Special warnings for each target species

None.

Special precautions for use

Special precautions for use in animals

Select the needle gauge based on the species and size of animal and check that the syringe and needle are mounted and secured correctly. It is important to ensure that any pressure created during delivery does not lead to the needle disconnecting from the syringe. The amount of the drug to be administered may mean that a smaller gauge needle may not deliver the necessary volume at an appropriate speed. Any volume administered outside the vein will reduce the efficacy of the dose.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Care should be taken to ensure the pressure on the syringe is not too great to avoid accidental spraying of the face and eyes (see 'Special precautions for use in animals'). Wear suitable protective gloves and glasses when handling the product. Avoid accidental self-administration and self-injection. In the case of accidental self-administration, by injection, ingestion or skin absorption, seek URGENT medical attention, advising medical service of barbiturate poisoning.

In the event of an accident the following action should be taken:

Skin - Wash immediately with water and then thoroughly with soap and water.

Eyes - Wash immediately with cold water and obtain medical attention.

Ingestion - Obtain medical attention immediately. Wash out mouth. Keep warm and rest.

Accidental self-injection - Obtain URGENT medical attention, advising medical services of barbiturate poisoning. Do not leave patient unattended.

Advice to Doctor - Maintain airways and give symptomatic and supportive treatment.

Interaction with Other Medicaments

The effects of the product are increased by concomitant administration of sedatives (xylazine or acepromazine).

Pregnancy and Lactation

Not applicable

Side Effects

None

Symptoms and Treatment of Overdose

In the event of accidental administration to an animal not presented for euthanasia, measures such as artificial respiration, administration of oxygen and the use of analeptics are appropriate.

Storage Condition

Store below 30°C.

Shelf life

3 years. Discard 6 months after first opening.

Withdrawal period

Not to be used in animals intended for human consumption.

Packing

20ml, 50ml, 100ml, 250ml

Manufacturer & Product Registration Holder

Range Pharma Sdn Bhd

No. 1, Jalan TSB 11, Taman Industri Sg. Buloh,
47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia.

Tel: 603-61568708 Fax: 603-61568707

Email: support@vetscareline.com

Revised date: 8.6.2026