

DEGREE IN COMPUTER ENGINEERING

PHYSICS EXAM

10th JANUARY 2011

Surname:

Name:

Group 89



E1 ☐ E2 ☐ E3 ☐ E4 ☐ Q1 ☐ Q2 ☐ Q3 ☐

- 1.- The exam has 4 exercises and 3 questions.
- 2.- The corresponding marks are attached to each exercise or question.
- 3.- Each exercise or question must be solved on a separate sheet.
- 4.- On each square above, mark with an X the exercises or questions that have **NOT** been solved.

CONSTANTS:

Charge of the electron: $-1,6 \times 10^{-19} \text{ C}$

Permittivity of free space: $\epsilon_0 = 8,85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$

Permeability of free space: $\mu_0 = 4 \pi \times 10^{-7} \text{ N A}^{-2}$

Mass of the electron: $9,11 \times 10^{-31} \text{ kg}$.

$h = 6.626 \times 10^{-34} \text{ Js}$

Mass of the proton: $1,67 \times 10^{-27} \text{ kg}$

$c = 2.998 \times 10^8 \text{ m/s}$

EXERCISES:


E1. (2p) A proton starting from rest is accelerated by a uniform electric field of magnitude 640 N/C. After some time, the speed of the proton is $1.2 \times 10^6 \text{ m/s}$.

- a) How long does it take for the proton to reach that speed?
- b) Find the distance travelled by the proton until it reaches that speed.
- c) Find the kinetic energy of the proton when it reaches that speed in J and eV.

E2. (2p) A small sphere of mass m and charge q hanging from a massless string is located inside a parallel-plate capacitor with plates of area A and distance between the plates d (as shown in the figure).


- a) Find the capacitance of the capacitor, the electric field inside the capacitor and the charge on the plates.
- b) Find the charge q of the sphere knowing that the pendulum is in equilibrium when the string forms an angle α with the vertical.

DATA: $A = 1.13 \text{ m}^2$; $d = 4 \text{ cm}$; $V_0 = 500 \text{ V}$, $m = 2 \text{ g}$; $\alpha = 30^\circ$



E3. (2p) A solid sphere of radius R_1 is uniformly charged with a volume charge density ρ_0 . A metallic spherical shell of inner radius R_2 and outer radius R_3 concentric with the solid sphere is charged with Q (see figure). Find the electric field \vec{E} (expressed in rectangular coordinates) at points: A (5,0,0); B (20,0,0); C (38,0,0) and D (25,35,0) (all coordinates are expressed in cm).

DATA: $\rho_0 = -4.8 \times 10^{-3} \text{ C/m}^3$; $R_1 = 10 \text{ cm}$; $R_2 = 30 \text{ cm}$; $R_3 = 40 \text{ cm}$; $Q = 30 \mu\text{C}$




E4. (2p) Three infinite conducting wires parallel to the Z axis carry currents $I_1 = 2 \text{ A}$, $I_2 = 5 \text{ A}$, $I_3 = 10 \text{ A}$. The location of the wires and the direction of the current intensities are shown in the figure attached.

a) Find \vec{B} at (0, 0, 0).

b) Find the magnetic force \vec{F} exerted on an electron located at (0,0,0) knowing that it travels with a velocity $\vec{v} = 3 \times 10^4 \vec{i} + 5 \times 10^4 \vec{j} \text{ (m/s)}$.

Express all vectors in rectangular coordinates.



QUESTIONS:

Q1. (1 p) Describe the conduction properties of an intrinsic semiconductor (both at 0 K and at $T > 0 \text{ K}$). Use the band theory of solids in the explanation.

Q2. (1p) Explain the physical behaviour of a semiconductor diode (see the characteristic curve attached). Consider both forward and reverse bias in the explanation.

