IMPROVED BOUNDS AND PROOF-CLASS BARRIERS FOR THE DIVISIBILITY ANTICHAIN SATURATION GAME (ERDŐS PROBLEM 872)

OM BUDDHDEV

Abstract. We study the saturation game on the divisibility poset of $\{2, 3, \ldots, n\}$. Two players alternately choose integers subject to the condition that the chosen set remains primitive; Prolonger moves first and maximizes the final number of moves, while Shortener minimizes it. We prove

$$\left(\frac{1}{8} - o(1)\right) \frac{n \log\log n}{\log n} \leq L(n) \leq \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n = (0.1897112 + o(1))n < 0.19n,$$

as Theorems 4.4 and 7.8. The lower bound follows from a two-phase Prolonger strategy and a bipartite maximum-degree capture lemma, while the upper bound uses an explicit fourth-order Bonferroni constant $\mathcal{W}_4$ coming from the local logarithmic density of Shortener's legal-prime prefix. We also prove a polynomial lower bound for the weighted shield functional, an exact $5n/24 + O(1)$ first-hit cover theorem for the lower half, and the intermediate upper bounds $13n/36 + o(n)$ and $5n/16 + o(n)$. A conditional lower bound of order $n(\log\log n)^2/\log n$ is included under a restricted finite safe-edge hypothesis which is stated explicitly. The paper also records three proof-class barriers for local relaxations, Johnson-shadow separators, and separator-only carrier strategies.

1. Introduction

Let $n \geq 2$. The divisibility antichain saturation game is played on

$$\{2, 3, \ldots, n\}.$$

Two players alternately choose previously unchosen integers, and the chosen set must remain an antichain under divisibility. Thus no chosen integer may divide another chosen integer. The game stops when the current antichain is maximal. Prolonger moves first and tries to maximize the final number of chosen integers; Shortener tries to minimize it. We denote the value of the game by $L(n)$.

This game is one of the finite saturation games arising from Erdős's questions on primitive sets [Erd92]; saturation games of this kind are part of a broader extremal-game tradition [FS92, BHW16]. A primitive set is precisely an antichain in the divisibility poset, and primitive sets have a long parallel history in multiplicative number theory [CE90, Lic23, MP10, Kuc24]. The upper half

$$U_n := \{m : \ n/2 < m \leq n\}$$

is the canonical maximum antichain, so the trivial scale is linear. The central question is whether optimal play forces a positive fraction of this linear scale, or whether Shortener can force $o(n)$ moves.

1.1. Prior work.

Erdős posed the problem in 1992 [Erd92], and it is catalogued as problem #872 in Bloom's list [Blo26]. No non-trivial upper bound appeared in the public record until early 2026. The game is a number-theoretic analogue of the triangle-free saturation game of Hajnal: for that game, Füredi and Seress [FS92] proved a lower bound of order $n \log n$, while Biró, Horn, and Wildstrom [BHW16] proved an upper bound below $0.215n^2$.

Date: April 21, 2026.

2020 *Mathematics Subject Classification.* Primary 91A46; Secondary 05D05, 11N25.

Key words and phrases. Saturation game, primitive set, divisibility poset, antichain, sieve methods, formal verification, Erdős problem 872.

The ambient combinatorial setting interacts with the literature on primitive sets and divisibility antichains. We use the primitive-set viewpoint of Cameron and Erdős [CE90], the weighted extremal perspective from Lichtman's proof of the Erdős primitive set conjecture [Lic23], and supporting multiplicative-structure estimates from [Kuc24, MP10].

The public upper-bound record for the present game developed rapidly in early 2026. Price and GPT-5.2 Pro gave the first linear upper bound,

$$L(n) \le \left(\tfrac{23}{48} + o(1)\right) n,$$

and introduced the terminology *Prolonger* and *Shortener* [PG26]. Subsequent forum refinements by Adenwalla, StijnC, natso26, Xiao_Hu, Desmond Weisenberg, and others improved this to

$$L(n) \le \left(\tfrac{419}{1008} + o(1)\right) n \le 0.416n$$

[Erd26]. The results in this paper substantially strengthen that record and add the first unconditional nontrivial lower bound.

1.2. **Main results.** Our main result is a new linear upper bound with an explicit constant below 0.19. Let

$$\rho(u) := \frac{1}{(\lfloor 1/u \rfloor + 1)u} \qquad (0 < u \le 1),$$

and define

$$J_r := \frac{1}{r!} \int_{\substack{u_1+\cdots+u_r \le 1 \\ 0<u_i\le 1}} \prod_{i=1}^{r} \rho(u_i)\, du_1 \cdots du_r \qquad (r \ge 1).$$

Set

$$\mathcal{W}_4 := 1 - J_1 + J_2 - J_3 + J_4.$$

The numerical value is

$$\frac{\mathcal{W}_4}{2} = 0.1897112\ldots < 0.19.$$

Theorem 1.1 (Main upper bound)**.** *Shortener has a strategy forcing*

$$L(n) \le \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n < 0.19n.$$

The proof of Theorem 1.1 is given in Section 7. Shortener plays a long prefix of the smallest legal odd primes. The local law governing which primes can be kept legal has density $\rho(u)$ on the logarithmic scale $p = n^u$. We build a monotone envelope for this density, invert it to obtain a comparison sequence, round the comparison sequence to genuine primes without losing the first four factorial moments, and then apply a fourth-order Bonferroni inequality to the odd-part compression of the final antichain.

Theorem 1.2 (Unconditional fan lower bound)**.** *Prolonger has a strategy forcing*

$$L(n) \ge \left(\frac{1}{8} - o(1)\right) \frac{n \log\log n}{\log n}.$$

The proof of Theorem 1.2 is given in Theorem 4.4. Prolonger first works through the small odd primes $a \le n^\delta$, making them unavailable by legal upper-half multiples unless Shortener chooses the singleton first. A harmonic charging argument shows that the small primes made unavailable by proper multiples still have reciprocal mass at least half of the total small-prime mass. The remaining upper-half targets ab form a fan graph, and a maximum-degree right-capture lemma forces at least half of the surviving fan edges to become actual game moves.

The lower-bound side also contains a weighted obstruction theorem for the standard shield-reduction approach. For $x \le n/2$, let

$$M_n(x) := \{u \in U_n : \ x \mid u\}, \qquad w_n(x) := |M_n(x)| - 1.$$

For $P \subseteq U_n$, put

$$\mathcal{L}(P) := \{x \le n/2 : \ x \nmid u \text{ for every } u \in P\},$$

and let

$$\beta_n(P) := \max\left\{\sum_{x\in B} w_n(x) : B \subseteq \mathcal{L}(P) \text{ is a divisibility antichain}\right\}.$$

Theorem 1.3 (Polynomial shield-weight lower bound)**.** *Fix $0 < \alpha < 1$. Uniformly for every $P \subseteq U_n$ with $|P| \le n^\alpha$,*

$$\beta_n(P) \ge \left(\frac{1}{2}\log\frac{1}{\alpha} - o(1)\right) n.$$

Equivalently, if

$$k_c^\star(n) := \min\{|P| : \ P \subseteq U_n, \ \beta_n(P) \le cn\},$$

then for every fixed $c > 0$,

$$k_c^\star(n) \ge n^{e^{-2c}-o(1)}.$$

The next theorem explains why this weighted quantity is the right one for shield-prefix arguments.

Theorem 1.4 (Shield reduction framework)**.** *Let A be any terminal antichain and let $P \subseteq A \cap U_n$. Then*

$$|A| \ge |U_n| - \beta_n(P).$$

Consequently, any proof which uses a prefix $P \subseteq U_n$ and this reduction to force $|A| \ge (1/2 - c)n + o(n)$ must have

$$|P| \ge n^{e^{-2c}-o(1)}.$$

The proof of the reduction is elementary and is given in Section 3; Theorem 1.3 is proved in Section 4. These structural results show that a short upper-half shield prefix cannot certify a linear lower bound through the weighted shadow method.

We also determine exactly the size of the first-hit cover of the lower half by upper-half multiples. Define

$$\tau(n) := \min\{|C| : \ C \subseteq U_n, \ \forall x \le n/2 \ \exists u \in C \text{ with } x \mid u\}.$$

Theorem 1.5 (Exact first-hit cover)**.**

$$\tau(n) = \frac{5}{24}n + O(1).$$

More precisely, the cover

$$H_n = \{u \in U_n : \ u \equiv 2 \pmod 4\} \cup \{u \in U_n : \ u > 2n/3, \ u \equiv 0 \pmod 4\}$$

has size $5n/24 + O(1)$, and there is a matching packing of lower-half witnesses of the same size up to $O(1)$.

Before the fourth-order argument, the same prime-prefix method gives two simpler upper bounds.

Theorem 1.6 (Intermediate upper bounds)**.** *Shortener has explicit strategies forcing*

$$L(n) \le \left(\frac{13}{36} + o(1)\right) n$$

and

$$L(n) \le \left(\frac{5}{16} + o(1)\right) n.$$

Finally, we include a conditional lower-bound theorem, not as a headline result but as a useful finite-combinatorial target. The natural general safe-edge hypothesis is false even inside the arithmetic residual hypergraphs, so the theorem is conditional on a narrower hypothesis for states generated by the activation and residual strategies in Section 4. Under that restricted hypothesis, the residual slot construction gives

$$L(n) \geq c\,\frac{n(\log\log n)^2}{\log n}$$

for some absolute $c > 0$. The finite graph, hypergraph, and arithmetic embedding cores of this theorem are formally verified; the restricted safe-edge hypothesis and the asymptotic activation wrapper remain in prose and are stated explicitly rather than suppressed.

Several parts of the paper are accompanied by Lean formalizations. The Shield Reduction theorem, the structural identities underlying the exact 5/24 cover, and the 13/36 upper-bound core have zero-sorry Lean verifications. Theorem A and the 5/16 upper bound have Lean artifacts whose remaining sorries are standard analytic-number-theory inputs or their assembly. The finite graph/hypergraph/embedding cores of the conditional T2 theorem are Lean-verified, while the restricted safe-edge hypothesis and asymptotic wrapper remain prose. For the $0.19n$ theorem, the endgame reduction from moment convergence to the strict inequality is zero-sorry Lean-verified; the moment-convergence hypotheses are the prose content of Section 7. A full status table appears in Appendix C.

The paper is organized as follows. Section 2 fixes notation. Section 3 proves the shield reduction. Section 4 proves Theorem A, the unconditional fan lower bound, and states the conditional T2 theorem. Section 5 proves the 5/24 cover theorem. Section 6 proves the 13/36 and 5/16 upper bounds. Section 7 proves the main $0.19n$ bound. Section 8 records three obstructions to extending the present methods. Deferred proofs, numerical details, and formalization metadata appear in the appendices.

2. Notation and Setup

All divisibility statements in this paper are inside the finite board

$$V_n := \{2, 3, \ldots, n\}.$$

A set $A \subseteq V_n$ is *primitive* if no two distinct elements of A are comparable under divisibility. Equivalently, A is an antichain in the divisibility poset. A primitive set is *maximal* if no element of $V_n \setminus A$ can be added while preserving primitiveness.

The game value $L(n)$ is the final size of the primitive set under optimal play when Prolonger moves first and maximizes the final size, while Shortener minimizes it.

We use the lower–upper split

$$L_n := \{m \in \mathbb{Z} :\ 2 \leq m \leq \lfloor n/2 \rfloor\}, \qquad U_n := \{m \in \mathbb{Z} :\ n/2 < m \leq n\}.$$

The upper half U_n is an antichain, since two distinct elements of U_n cannot divide each other.

For $x \in L_n$, the upper shadow and shadow weight are

$$M_n(x) := \{u \in U_n :\ x \mid u\}, \qquad w_n(x) := |M_n(x)| - 1.$$

The subtraction by 1 is convenient because a lower-half move x adds one move but may remove $|M_n(x)|$ possible upper-half moves from a terminal position.

For $P \subseteq U_n$, define

$$\mathcal{L}_n(P) := \{x \in L_n :\ x \nmid u \text{ for every } u \in P\},$$

and

$$\beta_n(P) := \max\left\{\sum_{x \in B} w_n(x) : B \subseteq \mathcal{L}_n(P) \text{ is a divisibility antichain}\right\}.$$

The maximum is over a finite set, and is understood to be 0 when $\mathcal{L}_n(P)$ is empty.

We use standard arithmetic notation. The functions $\pi(x)$ and $\vartheta(x)$ denote the prime-counting function and Chebyshev's function, respectively. For an integer m, $P^-(m)$ and $P^+(m)$ denote the least and greatest prime factors when $m > 1$, $\Omega(m)$ denotes the number of prime factors counted with multiplicity, and $\omega(m)$ denotes the number of distinct prime factors. The symbol $\mathbb{P}$ denotes the set of primes. As usual, $O(\cdot)$, $o(\cdot)$, $\ll$, and $\gg$ are interpreted as $n \to \infty$ unless a different parameter is explicitly specified; subscripts indicate dependence of the implied constants.

3. The Shield Reduction

The shield reduction turns terminal game positions into a weighted lower-half antichain problem.

Theorem 3.1 (Shield reduction)**.** *Let $A \subseteq V_n$ be a maximal primitive set and let $P \subseteq A \cap U_n$. Then*

$$|A| \geq |U_n| - \beta_n(P).$$

Proof. Put

$$B := A \cap L_n.$$

Then B is a divisibility antichain. Since $P \subseteq A$ and A is primitive, no $x \in B$ divides any element of P. Hence

$$B \subseteq \mathcal{L}_n(P).$$

We next determine $A \cap U_n$. We claim that

$$A \cap U_n = U_n \setminus \bigcup_{x \in B} M_n(x). \tag{3.1}$$

The inclusion "$\subseteq$" follows from primitiveness: if $u \in A \cap U_n$ and $x \in B$, then $x \nmid u$.

For the reverse inclusion, let $u \in U_n \setminus \bigcup_{x \in B} M_n(x)$. Suppose $u \notin A$. Since A is maximal, some $a \in A$ is comparable with u. If $u \mid a$, then $a \in U_n$ and $a < 2u$, so $a = u$, contradicting $u \notin A$. If $a \mid u$ and $a \in U_n$, again $a = u$. Thus $a \in L_n$, so $a \in B$, and $u \in M_n(a)$, contrary to the choice of u. This proves (3.1).

Using (3.1) and the union bound,

$$\begin{aligned} |A| &= |B| + |A \cap U_n| \\ &\geq |B| + |U_n| - \sum_{x \in B} |M_n(x)| \\ &= |U_n| - \sum_{x \in B} (|M_n(x)| - 1) \\ &= |U_n| - \sum_{x \in B} w_n(x). \end{aligned}$$

Since $B \subseteq \mathcal{L}_n(P)$ is an antichain, the final sum is at most $\beta_n(P)$. Therefore

$$|A| \geq |U_n| - \beta_n(P).$$

□

Corollary 3.2 (Shield-prefix obstruction)**.** *Fix $c > 0$. If a shield-prefix argument using Theorem 3.1 forces*

$$|A| \geq |U_n| - cn + o(n)$$

through a set $P \subseteq U_n$, then necessarily

$$|P| \geq n^{e^{-2c} - o(1)}.$$

Proof. Such an argument requires $\beta_n(P) \le cn + o(n)$. Applying Theorem 4.1 with $|P| \le n^\alpha$ gives

$$\frac{1}{2}\log\frac{1}{\alpha} \le c + o(1),$$

and hence $\alpha \ge e^{-2c} - o(1)$. □

4. Lower Bounds and Conditional Slot Construction

This section proves the polynomial shield-weight obstruction and an unconditional fan-capture lower bound for the game, and then states the conditional T2 lower bound. The conditional result is deferred to Appendix A.

4.1. The polynomial shield-weight lower bound.

Theorem 4.1 (Polynomial shield-weight lower bound)**.** *Fix $0 < \alpha < 1$. Uniformly over all $P \subseteq U_n$ with $|P| \le n^\alpha$,*

$$\beta_n(P) \ge \left(\frac{1}{2}\log\frac{1}{\alpha} - o(1)\right) n.$$

Proof. Fix δ with $\alpha < \delta < 1$. Let

$$Q_\delta(P) := \{p \le n^\delta : \ p \text{ prime and } p \nmid u \text{ for every } u \in P\}.$$

For all sufficiently large n, every $p \le n^\delta$ lies in L_n. The set $Q_\delta(P)$ is an antichain and is contained in $\mathcal{L}_n(P)$, so

$$\beta_n(P) \ge \sum_{p \in Q_\delta(P)} w_n(p). \tag{4.1}$$

For $p \le n^\delta$,

$$w_n(p) = \left\lfloor \frac{n}{p} \right\rfloor - \left\lfloor \frac{n}{2p} \right\rfloor - 1 = \frac{n}{2p} + O(1).$$

Since $\pi(n^\delta) = o(n)$, (4.1) gives

$$\beta_n(P) \ge \frac{n}{2} \sum_{p \in Q_\delta(P)} \frac{1}{p} - o(n). \tag{4.2}$$

Let

$$C_\delta(P) := \{p \le n^\delta : \ p \text{ prime and } p \mid u \text{ for some } u \in P\}.$$

Then $Q_\delta(P)$ and $C_\delta(P)$ partition the primes up to n^δ.

We first bound the reciprocal mass of $C_\delta(P)$. For each $p \in C_\delta(P)$ choose one $u(p) \in P$ divisible by p. For any fixed $u \in P$, the primes assigned to u are distinct divisors of u, so

$$\sum_{p:\, u(p)=u} \log p \le \log u \le \log n.$$

Summing over $u \in P$ gives

$$\sum_{p \in C_\delta(P)} \log p \le |P| \log n \le n^\alpha \log n. \tag{4.3}$$

Among all finite prime sets satisfying a log-budget $\sum_{p\in S} \log p \le B$, the reciprocal sum $\sum_{p\in S} 1/p$ is maximized by taking the smallest primes first. Indeed, if S contains a prime q and omits a smaller prime $p < q$, then replacing q by p decreases the log-cost and increases the reciprocal sum. Iterating this exchange proves the claim.

Let z_n be the largest value for which

$$\vartheta(z_n) = \sum_{p \le z_n} \log p \le n^\alpha \log n.$$

By (4.3) and the exchange argument,

$$\sum_{p\in C_\delta(P)} \frac{1}{p} \le \sum_{p\le z_n} \frac{1}{p}. \tag{4.4}$$

The prime number theorem gives

$$z_n = (1+o(1))n^\alpha \log n,$$

and hence

$$\log\log z_n = \log\log n + \log\alpha + o(1).$$

Mertens' theorem for primes yields

$$\begin{aligned}\sum_{p\in Q_\delta(P)} \frac{1}{p} &= \sum_{p\le n^\delta} \frac{1}{p} - \sum_{p\in C_\delta(P)} \frac{1}{p}\\ &\ge \sum_{p\le n^\delta} \frac{1}{p} - \sum_{p\le z_n} \frac{1}{p}\\ &= \log\log(n^\delta) - \log\log z_n + o(1)\\ &= \log\frac{\delta}{\alpha} + o(1).\end{aligned}$$

Substituting this into (4.2) gives

$$\beta_n(P) \ge \left(\frac{1}{2}\log\frac{\delta}{\alpha} - o(1)\right) n.$$

Letting $\delta \uparrow 1$ proves the theorem. □

Corollary 4.2 (Barrier exponent)**.** *For every fixed $c > 0$,*

$$k_c^\star(n) := \min\{|P| : \ P \subseteq U_n, \ \beta_n(P) \le cn\} \ge n^{e^{-2c}-o(1)}.$$

Proof. If $|P| \le n^\alpha$ and $\beta_n(P) \le cn$, then Theorem 4.1 gives

$$\frac{1}{2}\log\frac{1}{\alpha} \le c + o(1).$$

Thus $\alpha \ge e^{-2c} - o(1)$, which is the claimed exponent lower bound. □

4.2. **An unconditional fan-capture lower bound.** The shield obstruction above does not itself give a game lower bound. We now record a direct unconditional lower-bound strategy for Prolonger. The mechanism is different: Prolonger first makes a large reciprocal-mass set of small primes unavailable, then uses the resulting fan graph of upper-half products.

Lemma 4.3 (Maximum-degree right capture)**.** *Let $G = (L \sqcup R, E)$ be a finite bipartite graph. Consider the following finite capture game on the live graph. Maker moves first. On each Maker move, Maker chooses a right vertex of maximum positive live degree, captures all currently live edges incident to it, and removes that right vertex and its incident live edges. On each Breaker move, Breaker may delete one right vertex and all live edges incident to it, delete one live edge, or do nothing. The game stops when no live edge remains.*

Let C be the number of edges captured by Maker, and let X be the number of edges individually deleted by Breaker. Then, regardless of Breaker's play,

$$C + X \ge \frac{|E|}{2}.$$

Proof. Index Maker's nontrivial moves by i. Let d_i be the live degree of the right vertex captured by Maker on move i. Thus this move contributes d_i to C. Since Maker chose a maximum-degree right vertex, after Maker removes it every remaining right vertex has live degree at most d_i. Therefore, if Breaker replies by deleting a right vertex in the same round, that deletion removes at most d_i live edges. If Breaker instead deletes one edge or does nothing, the number of right-deleted edges in that round is 0.

Let D_{right} be the total number of edges deleted through right-vertex deletions. Summing the preceding round-by-round bound gives

$$D_{\text{right}} \leq \sum_i d_i = C.$$

Every original edge is eventually in exactly one of three classes: captured by Maker, deleted as part of a right-vertex deletion, or individually deleted by Breaker. Hence

$$|E| = C + D_{\text{right}} + X \leq 2C + X \leq 2(C + X),$$

which proves the claim. □

Theorem 4.4 (Unconditional fan-capture lower bound)**.** *For every fixed $0 < \delta < 1/2$ and every $\varepsilon > 0$, for all sufficiently large n, Prolonger has a strategy forcing*

$$L(n) \geq \left(\frac{1}{8} - \varepsilon\right) \frac{n \log\log n}{\log n}.$$

Equivalently,

$$L(n) \geq \left(\frac{1}{8} - o(1)\right) \frac{n \log\log n}{\log n}.$$

Proof. Fix $0 < \delta < 1/2$ and put $y := n^\delta$. Let

$$\mathcal{A} := \{a \leq y : \ a \in \mathbb{P}, \ a \text{ odd}\}.$$

For each $a \in \mathcal{A}$, define the right-prime interval

$$J_a := \left(\frac{n}{2a}, \frac{n}{a}\right] \cap \mathbb{P}.$$

If $b \in J_a$, then $ab \in U_n$. Also $b > y$ for all sufficiently large n, because $b > n/(2y) = \frac{1}{2}n^{1-\delta}$ and $\delta < 1/2$.

We shall use the fan graph with left vertices in $\mathcal{A}$, right vertices the primes appearing in the intervals J_a, and edges (a, b) representing the upper-half target ab.

Phase 1: activating small primes. As long as some prime in $\mathcal{A}$ is still a legal singleton at the start of a Prolonger turn, Prolonger takes the smallest such prime a and plays a legal target ab with $b \in J_a$. Such a target exists. Indeed, before any Phase 1 Prolonger move there have been $O(|\mathcal{A}|)$ previous Phase 1 moves, while uniformly for $a \in \mathcal{A}$ the prime number theorem gives

$$|J_a| = (1 + o_\delta(1)) \frac{n}{2a \log(n/a)} \gg_\delta \frac{n^{1-\delta}}{\log n}.$$

This is larger than $|\mathcal{A}| = O_\delta(n^\delta / \log n)$ because $\delta < 1/2$. If a is still legal, a previously chosen number can make a candidate ab illegal only by being b itself or ab itself; each previous move rules out at most one candidate b. Therefore some legal ab remains.

Phase 1 ends when no prime in $\mathcal{A}$ is a legal singleton. If this happens immediately after a Prolonger move, include Shortener's next move, if there is one, in the Phase 1 damage accounting below before Phase 2 starts. This convention only adds one Shortener move.

Define

$$A_0 := \{a \in \mathcal{A} : \text{ during Phase 1, } a \text{ is made illegal by a chosen proper multiple}\}$$

and

$$\mathcal{D} := \{a \in \mathcal{A} : \text{ Shortener chooses the singleton } a \text{ during Phase 1}\}.$$

Prolonger never plays a singleton from $\mathcal{A}$, and a prime singleton can become illegal only by being chosen itself or by having a proper multiple chosen. Thus

$$\mathcal{A} = A_0 \sqcup \mathcal{D}.$$

Let $P \subseteq A_0$ be the set of primes activated by Prolonger's Phase 1 moves. Pair each $a' \in \mathcal{D}$ with the immediately preceding Prolonger activation $a \in P$. The pairs are distinct because turns alternate. Since Prolonger always activates the smallest currently live prime, and since legality only decreases as play proceeds, $a \le a'$. Therefore

$$\sum_{a' \in \mathcal{D}} \frac{1}{a'} \le \sum_{a \in P} \frac{1}{a} \le \sum_{a \in A_0} \frac{1}{a}.$$

Using $\mathcal{A} = A_0 \sqcup \mathcal{D}$ and Mertens' theorem for primes,

$$\sum_{a \in A_0} \frac{1}{a} \ge \frac{1}{2} \sum_{a \in \mathcal{A}} \frac{1}{a} = \left(\frac{1}{2} - o(1)\right) \log\log n. \tag{4.5}$$

Phase 1 damage and the surviving fan. Let $E^\star$ be the raw fan-edge set over A_0:

$$E^\star := \{(a, b) : a \in A_0,\ b \in J_a\}.$$

The same uniform prime number theorem estimate gives

$$\begin{aligned} |E^\star| &= \sum_{a \in A_0} |J_a| \\ &\ge \left(\frac{1}{2} - o_\delta(1)\right) \frac{n}{\log n} \sum_{a \in A_0} \frac{1}{a} \\ &\ge \left(\frac{1}{4} - o_\delta(1)\right) \frac{n \log\log n}{\log n}, \end{aligned} \tag{4.6}$$

by (4.5).

We now remove from $E^\star$ the edges whose targets have already become unavailable before the first Phase 2 Prolonger move, and call the remaining edge set E_0. If $a \in A_0$, the singleton a was not chosen. Since $ab > n/2$, the target ab has no proper multiple in $\{2, \dots, n\}$, and its only relevant proper divisors are a and b. Hence a Phase 1 move can destroy a raw fan edge (a, b) only by choosing the right singleton b, which removes a whole right star, or by choosing the exact target ab, which removes one edge.

There are $O(|\mathcal{A}|)$ Phase 1 moves, including the possible intervening Shortener move. Right-singleton choices therefore destroy at most $O(|\mathcal{A}|^2)$ raw fan edges, and exact-target choices destroy at most $O(|\mathcal{A}|)$ raw fan edges. Since

$$|\mathcal{A}| = O_\delta\left(\frac{n^\delta}{\log n}\right),$$

the total damage is

$$O_\delta\left(\frac{n^{2\delta}}{\log^2 n}\right) = o_\delta\left(\frac{n}{\log n}\right) = o_\delta\left(\frac{n \log\log n}{\log n}\right).$$

Consequently

$$|E_0| \ge \left(\frac{1}{4} - o_\delta(1)\right) \frac{n \log\log n}{\log n}. \tag{4.7}$$

Phase 2: capturing right stars. On the surviving fan graph $G_0 = (A_0 \sqcup R_0, E_0)$, Prolonger follows the maximum-degree right-capture strategy from Lemma 4.3. That is, on each Phase 2 turn he

chooses a right prime b of maximum positive current degree and plays one legal target a_0b incident to it.

This move is legal. The left singleton a_0 is already illegal from Phase 1, but it was not chosen; the right singleton b has not been chosen, because otherwise the whole right star would have been removed from E_0; and the target a_0b itself is still available by the definition of E_0. Since $a_0b > n/2$, no proper multiple of it is on the board, and the only relevant proper divisors are a_0 and b.

Shortener's reply has one of the three effects allowed in Lemma 4.3: choosing a right singleton deletes a right vertex and its live star; choosing an exact fan target ab deletes that one live edge if it is still live, and is a do-nothing move relative to the live graph if ab was already captured in an earlier right-star capture by Prolonger; any other legal move deletes no live fan edge. Therefore the lemma applies. Let C be the number of fan edges captured by Prolonger's right-star moves in Phase 2, and let X be the number of live fan edges individually deleted by Shortener as exact targets in Phase 2. Then

$$C + X \geq \frac{|E_0|}{2}. \tag{4.8}$$

It remains to translate captured graph edges back to actual game moves. When Prolonger captures a right prime b by playing some a_0b, the capture-move edge (a_0, b) is realized as Prolonger's current game move. For every *other* edge (a, b) live at that capture time, with $a \neq a_0$, the singleton a is already illegal from Phase 1 but was not chosen, and the target ab has not yet been chosen. Moreover ab has no proper multiple in $\{2, \ldots, n\}$ and its only relevant proper divisors are a and b; any proper multiple of a that was chosen in Phase 1 is not comparable with this surviving target. Therefore ab remains legal until ab itself is chosen; terminal maximality forces it to be chosen eventually. Distinct fan targets all lie in U_n, so they are mutually incomparable unless they are equal, and the graph edges are distinct. Thus the captured edges account for C distinct eventual game moves (one of which, a_0b, is Prolonger's own capture play), while the individually deleted edges counted by X are themselves distinct game moves realized by Shortener's exact-target choices. Hence

$$L(n) \geq C + X \geq \frac{|E_0|}{2} \geq \left(\frac{1}{8} - o_\delta(1)\right) \frac{n \log\log n}{\log n},$$

by (4.7) and (4.8). Since δ was fixed, this proves the theorem. □

4.3. **The conditional T2 theorem.** The preceding fan-capture theorem gives an unconditional game lower bound. A different, potentially larger lower-bound mechanism uses residual upper-half targets of the form acb, where a, c are small primes and b lies in a corresponding large-prime interval. The resulting residual game is modeled by a scored rank-three slot hypergraph.

The finite potential argument which one would like to use in this slot hypergraph is not true for arbitrary finite capture games. The counterexample in Proposition A.1 shows that a unique max-gain Maker move need not dominate every subsequent Breaker deletion, and Proposition A.2 shows that the general safe-edge property already fails inside the residual arithmetic games. We therefore state the lower bound conditionally on a restricted property for the states generated by the specific activation and residual strategies.

Definition 4.5 (Restricted safe-edge hypothesis)**.** In each finite graph or scored rank-three hypergraph state actually reached by the T2 activation strategy and subsequent residual slot construction, assume that whenever a positive-weight live edge exists, Maker has a legal live edge f such that, after Maker plays f, every legal Breaker reply leaves the scaled potential Q at least as large as it was before Maker's move.

Theorem 4.6 (Conditional T2 lower bound)**.** *Assume the restricted safe-edge hypothesis for the finite graph and residual slot-hypergraph states generated by the construction in Appendix A. Then*

for every fixed $0 < \delta < 1/4$ *there is a constant* $c_\delta > 0$ *such that*

$$L(n) \geq c_\delta \frac{n(\log\log n)^2}{\log n}$$

for all sufficiently large n.

Proof roadmap. The full proof is in Appendix A. The argument has four ingredients.

First, the target family

$$\mathcal{T} = \{acb : \ a < c \leq n^\delta, \ b \in (n/(2ac), n/(ac)] \cap \mathbb{P}\}$$

has total mass

$$\sum_{a<c} |(n/(2ac), n/(ac)] \cap \mathbb{P}| \gg_\delta \frac{n(\log\log n)^2}{\log n}.$$

Second, an activation graph game secures a positive fraction of this mass, up to a deletion error which is

$$O\left(\frac{n^{4\delta}}{(\log n)^4}\right) = o\left(\frac{n(\log\log n)^2}{\log n}\right)$$

when $\delta < 1/4$. Third, once a pair (a, c) is secured, each residual target acb has only the three harmful slot divisors b, ab, cb; exact target plays score only their own target. Fourth, the restricted safe-edge hypothesis gives a one-eighth finite potential bound in the residual scored slot hypergraph. These ingredients produce a positive constant multiple of the initial target mass as actual game moves. □

5. The Exact 5/24 First-Hit Cover

This section determines the minimum number of upper-half elements needed to hit every lower-half element by divisibility.

Theorem 5.1 (First-hit cover constant)**.** *Let*

$$\tau(n) := \min\{|C| : \ C \subseteq U_n, \ \forall x \in L_n \ \exists u \in C \ \textit{with } x \mid u\}.$$

Then

$$\tau(n) = \frac{5}{24}n + O(1).$$

Proof. We prove matching upper and lower bounds.

For the upper bound, define

$$H_n := \{u \in U_n : \ u \equiv 2 \pmod 4\} \cup \{u \in U_n : \ u > 2n/3, \ u \equiv 0 \pmod 4\}.$$

We show that every $x \in L_n$ divides some element of H_n.

If $x > n/3$, then $2x \in U_n$. If x is odd, then $2x \equiv 2 \pmod 4$; if x is even, then $2x \equiv 0 \pmod 4$ and $2x > 2n/3$. Hence $2x \in H_n$.

If $n/4 < x \leq n/3$, then for odd x the element $2x$ lies in U_n and is 2 modulo 4. For even x, the element $3x$ satisfies $3x > 3n/4 > 2n/3$ and $3x \leq n$; it is even, and is therefore either 2 or 0 modulo 4, with the latter case covered by the condition $3x > 2n/3$. Thus $3x \in H_n$.

If $n/6 < x \leq n/4$, then $4x \in U_n$, $4x > 2n/3$, and $4x \equiv 0 \pmod 4$, so $4x \in H_n$.

Finally suppose $x \leq n/6$. The interval

$$\left(\frac{2n}{3x}, \frac{n}{x}\right]$$

has length $n/(3x) \geq 2$, and hence contains an even integer k. Then $u = kx$ satisfies $2n/3 < u \leq n$ and is even. If $u \equiv 2 \pmod 4$ then $u \in H_n$ by the first clause; if $u \equiv 0 \pmod 4$ then $u \in H_n$ by the second clause. In either case $x \mid u$.

The two clauses defining H_n are disjoint. Therefore

$$\begin{aligned}|H_n| &= \#\{u \in (n/2, n] : u \equiv 2 \pmod 4\} \\ &\quad + \#\{u \in (2n/3, n] : u \equiv 0 \pmod 4\} \\ &= \frac{n}{8} + \frac{n}{12} + O(1) = \frac{5n}{24} + O(1).\end{aligned}$$

Thus $\tau(n) \le 5n/24 + O(1)$.

For the lower bound, define

$$P_n := \{m : \ n/3 < m \le n/2\} \cup \{m : \ n/4 < m \le n/3, \ m \text{ odd}\}.$$

Then

$$|P_n| = \frac{n}{6} + \frac{1}{2}\left(\frac{n}{3} - \frac{n}{4}\right) + O(1) = \frac{5n}{24} + O(1).$$

The set P_n is an antichain. Indeed, if $x < y$ and $x \mid y$, then $y \ge 2x$. If $y \le n/3$, then $x \le n/6 < n/4$, impossible for $x \in P_n$. If $n/3 < y \le n/2$, then $x \le n/4$, again impossible for $x \in P_n$.

We claim that any $u \in U_n$ is divisible by at most one element of P_n. If $x \in P_n$ divides u, then $x > n/4$ and $u \le n$, so

$$1 < \frac{u}{x} < 4.$$

Thus u/x is either 2 or 3. If two distinct elements of P_n divide u, they must be $u/2$ and $u/3$. In particular $6 \mid u$. But then $u/3$ is even. It cannot lie in the first component $(n/3, n/2]$ because $u/3 \le n/3$, and it cannot lie in the second component because that component requires oddness. Hence at most one element of P_n divides u.

Now let $C \subseteq U_n$ be any cover of L_n. Each $x \in P_n$ must divide some element of C, and the preceding paragraph shows that different $x \in P_n$ require different elements of C. Therefore

$$|C| \ge |P_n| = \frac{5n}{24} + O(1).$$

This proves the matching lower bound. □

6. Intermediate Upper Bounds

The main upper bound in Section 7 refines a simpler odd-prime-prefix method. We present the two intermediate versions because their proofs isolate the compression and Bonferroni ideas used later.

6.1. Odd-part compression.

The following elementary lemma is used throughout the upper-bound arguments.

Lemma 6.1 (Odd-part injection)**.** *Let A be a divisibility antichain. The map*

$$\varphi(x) := \frac{x}{2^{v_2(x)}}$$

is injective on A.

Proof. If $\varphi(x) = \varphi(y) = m$, then $x = 2^a m$ and $y = 2^b m$. If $a < b$, then $x \mid y$; if $b < a$, then $y \mid x$. Since A is an antichain, both cases force $x = y$. □

6.2. The 13/36 bound.

Theorem 6.2. *Shortener has a strategy forcing*

$$L(n) \le \left(\frac{13}{36} + o(1)\right) n.$$

Proof. Let

$$k := \left\lfloor \frac{\sqrt{n}}{\log n} \right\rfloor.$$

For her first k turns, Shortener plays the smallest legal odd prime; after that she plays arbitrarily. Let

$$q_1 < q_2 < \cdots < q_k$$

be the odd primes she obtains.

We first bound these primes. Fix $\varepsilon > 0$. If, before Shortener's jth prefix move, every odd prime up to

$$X = (3/2 + \varepsilon) j \log n$$

were blocked, then the total odd-prime Chebyshev mass up to X would be covered by the earlier Shortener primes together with the odd prime divisors of Prolonger's first j moves. Each Prolonger move contributes at most $\log n$ of prime-divisor log-mass. By induction on j, the previous Shortener primes contribute at most

$$\sum_{i<j} \log q_i \le \left(\frac{1}{2} + o(1)\right) j \log n,$$

because $j \le \sqrt{n}/\log n$. Thus the blocked log-mass is at most $(3/2 + o(1)) j \log n$, while the prime number theorem gives

$$\vartheta_{\mathrm{odd}}(X) = (3/2 + \varepsilon + o(1)) j \log n,$$

a contradiction for large n. Hence

$$q_j \le (3/2 + \varepsilon) j \log n$$

uniformly for $j \le k$, and therefore

$$\sum_{j=1}^{k} \frac{1}{q_j} \ge \frac{1}{3} - o(1). \tag{6.1}$$

Let A be the final antichain. All elements other than the prefix primes have odd part not divisible by any q_j. By Lemma 6.1,

$$|A| \le k + N_D(n),$$

where $D = \{q_1, \ldots, q_k\}$ and $N_D(n)$ is the number of odd integers $m \le n$ not divisible by any prime in D.

Choose t minimal such that

$$s_t := \sum_{j \le t} \frac{1}{q_j} \ge \frac{1}{3} - o(1),$$

and put $E = \{q_1, \ldots, q_t\}$. Since every q_j is an odd prime, $1/q_j \le 1/3$, so

$$s_t \in \left[\frac{1}{3} - o(1),\ \frac{2}{3} + o(1)\right] \subset [0, 1]$$

for large n. Because $E \subseteq D$, we have $N_D(n) \le N_E(n)$.

Second-order Bonferroni on the events $q \mid m$ over odd $m \le n$ gives

$$N_E(n) \le \frac{n}{2}\left(1 - s_t + \frac{s_t^2}{2}\right) + O(t^2).$$

Here $t \le k$, so $O(t^2) = O(n/\log^2 n) = o(n)$. The function $f(s) = 1 - s + s^2/2$ is decreasing on $[0,1]$, and therefore

$$f(s_t) \le f(1/3 - o(1)) = \frac{13}{18} + o(1).$$

Consequently

$$|A| \le k + N_D(n) \le \frac{13}{36}n + o(n).$$

□

6.3. **The** 5/16 **bound.**

Theorem 6.3. *Shortener has a strategy forcing*

$$L(n) \le \left(\frac{5}{16} + o(1)\right) n.$$

Proof. Fix $A_0 > 2$ and set

$$k := \left\lfloor \frac{n}{2A_0 \log n} \right\rfloor.$$

Shortener plays the smallest legal odd prime for her first k turns. The same Chebyshev log-mass induction, now with $j \le n/(2A_0 \log n)$, gives

$$q_j \le A_0 j \log n \qquad (1 \le j \le k).$$

Indeed, if all odd primes below $A_0 j \log n$ were blocked, Prolonger's first j moves would contribute at most $j \log n$, and the previous Shortener primes would contribute at most $(1 + o(1)) j \log n$; this is less than $A_0 j \log n$ because $A_0 > 2$.

It follows that

$$\sum_{j=1}^{k} \frac{1}{q_j} \ge \frac{1}{A_0} - o(1).$$

As in the proof of Theorem 6.2, use the odd-part injection and truncate the prime set at the first partial sum

$$s_t \ge \frac{1}{A_0} - o(1).$$

Since $1/q_j \le 1/3$ and $A_0 > 2$, the partial sum remains in $[0,1]$. The second-order Bonferroni estimate gives

$$L(n) \le \frac{n}{2}\left(1 - \frac{1}{A_0} + \frac{1}{2A_0^2}\right) + o(n).$$

Letting $A_0 \downarrow 2$ yields

$$L(n) \le \frac{n}{2}\left(1 - \frac{1}{2} + \frac{1}{8}\right) + o(n) = \frac{5}{16}n + o(n).$$

The pair-intersection error is $o(n)$ because only prime pairs with product at most n contribute an $O(1)$ floor error, and

$$\#\{p < q : \ pq \le n\} \ll \sum_{p \le \sqrt{n}} \pi(n/p) \ll \frac{n}{\log n} \sum_{p \le \sqrt{n}} \frac{1}{p} = o(n).$$

□

7. The Main Upper Bound

In this section we prove the paper's headline upper bound

$$L(n) \le \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n < 0.19n.$$

The proof uses the following explicit strategy. Shortener plays the smallest legal odd prime for a long initial prefix of

$$K = \left\lfloor \frac{(1-\varepsilon)n}{2\log n} \right\rfloor$$

turns. Let

$$q_1 < q_2 < \cdots < q_K$$

be the odd primes she plays during that prefix. The argument has four layers:

(1) a local density law for the counting function of the q_j;
(2) a monotone envelope and inversion producing a comparison sequence $b_j \ge q_j$ with explicit moment limits;
(3) a fourth-order Bonferroni comparison theorem for arbitrary increasing odd-prime comparison sequences;
(4) a prime-rounding bridge that replaces the real sequence b_j by actual odd primes p_j, using the key half-density slack supplied by the prime-rounding bridge of Theorem 7.7.

7.1. The local density.

Write

$$S_K(X) := \#\{j \le K : q_j \le X\}, \qquad u := \frac{\log X}{\log n}.$$

On the interval

$$u \in \left(\frac{1}{h+1}, \frac{1}{h}\right],$$

the local prime-counting law identifies the limiting captured-prime density as

$$\rho(u) = \frac{1}{(\lfloor 1/u \rfloor + 1)u} = \frac{1}{(h+1)u}.$$

Lemma 7.1 (Local prime-count per range)**.** *Let Shortener follow the smallest-legal-odd-prime strategy. Fix an integer $h \ge 1$ and real numbers $2 \le Y < X \le n$ with*

$$Y^{h+1} > n.$$

Let $S(X)$ be the number of odd primes at most X which Shortener has played by the first time her next odd-prime move would exceed X (or by terminal time, if no such move occurs). Then

$$S(X) \ge \frac{\pi(X) - \pi(Y)}{h+1} - O_h(1).$$

In particular, if $n^{1/(h+1)} < Y < X \le n^{1/h}$, then every prime $p \in (Y, X]$ lies in the packet regime $p^h \le n < p^{h+1}$ and the same lower bound holds.

Proof. Shortener's odd-prime moves are increasing: once a prime has become illegal it cannot become legal later, and the strategy always chooses the smallest legal odd prime. At the stopping time in the definition of $S(X)$, no odd prime $p \le X$ is still legal. Therefore every odd prime $p \in (Y, X]$ has one of two forms: either Shortener herself has played p, or some earlier Prolonger move was divisible by p.

Let

$$M := \#\{p \in \mathbb{P} : \ Y < p \le X, \ p \text{ odd}\}.$$

The number of primes counted by M that Shortener has played is at most $S(X)$. There are at most $S(X) + 1$ Prolonger moves before the stopping time. Each such move is an integer at most n,

so it is divisible by at most h distinct primes from $(Y, X]$: otherwise it would be at least $Y^{h+1} > n$. Hence

$$M \le S(X) + h(S(X) + 1) = (h + 1)S(X) + h.$$

Thus

$$S(X) \ge \frac{M - h}{h + 1}.$$

Finally $M = \pi(X) - \pi(Y) + O(1)$, the $O(1)$ accounting only for the prime 2 and endpoint conventions. This gives the claimed bound. If $n^{1/(h+1)} < Y < X \le n^{1/h}$ and $p \in (Y, X]$, then $p^h \le X^h \le n$ while $p^{h+1} > Y^{h+1} > n$, so this is exactly the stated packet regime. □

Proposition 7.2 (Local density away from breakpoints)**.** *Fix $H \ge 2$, and set*

$$\tau_H := \frac{1}{8H^2}.$$

For $1 \le h \le H - 1$, define

$$\alpha_h := \frac{1}{h + 1} + \tau_H, \qquad \beta_h := \frac{1}{h} - \tau_H, \qquad I_h := [\alpha_h, \beta_h].$$

Then there exists a function $\xi_H(n) \to 0$ such that, for all sufficiently large n, for every $1 \le h \le H-1$ and every $u \in I_h$,

$$S_K(n^u) \ge \big(1 - \xi_H(n)\big)\frac{n^u}{(h + 1)u \log n}.$$

Proof. The intervals I_h are nonempty, because

$$\beta_h - \alpha_h = \frac{1}{h} - \frac{1}{h + 1} - 2\tau_H \ge \frac{1}{H(H - 1)} - \frac{1}{4H^2} > 0.$$

Fix $1 \le h \le H - 1$ and $u \in I_h$, and put

$$\eta := \frac{\tau_H}{2}, \qquad X := n^u, \qquad Y := n^{u-\eta}.$$

Since $u \in I_h$,

$$u - \eta \ge \frac{1}{h + 1} + \frac{\tau_H}{2} > \frac{1}{h + 1}, \qquad u \le \frac{1}{h} - \tau_H < \frac{1}{h}.$$

Hence the entire interval $[u - \eta, u]$ stays inside $\big(\frac{1}{h+1}, \frac{1}{h}\big)$. Thus $Y^{h+1} > n$, and Lemma 7.1 yields

$$S(X) \ge \frac{\pi(X) - \pi(Y)}{h + 1} - O_H(1),$$

where $S(X)$ denotes the number of odd primes played by Shortener up to the first time her move exceeds X.

We claim that $S(X) = S_K(X)$ for all $u \in \bigcup_{h=1}^{H-1} I_h$ once n is large enough. Indeed, $u \le \beta_1 = 1 - \tau_H$, so by the prime number theorem,

$$\pi(n^u) \ll \frac{n^{1-\tau_H}}{\log n} = o\left(\frac{n}{\log n}\right) = o(K).$$

Therefore all odd primes at most $X = n^u$ are exhausted before the prefix of length K, and the local counting function $S(X)$ coincides with the truncated counting function $S_K(X)$.

Substituting $X = n^u$ and $Y = n^{u-\eta}$ gives

$$S_K(n^u) \ge \frac{\pi(n^u) - \pi(n^{u-\eta})}{h + 1} - O_H(1).$$

Now u ranges over the fixed compact interval I_h, so the prime number theorem is uniform there:

$$\pi(n^u) = \big(1 + o_H(1)\big)\frac{n^u}{u \log n}.$$

Also $u-\eta \le u-\tau_H/2$, hence

$$\frac{\pi(n^{u-\eta})}{n^u/(u\log n)} \ll_H u\, n^{-\eta} = o_H(1)$$

uniformly for $u \in I_h$. Since there are only finitely many blocks $1 \le h \le H-1$, we may absorb the error terms into a single function $\xi_H(n) \to 0$, valid simultaneously for every h and every $u \in I_h$. This gives

$$S_K(n^u) \ge \big(1-\xi_H(n)\big)\frac{n^u}{(h+1)u\log n},$$

as claimed. □

7.2. **Moment constants and the Bonferroni coefficient.** Define

$$J_r := \frac{1}{r!}\int_{\substack{u_1+\cdots+u_r\le 1\\ u_i\in(0,1]}} \prod_{i=1}^{r}\rho(u_i)\,du_1\cdots du_r,$$

and

$$\mathcal{W}_4 := 1 - J_1 + J_2 - J_3 + J_4.$$

Numerical evaluation gives

$$\mathcal{W}_4 = 0.3794224, \qquad \frac{\mathcal{W}_4}{2} = 0.1897112 < 0.19.$$

For any increasing sequence $b_1 \le \cdots \le b_K$ of positive reals, define the truncated factorial-moment sums

$$T_r^{(b)}(n) := \sum_{\substack{1\le j_1<\cdots<j_r\le K\\ b_{j_1}\cdots b_{j_r}\le n}} \frac{1}{b_{j_1}\cdots b_{j_r}} \qquad (1\le r\le 4).$$

7.3. **Monotone envelope and inversion.** The local density from Proposition 7.2 is piecewise-defined, so one must build a monotone global comparison sequence before applying the Bonferroni comparison.

Proposition 7.3 (Envelope and inversion)**.** *Fix $H \ge 2$, and define $\tau_H, \alpha_h, \beta_h, I_h$ as in Proposition 7.2. Put*

$$G_H := \bigcup_{h=1}^{H-1} I_h,$$

and for $1 \le r \le 4$ define

$$J_r^{(H)} := \frac{1}{r!}\int_{\substack{u_1+\cdots+u_r\le 1\\ u_i\in G_H}} \prod_{i=1}^{r}\rho(u_i)\,du_1\cdots du_r.$$

Then for every sufficiently large n there exists a monotone function $C_{H,n} : [2,n] \to \{0,1,\dots,K\}$ such that:

(1) *$C_{H,n}(X) \le S_K(X)$ for all $2 \le X \le n$;*
(2) *if*
$$b_j := \inf\{X \in [2,n] : C_{H,n}(X) \ge j\},$$
then $q_j \le b_j$ for every $1 \le j \le K$;
(3) *for every fixed $r \in \{1,2,3,4\}$,*
$$T_r^{(b)}(n) = J_r^{(H)} + o_H(1) \qquad (n\to\infty);$$
(4) *for every fixed $r \in \{1,2,3,4\}$,*
$$0 \le J_r - J_r^{(H)} \le \frac{5r}{4H\,r!}.$$

In particular, there exists an integer-valued function $H_*(n) \to \infty$ and a corresponding choice of comparison sequences

$$b_1^{(n)} \le \cdots \le b_K^{(n)}$$

obtained from the above construction with $H = H_*(n)$ such that

$$T_r^{(b^{(n)})}(n) = J_r + o(1) \qquad (1 \le r \le 4).$$

Proof. By Proposition 7.2, there is a function $\xi_H(n) \to 0$ such that for every large enough n, every $1 \le h \le H-1$, and every $u \in I_h$,

$$S_K(n^u) \ge \big(1 - \xi_H(n)\big)\frac{n^u}{(h+1)u\log n}.$$

Define

$$A_{n,h}(u) := \big(1 - \xi_H(n)\big)\frac{n^u}{(h+1)u\log n} \qquad (u \in I_h).$$

For fixed H and all sufficiently large n, the functions $A_{n,h}$ are strictly increasing on I_h, since

$$A'_{n,h}(u) = \big(1 - \xi_H(n)\big)\frac{n^u}{(h+1)u}\left(1 - \frac{1}{u\log n}\right),$$

and $u \ge \alpha_{H-1} > 1/H$.

Set

$$L_h := \lfloor A_{n,h}(\alpha_h)\rfloor, \qquad R_h := \lfloor A_{n,h}(\beta_h)\rfloor \qquad (1 \le h \le H-1).$$

Then $L_h \le R_h$. Moreover, for $2 \le h \le H-1$,

$$\frac{A_{n,h-1}(\alpha_{h-1})}{A_{n,h}(\beta_h)} = n^{2\tau_H} \cdot \frac{(h+1)(1/h - \tau_H)}{h(1/h + \tau_H)} \to \infty,$$

so $R_h \le L_{h-1}$ for all large n. Also

$$R_1 \ll \frac{n^{1-\tau_H}}{\log n} = o(K),$$

again for large n. Thus, after increasing n if necessary,

$$0 \le L_{H-1} \le R_{H-1} \le L_{H-2} \le R_{H-2} \le \cdots \le L_1 \le R_1 < K.$$

We now define the cumulative lower envelope by

$$C_{H,n}(X) := \begin{cases} 0, & 2 \le X < n^{\alpha_{H-1}}, \\ \lfloor A_{n,h}(\log_n X)\rfloor, & n^{\alpha_h} \le X \le n^{\beta_h} \quad (1 \le h \le H-1), \\ R_h, & n^{\beta_h} < X < n^{\alpha_{h-1}} \quad (2 \le h \le H-1), \\ R_1, & n^{\beta_1} < X < n, \\ K, & X = n. \end{cases}$$

This function is monotone nondecreasing: it is nondecreasing on each genuine block because $A_{n,h}$ is increasing, it is constant on each gap, and the relations $R_h \le L_{h-1}$ guarantee monotonicity across adjacent pieces.

We next prove that $C_{H,n}(X) \le S_K(X)$ for all $2 \le X \le n$. If $X \in [n^{\alpha_h}, n^{\beta_h}]$, then

$$C_{H,n}(X) = \lfloor A_{n,h}(\log_n X)\rfloor \le S_K(X) \qquad \text{by the definition of } A_{n,h}.$$

If $X \in (n^{\beta_h}, n^{\alpha_{h-1}})$ with $2 \le h \le H-1$, then

$$C_{H,n}(X) = R_h \le S_K(n^{\beta_h}) \le S_K(X),$$

since S_K is nondecreasing. The same argument gives $C_{H,n}(X) = R_1 \le S_K(X)$ on (n^{β_1}, n), while $C_{H,n}(X) = 0 \le S_K(X)$ below $n^{\alpha_{H-1}}$ and $C_{H,n}(n) = K = S_K(n)$. This proves part (1).

Define

$$b_j := \inf\{X \in [2,n] : C_{H,n}(X) \geq j\} \qquad (1 \leq j \leq K).$$

Because $C_{H,n}$ is nondecreasing, the sequence $(b_j)_{j=1}^K$ is nondecreasing. Moreover,

$$S_K(b_j) \geq C_{H,n}(b_j) \geq j,$$

so the jth prime in the prime prefix satisfies $q_j \leq b_j$. This proves part (2).

For the moment asymptotics, write

$$u_j := \log_n b_j, \qquad w_j := \frac{1}{b_j} = n^{-u_j}, \qquad \mu_{H,n} := \sum_{j=1}^{K} w_j\, \delta_{u_j}.$$

Split $\mu_{H,n} = \nu_{H,n} + \lambda_{H,n}$, where $\nu_{H,n}$ is the contribution from the genuine interval blocks $L_h < j \leq R_h$, and $\lambda_{H,n}$ is the contribution from the flat filler blocks.

The filler blocks have negligible reciprocal mass. For the bottom block,

$$|\lambda_{H,n}^{\text{bot}}| = L_{H-1} n^{-\alpha_{H-1}} \leq A_{n,H-1}(\alpha_{H-1}) n^{-\alpha_{H-1}} = \frac{1-\xi_H(n)}{H\alpha_{H-1}\log n} = O_H\left(\frac{1}{\log n}\right).$$

For a gap block between I_h and I_{h-1}, where $2 \leq h \leq H-1$,

$$|\lambda_{H,n}^{(h)}| = (L_{h-1} - R_h) n^{-\alpha_{h-1}} \leq L_{h-1} n^{-\alpha_{h-1}} \leq \frac{1-\xi_H(n)}{h\,\alpha_{h-1}\log n} = O_H\left(\frac{1}{\log n}\right).$$

For the top block,

$$|\lambda_{H,n}^{\text{top}}| = (K - R_1) n^{-1} \leq \frac{K}{n} = O\left(\frac{1}{\log n}\right).$$

Since there are only $O_H(1)$ filler blocks,

$$|\lambda_{H,n}| = O_H\left(\frac{1}{\log n}\right) = o_H(1).$$

We next analyze the genuine blocks. Fix $1 \leq h \leq H-1$. On the block $L_h < j \leq R_h$, the value $u_j =: u_{h,j} \in I_h$ is defined by $A_{n,h}(u_{h,j}) = j$. Therefore, for every $f \in C^1([0,1])$,

$$\int f\, d\nu_{H,n}^{(h)} = \sum_{j=L_h+1}^{R_h} f(u_{h,j}) n^{-u_{h,j}} = \int_{\alpha_h}^{\beta_h} f(u) n^{-u}\, d\lfloor A_{n,h}(u)\rfloor.$$

Let

$$R_{n,h}(u) := \lfloor A_{n,h}(u)\rfloor - A_{n,h}(u), \qquad |R_{n,h}(u)| \leq 1.$$

Applying Stieltjes integration by parts with $g(u) := f(u)n^{-u}$ gives

$$\int_{\alpha_h}^{\beta_h} g(u)\, d\lfloor A_{n,h}(u)\rfloor = \int_{\alpha_h}^{\beta_h} g(u)\, dA_{n,h}(u) + E_{n,h}(f),$$

where

$$|E_{n,h}(f)| \leq 2\|g\|_\infty + \int_{\alpha_h}^{\beta_h} |g'(u)|\, du.$$

Since

$$g'(u) = \big(f'(u) - (\log n) f(u)\big) n^{-u},$$

we obtain

$$|E_{n,h}(f)| \leq 2\|f\|_\infty n^{-\alpha_h} + (\beta_h - \alpha_h)\|f'\|_\infty n^{-\alpha_h} + (\beta_h - \alpha_h)(\log n)\|f\|_\infty n^{-\alpha_h} = o_H(1).$$

On the other hand,

$$n^{-u} A'_{n,h}(u) = \big(1 - \xi_H(n)\big)\rho(u)\left(1 - \frac{1}{u\log n}\right) \qquad (u \in I_h),$$

because $\rho(u) = 1/((h+1)u)$ on I_h. Hence

$$\int_{\alpha_h}^{\beta_h} f(u) n^{-u}\, dA_{n,h}(u) = \big(1 - \xi_H(n)\big) \int_{\alpha_h}^{\beta_h} f(u)\rho(u) \left(1 - \frac{1}{u \log n}\right) du = \int_{I_h} f(u)\rho(u)\, du + o_H(1),$$

uniformly for fixed f. Combining the last two displays yields

$$\int f\, d\nu_{H,n}^{(h)} = \int_{I_h} f(u)\rho(u)\, du + o_H(1).$$

Summing over h and using $|\lambda_{H,n}| = o_H(1)$, we find that for every $f \in C^1([0,1])$,

$$\int f\, d\mu_{H,n} = \int_{G_H} f(u)\rho(u)\, du + o_H(1).$$

Taking $f \equiv 1$ gives $\mu_{H,n}([0,1]) = O_H(1)$.

Fix $r \in \{1,2,3,4\}$, and write

$$\Sigma_r := \{(u_1, \dots, u_r) \in [0,1]^r : u_1 + \cdots + u_r \le 1\}.$$

Then

$$\int_{\Sigma_r} d\mu_{H,n}^{\otimes r} = \sum_{j_1,\dots,j_r} \mathbf{1}_{\{u_{j_1} + \cdots + u_{j_r} \le 1\}} \prod_{t=1}^{r} w_{j_t}.$$

The contribution of distinct ordered r-tuples is exactly $r!\, T_r^{(b)}(n)$, because $u_{j_1} + \cdots + u_{j_r} \le 1$ is equivalent to $b_{j_1} \cdots b_{j_r} \le n$. The repeated-index contribution is negligible:

$$\max_j w_j \le n^{-\alpha_{H-1}} \to 0,$$

while $\mu_{H,n}([0,1]) = O_H(1)$ by the preceding $f \equiv 1$ bound. Therefore

$$0 \le \int_{\Sigma_r} d\mu_{H,n}^{\otimes r} - r!\, T_r^{(b)}(n) \le \binom{r}{2} \big(\max_j w_j\big) \mu_{H,n}([0,1])^{r-1} = o_H(1).$$

The same Riemann-sum convergence tensorizes: for every continuous F on $[0,1]^r$,

$$\int F\, d\mu_{H,n}^{\otimes r} \to \int_{G_H^r} F(u_1, \dots, u_r) \prod_{i=1}^{r} \rho(u_i)\, du_1 \cdots du_r.$$

The indicator of Σ_r restricted to G_H^r is Riemann integrable, because its discontinuities lie on the finitely many coordinate breakpoints and on the hyperplane $u_1 + \cdots + u_r = 1$. Approximating this indicator from above and below by continuous functions therefore gives

$$\int_{\Sigma_r} d\mu_{H,n}^{\otimes r} \to \int_{\substack{u_1 + \cdots + u_r \le 1 \\ u_i \in G_H}} \prod_{i=1}^{r} \rho(u_i)\, du_1 \cdots du_r = r!\, J_r^{(H)}.$$

Combining the last two displays proves part (3):

$$T_r^{(b)}(n) = J_r^{(H)} + o_H(1).$$

It remains to control the excised breakpoint region. Since $0 < \rho(u) < 1$ on $(0,1]$, we have

$$0 \le J_r - J_r^{(H)} \le \frac{1}{r!}\, \lambda_r\big(\{(u_1, \dots, u_r) \in [0,1]^r : \exists i,\ u_i \notin G_H\}\big),$$

where λ_r denotes r-dimensional Lebesgue measure. Moreover,

$$|(0,1] \setminus G_H| = \frac{1}{H} + 2(H-1)\tau_H \le \frac{1}{H} + \frac{1}{4H} = \frac{5}{4H}.$$

By the union bound in $[0,1]^r$,

$$\lambda_r\big(\{(u_1, \dots, u_r) \in [0,1]^r : \exists i,\ u_i \notin G_H\}\big) \le r\, |(0,1] \setminus G_H| \le \frac{5r}{4H}.$$

This proves part (4), and the final statement follows immediately by first letting $n \to \infty$ for fixed H, and then letting $H \to \infty$.

To obtain a single comparison sequence for each n, choose N_H so large that for all $n \geq N_H$ the above construction is valid and, for every $1 \leq r \leq 4$,

$$\left|T_r^{(b)}(n) - J_r^{(H)}\right| \leq \frac{1}{H}.$$

After enlarging if necessary, we may assume N_H is increasing in H. Define

$$H_*(n) := \max\{H \geq 2 : N_H \leq n\},$$

and let $b_1^{(n)} \leq \cdots \leq b_K^{(n)}$ be the comparison sequence produced with parameter $H = H_*(n)$. Then $H_*(n) \to \infty$, and for every $1 \leq r \leq 4$,

$$\left|T_r^{(b^{(n)})}(n) - J_r\right| \leq \left|T_r^{(b^{(n)})}(n) - J_r^{(H_*(n))}\right| + \left|J_r^{(H_*(n))} - J_r\right| \leq \frac{1}{H_*(n)} + \frac{5r}{4H_*(n)r!},$$

which tends to 0. This proves the final statement. □

7.4. **The prime-sequence Bonferroni comparison theorem.** The comparison theorem must be stated for *prime sequences*, not for arbitrary reals; the correct statement is the following.

Theorem 7.4 (Prime-sequence Bonferroni–4 comparison)**.** *Let $q_1 < \cdots < q_K$ be the actual odd primes played by Shortener during the prime prefix, and let $p_1 < \cdots < p_K$ be any increasing sequence of odd primes with*

$$q_j \leq p_j \qquad (1 \leq j \leq K).$$

Assume that for $1 \leq r \leq 4$,

$$T_r^{(p)}(n) = \Lambda_r + o(1).$$

Then

$$L(n) \leq \frac{n}{2}\left(1 - \Lambda_1 + \Lambda_2 - \Lambda_3 + \Lambda_4 + o(1)\right).$$

Proof. Let A be the final played set. If the game ends before Shortener completes the prefix of length K, then $L(n) = o(n)$ and there is nothing to prove, so assume $q_1, \ldots, q_K$ were all played.

Define the odd-part map

$$\varphi(x) := \frac{x}{2^{v_2(x)}}.$$

If $\varphi(x) = \varphi(y) = m$, then $x = 2^a m$ and $y = 2^b m$. If $a \neq b$, one of x and y divides the other, contradicting that A is an antichain. So φ is injective on A.

Since each $q_j \in A$, no other element of A can be divisible by any q_j. Because q_j is odd, divisibility by q_j is preserved under the odd-part map. Hence

$$\varphi(A) \subseteq \{m \leq n : m \text{ odd and } q_j \nmid m \text{ for all } j\} \cup \{q_1, \ldots, q_K\}.$$

Therefore

$$L(n) \leq N(q_1, \ldots, q_K) + K = N(q_1, \ldots, q_K) + o(n),$$

where $N(q_1, \ldots, q_K)$ counts odd integers up to n avoiding all q_j.

Now replace the actual sequence q_j by the larger prime sequence p_j. The count of odd integers avoiding a finite prime set is monotone in each prime, so

$$N(q_1, \ldots, q_K) \leq N(p_1, \ldots, p_K).$$

Applying fourth-order Bonferroni to the divisibility events $p_j \mid m$ over odd integers $m \leq n$ gives

$$N(p_1, \ldots, p_K) \leq \frac{n}{2}\left(1 - T_1^{(p)}(n) + T_2^{(p)}(n) - T_3^{(p)}(n) + T_4^{(p)}(n)\right) + O(M_2 + M_3 + M_4),$$

where M_r is the number of odd integers $\le n$ with at least r distinct prime divisors from the set $\{p_j\}$. The standard squarefree divisor-count bound gives

$$M_r = O_r\left(\frac{n(\log\log n)^{r-1}}{\log n}\right) = o(n) \qquad (2 \le r \le 4),$$

so the error is $o(n)$. Substituting $T_r^{(p)}(n) = \Lambda_r + o(1)$ yields the stated inequality. □

7.5. **The prime-rounding bridge.**

Lemma 7.5 (Fixed-ratio cell reciprocal mass)**.** *Fix $H \ge 2$, and let $b_1, \dots, b_K$ be the comparison sequence obtained from Proposition 7.3 with this fixed H. For every fixed $c > 1$, uniformly in $Y > 0$,*

$$\sum_{\substack{1\le j\le K\\ Y\le b_j<cY}} \frac{1}{b_j} = O_{H,c}\left(\frac{1}{\log n}\right).$$

Proof. If the cell $[Y, cY)$ lies inside a genuine block $[n^{\alpha_h}, n^{\beta_h}]$, then $B_{H,n}(X) := \#\{j : b_j \le X\}$ agrees with $\lfloor A_{n,h}(\log_n X)\rfloor$ on the block. As in the proof of Proposition 7.3,

$$\frac{d}{dX} A_{n,h}(\log_n X) = \frac{1-\xi_H(n)}{(h+1)\log X}\left(1 - \frac{1}{\log X}\right) \ll_H \frac{1}{\log n},$$

because every genuine block has $X \ge n^{\alpha_{H-1}} > n^{1/H}$. Hence

$$\#\{j : Y \le b_j < cY,\ j \text{ genuine}\} \ll_{H,c} \frac{Y}{\log n} + 1.$$

Dividing by Y gives a reciprocal contribution $O_{H,c}(1/\log n)$; the endpoint $+1$ is harmless since $Y \ge n^{1/H}$ in a genuine block.

A fixed-ratio cell can meet at most one breakpoint for all large n, because successive breakpoints in the envelope are separated by a factor which is a positive power of n. If the cell is adjacent to a breakpoint, split its genuine part into the at most two adjacent genuine pieces and apply the preceding estimate to each. The only additional contribution is the flat atom at the breakpoint. At $A_h := n^{\alpha_h}$ the flat multiplicity is at most L_h, so

$$\frac{L_h}{A_h} \le \frac{A_{n,h}(\alpha_h)}{n^{\alpha_h}} = O_H\left(\frac{1}{\log n}\right).$$

The top flat block at $b_j = n$ contributes at most

$$\frac{K}{n} = O\left(\frac{1}{\log n}\right).$$

Cells lying wholly in gaps have no genuine contribution and only the same possible flat-atom contribution. These cases cover all Y. □

It remains to convert the real comparison sequence b_j from Proposition 7.3 into an actual increasing odd-prime sequence. The naive claim

$$p_j/b_j = 1 + O(1/\log b_j)$$

uniformly in j is *not* justified by the prime number theorem alone and must not be used. The correct bridge is the half-density construction of Theorem 7.7.

Proposition 7.6 (Fixed-parameter queued prime rounding)**.** *Fix $H \ge 2$ and $\lambda > 1$, and choose $a > 1$ with $a^2 \le \lambda$. Let $b_1, \dots, b_K$ be the comparison sequence obtained from Proposition 7.3 with this fixed H. Then there is an increasing sequence of odd primes $p_1 < \cdots < p_K$ such that*

$$q_j \le b_j \le p_j \qquad (1 \le j \le K),$$

and, for every fixed $1 \le r \le 4$,

$$T_r^{(p)}(n) = T_r^{(b)}(n) + O_{H,r}(\lambda - 1) + o_{H,\lambda}(1). \tag{7.1}$$

Proof. Write

$$A_h := n^{\alpha_h}, \qquad B_h := n^{\beta_h} \qquad (1 \le h \le H-1).$$

Fix one genuine block $[A_h, B_h]$ and abbreviate $A = A_h$, $B = B_h$. Partition it into fixed multiplicative bins

$$I_k = [Aa^k, Aa^{k+1}) \qquad (k = 0, 1, \dots, M),$$

where the last bin is truncated at B. Let

$$d_k := \#\{j : b_j \in I_k,\ j \text{ lies in the genuine } h\text{-block}\}$$

and let f_h be the multiplicity of the flat atom at the left endpoint A_h. Thus f_{H-1} is the bottom flat atom, and for $h < H-1$ it is the flat atom in the gap just below A_h.

We first record the fixed-ratio estimates used by the queue. For every fixed $c > 1$, uniformly for $X \ge n^{1/H}$ with $[X, cX] \subseteq [A_h, B_h]$,

$$\#\{j : X \le b_j < cX,\ j \text{ genuine in block } h\} \le (1 + o_{H,c}(1)) \frac{(c-1)X}{(h+1)\log X} + O(1). \tag{7.2}$$

Indeed, on a genuine block $B_{H,n}(Y) = \lfloor A_{n,h}(\log_n Y) \rfloor$, and

$$\frac{d}{dY} A_{n,h}(\log_n Y) = \frac{1 - \xi_H(n)}{(h+1)\log Y} \left(1 - \frac{1}{\log Y}\right).$$

Integrating over $[X, cX]$ gives (7.2); the endpoint error is negligible at the scales used below. Similarly,

$$f_h \le (1 + o_H(1)) \frac{A_h}{(h+1)\log A_h}, \tag{7.3}$$

because $f_h \le L_h \le A_{n,h}(\alpha_h)$. Finally, if

$$P_k := \#\{p \text{ odd prime} : Aa^k \le p < Aa^{k+1}\},$$

then the prime number theorem in fixed multiplicative intervals gives, uniformly in k in this fixed block,

$$P_k = (1 + o_{H,a}(1)) \frac{(a-1)Aa^k}{\log(Aa^k)}. \tag{7.4}$$

Choose a small fixed $\eta > 0$. Since the demand has the factor $1/(h+1) \le 1/2$, we may choose an integer $s = s(a)$ such that

$$(1 - \eta)(a^{s+1} - 1) > (1 + \eta) \frac{a^s}{2}. \tag{7.5}$$

This is possible for every $a > 1$. For all large n, (7.2), (7.3), and the fact that s is fixed give

$$f_h + \sum_{k=0}^{s-1} d_k \le (1 + \eta) \frac{Aa^s}{2\log A},$$

while (7.4) gives

$$\sum_{\ell=0}^{s} P_\ell \ge (1 - \eta) \frac{A(a^{s+1} - 1)}{\log A}.$$

By (7.5), the queue containing the flat atom and the first s genuine bins clears by the end of prime bin s. More explicitly, the flat atom is eligible for primes in bin 0, and genuine demand from I_k is released only at prime bin $k+1$; processing prime bins in increasing order and assigning available

primes to the oldest unassigned demand therefore assigns all these early demands to primes at least as large as their corresponding b_j.

For each later full source bin I_k, $k \geq s$, assign its d_k elements to primes in the next prime bin $[Aa^{k+1}, Aa^{k+2})$. This is possible for all large n, because (7.2) and (7.4) give

$$d_k \leq (1 + o_{H,a}(1)) \frac{(a-1)Aa^k}{(h+1)\log(Aa^k)} < (1 - o_{H,a}(1)) \frac{(a-1)Aa^{k+1}}{\log(Aa^{k+1})} \leq P_{k+1},$$

using $1/(h+1) \leq 1/2 < a$. The final truncated source bin is handled in the same way, using the next full prime bin above the block.

Call exceptional the flat atoms, the first $s(a)$ genuine bins of every genuine block, and the top flat block $b_j = n$. Let E be the set of exceptional indices. By Lemma 7.5, each fixed multiplicative bin has b-reciprocal mass $O_{H,a}(1/\log n)$, and there are only $O_{H,a}(1)$ exceptional bins. The top flat block contributes $K/n = O(1/\log n)$. Thus

$$\sum_{j \in E} \frac{1}{b_j} = O_{H,a}\left(\frac{1}{\log n}\right). \tag{7.6}$$

Since every assigned prime is at least the corresponding b_j, the same bound holds with p_j in place of b_j.

The top flat block is assigned, in order, to odd primes in $[n, 2n]$. This interval contains $(1 + o(1))n/\log n$ odd primes, more than enough for the at most $K \leq n/(2\log n)$ top flat indices. Assignments from different genuine blocks do not overlap. The largest prime used for block h is at most $a^2 B_h$, while the next larger block begins at

$$A_{h-1} = n^{1/h+\tau_H} = n^{2\tau_H} B_h,$$

so $a^2 B_h < A_{h-1}$ for large n. Also $a^2 B_1 = o(n)$, so the top flat assignment above n comes last. The construction therefore produces one increasing prime sequence. For every nonexceptional index,

$$b_j \leq p_j \leq a^2 b_j \leq \lambda b_j. \tag{7.7}$$

Together with Proposition 7.3, this also gives $q_j \leq b_j \leq p_j$.

It remains to compare moments. The total reciprocal mass of the b_j is $O_H(1)$ by Proposition 7.3, and the same is true for the p_j by (7.6) and (7.7). Hence all r-tuples containing at least one exceptional index contribute

$$O_{H,r,a}\left(\frac{1}{\log n}\right) = o_{H,\lambda}(1)$$

to either $T_r^{(b)}(n)$ or $T_r^{(p)}(n)$.

For a nonexceptional tuple $I = \{j_1 < \cdots < j_r\}$, write

$$b_I := \prod_{j \in I} b_j, \qquad p_I := \prod_{j \in I} p_j.$$

By (7.7),

$$b_I \leq p_I \leq \lambda^r b_I, \qquad \lambda^{-r} \frac{1}{b_I} \leq \frac{1}{p_I} \leq \frac{1}{b_I}.$$

On tuples counted by both cutoffs, this changes the total reciprocal weight by $O_{H,r}(\lambda - 1)$, since the r-fold reciprocal mass is $O_H(1)$. The only cutoff discrepancy comes from tuples with

$$n/\lambda^r < b_I \leq n.$$

By the product-strip estimate proved from Lemma 7.5 above, these tuples have total b-reciprocal contribution $o_{H,\lambda}(1)$. This proves (7.1). □

Theorem 7.7 (Prime-rounding bridge)**.** *Fix $0 < \varepsilon < 1$, and let*

$$K = \left\lfloor \frac{(1-\varepsilon)n}{2\log n} \right\rfloor.$$

There exists a choice of comparison sequences

$$b_1^{(n)} \le \cdots \le b_K^{(n)}$$

from the construction of Proposition 7.3, and there exists an increasing sequence of odd primes

$$p_1 < \cdots < p_K$$

such that

$$q_j \le p_j \qquad (1 \le j \le K)$$

and

$$T_r^{(b^{(n)})}(n) = J_r + o(1) \qquad (1 \le r \le 4).$$

Moreover the primes may be chosen so that, for each fixed $1 \le r \le 4$,

$$T_r^{(p)}(n) = T_r^{(b^{(n)})}(n) + o(1) = J_r + o(1).$$

Proof. Choose a sequence $H_m \to \infty$ such that

$$|J_r^{(H_m)} - J_r| \le \frac{1}{m} \qquad (1 \le r \le 4).$$

For each m, choose $\lambda_m > 1$ so close to 1 that the $O_{H_m,r}(\lambda_m - 1)$ term in Proposition 7.6 is at most $1/m$ for $1 \le r \le 4$, and choose $a_m > 1$ with $a_m^2 \le \lambda_m$. For fixed m, enlarge N_m so that, for all $n \ge N_m$, the construction of Proposition 7.3 with $H = H_m$ is valid,

$$|T_r^{(b)}(n) - J_r^{(H_m)}| \le \frac{1}{m}, \qquad |T_r^{(p)}(n) - T_r^{(b)}(n)| \le \frac{2}{m}$$

for $1 \le r \le 4$, where the second inequality uses Proposition 7.6. We may take the N_m increasing. For $N_m \le n < N_{m+1}$ use the parameters $H = H_m$, $\lambda = \lambda_m$, and $a = a_m$. Then $H_m \to \infty$ and $\lambda_m \downarrow 1$, so the displayed bounds give

$$T_r^{(b^{(n)})}(n) = J_r + o(1), \qquad T_r^{(p)}(n) = T_r^{(b^{(n)})}(n) + o(1) = J_r + o(1)$$

for every fixed $1 \le r \le 4$. The inequality $q_j \le p_j$ follows from Proposition 7.6. □

7.6. **The finite Bonferroni–4 theorem.**

Theorem 7.8. *Under the piecewise-density prime-prefix strategy,*

$$L(n) \le \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n = (0.1897112 + o(1))n < 0.19n.$$

Proof. By Theorem 7.7, there are a comparison sequence $b_1 \le \cdots \le b_K$ and an increasing odd-prime sequence p_j with $q_j \le p_j$ such that

$$T_r^{(b)}(n) = J_r + o(1) \qquad (1 \le r \le 4).$$

The same theorem gives

$$T_r^{(p)}(n) = J_r + o(1) \qquad (1 \le r \le 4).$$

Applying Theorem 7.4 with $\Lambda_r = J_r$ gives

$$L(n) \le \frac{n}{2}\left(1 - J_1 + J_2 - J_3 + J_4 + o(1)\right) = \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n.$$

Numerical evaluation yields

$$\frac{\mathcal{W}_4}{2} = 0.1897112 < 0.19,$$

so the theorem follows. □

Remark 7.9. The corresponding formal artifact verifies the endgame reduction: once the first four moment terms are sufficiently close to the certified constants $J_1, \dots, J_4$, the strict inequality $L(n) < 0.19n$ follows eventually. Theorems 7.4 and 7.7 supply those moment hypotheses.

8. Obstructions to Further Improvement

This section records three barriers encountered by natural attempts to improve the upper bound. They are not used in the proof of Theorem 7.8; their purpose is to delineate which proof classes cannot by themselves resolve the asymptotic question.

8.1. **A local-relaxation barrier.**

The first obstruction says that bounded-round local relaxations can have an integrality gap at precisely the scale where one would hope for a short transversal argument. We state the barrier in its cleanest finite form.

Let $K_N^{(\ell)}$ be the complete ℓ-uniform hypergraph on the vertex set $[N]$. A transversal is a set of vertices meeting every ℓ-edge.

Proposition 8.1 (Sherali–Adams transversal barrier)**.** *Let $\ell = \lfloor N/\log N \rfloor$. The integral transversal number of $K_N^{(\ell)}$ is*

$$\tau(K_N^{(\ell)}) = N - \ell + 1 = N - o(N).$$

The fractional transversal number is

$$\tau_f(K_N^{(\ell)}) = \frac{N}{\ell} = (1 + o(1)) \log N.$$

Moreover, after any fixed number $r < \ell$ of Sherali–Adams rounds, the symmetric lifted solution still has objective at most

$$\frac{N}{\ell - r}.$$

Proof. A set $T \subseteq [N]$ meets every ℓ-set if and only if its complement has size at most $\ell - 1$. Hence

$$\tau(K_N^{(\ell)}) = N - \ell + 1.$$

For the fractional relaxation, give every vertex weight $1/\ell$. Each ℓ-edge then has total weight 1, so $\tau_f \le N/\ell$. Conversely, summing the edge constraints over all ℓ-subsets gives

$$\binom{N-1}{\ell-1} \sum_{i=1}^{N} x_i \ge \binom{N}{\ell},$$

and therefore $\sum_i x_i \ge N/\ell$. This proves $\tau_f = N/\ell$.

For the lifted statement, the same symmetric solution remains feasible after conditioning on any pattern involving fewer than ℓ vertices. If r vertices of an ℓ-edge have already been fixed outside the transversal, the residual constraint asks for weight at least 1 on an $(\ell - r)$-set; the uniform residual assignment $1/(\ell - r)$ satisfies it. The corresponding objective is $N/(\ell - r)$. This is the standard symmetric feasible solution for the r-round Sherali–Adams lift, and gives the claimed upper bound on the relaxation value. □

The proposition shows that local linear relaxations can remain logarithmic in size while the actual separator has linear size.

8.2. **A q-shadow dichotomy.** The next obstruction is an extremal statement for Johnson shadows. It explains why a sparse separator family cannot be forced by expansion alone unless a large pre-existing shadow is already present.

Let $X = \binom{[K]}{q}$ and $Y = \binom{[K]}{h}$, with $q < h$. A q-set Q covers an h-set H if $Q \subseteq H$. Let $R \subseteq Y$ be a live packet of density

$$\eta := \frac{|R|}{|Y|}.$$

Let $D \subseteq X$ be a previously forbidden q-shadow and let $C \subseteq X$ be a set of already claimed q-separators. Define

$$\delta_q := \frac{\binom{K-q}{h-q}}{\binom{K}{h}},$$

the density of h-sets containing a fixed q-set, and write $\sigma_q(D)$ for the fraction of Y covered by D. The normalized second singular value of the q-versus-h inclusion graph satisfies

$$\lambda_q^2 = \frac{q(K-h)}{h(K-q)}.$$

Proposition 8.2 (q-shadow covering dichotomy)**.** *Assume no legal q-separator captures at least*

$$\frac{1}{2}\,\delta_q|R|$$

members of R. Then

$$\sigma_q(D) + |C|\delta_q > 1 - \frac{4\lambda_q^2}{\eta}.$$

Proof. Let $A \subseteq X$ be the legal q-separators, meaning those not already excluded by D or C. By the union bound,

$$\frac{|A|}{|X|} \geq 1 - \sigma_q(D) - |C|\delta_q.$$

The expander-mixing inequality for the inclusion graph between X and Y gives

$$\left| e(A,R) - \frac{|A|}{|X|}\frac{|R|}{|Y|} e(X,Y) \right| \leq \lambda_q\, e(X,Y) \sqrt{\frac{|A|}{|X|}\frac{|R|}{|Y|}}.$$

If

$$\frac{|A|}{|X|} \geq \frac{4\lambda_q^2}{\eta},$$

then the average number of elements of R captured by a member of A is at least half the random expectation, namely at least $\frac{1}{2}\delta_q|R|$. This contradicts the hypothesis that no legal separator captures that many live packet members. Hence

$$\frac{|A|}{|X|} < \frac{4\lambda_q^2}{\eta},$$

and the displayed conclusion follows from the union bound. □

In the central regime relevant to top-facet packet arguments, the inequality says that separator starvation is possible only after the previous shadow or the already claimed separator set has become large. Pure expansion cannot supply an unlimited sequence of fresh separators.

8.3. **A separator-only limitation.** The last obstruction is a finite model for strategies which only play odd carrier separators and ignore dyadic shielding.

Proposition 8.3 (Separator-only limitation)**.** *Fix a finite set P of odd primes and a family $\mathcal{S} \subseteq 2^P$ of odd squarefree carrier supports. Suppose Shortener's planned moves are only odd squarefree separators with supports in $\mathcal{S}$. Then there are dyadic upper-half targets of the form*

$$2^a \prod_{p \in T} p$$

which remain untouched by all separators unless $\mathcal{S}$ covers the corresponding support T itself. In particular, any separator-only proof that does not control all dyadic lifts of the carrier family leaves a positive residual packet.

Proof. An odd separator can only be comparable with an upper-half target $2^a m$ through its odd part m. The dyadic exponent is invisible to the odd separator. Therefore, if a support T is not covered by the chosen odd separator family, every dyadic lift $2^a \prod_{p \in T} p$ with value in $(n/2, n]$ remains incomparable with all separator moves.

For any fixed odd part $m \leq n/2$, there is at least one dyadic lift $2^a m \in (n/2, n]$ unless $m = 1$; choose a maximal with $2^a m \leq n$. If $2^a m \leq n/2$, then $2^{a+1} m \leq n$, contradicting maximality. Thus each uncovered odd support produces an upper-half target not hit by the separator-only family. Summing over an uncovered packet gives the claimed residual. □

The proposition is deliberately modest: it does not rule out separator methods which also track dyadic lifts or exact upper targets. It rules out only the separator-only abstraction in which odd carrier supports are treated as if they were the whole divisibility geometry.

9. Conclusion

The main unconditional bounds of Theorems 4.4 and 7.8 bracket the divisibility antichain saturation game as

$$\left(\frac{1}{8} - o(1)\right) \frac{n \log\log n}{\log n} \leq L(n) \leq \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n.$$

The lower bound comes from a two-phase fan-capture strategy, while the upper bound improves the elementary linear picture by using the local density of Shortener's legal-prime prefix and a fourth-order Bonferroni comparison.

The conditional T2 theorem points toward a substantially larger lower-bound scale, but it is deliberately stated under a restricted safe-edge hypothesis: the general safe-edge property is false even inside the residual arithmetic games. The finite graph, hypergraph, and embedding components are robust; the remaining issue is whether the safe-edge property holds for states reached under the Prolonger activation strategy.

On the structural side, the shield reduction and Theorem A show that short upper-half shield prefixes cannot prove a linear game lower bound through the weighted shadow functional $\beta_n(P)$.

Three obstructions in Section 8 explain why the next improvements are unlikely to come from purely local relaxations, fixed shadow-cover dichotomies, or separator-only strategies. A sharper upper bound will need to use dynamic information about the actual divisibility game, not merely the static incidence geometry of low-rank shadows.

The central question remains the sharp order of growth of $L(n)$, and in particular whether $L(n) = o(n)$. The methods here give new constants, exact auxiliary structure, and a set of formal targets for future work, but they do not settle that asymptotic dichotomy.

Appendix A. Deferred Proofs

A.1. **Conditional proof of the T2 lower bound.** This appendix proves the conditional lower bound in Theorem 4.6. The proof is self-contained: all target families, auxiliary games, weights, activation rules, deletion errors, and asymptotic estimates are defined below. The opening counterexamples explain why the restricted finite safe-edge hypothesis is stated explicitly rather than hidden inside a max-gain argument.

Throughout this appendix n is large, $0 < \delta < 1/4$ is fixed, and

$$Y := n^{\delta}.$$

Let $\mathcal{P}_Y$ be the set of odd primes at most Y. For $a < c$ in $\mathcal{P}_Y$, put

$$I_{a,c} := \left(\frac{n}{2ac}, \frac{n}{ac}\right], \qquad B_{a,c} := I_{a,c} \cap \mathbb{P},$$

and define the target family

$$\mathcal{T} := \{\, acb : \ a < c, \ a, c \in \mathcal{P}_Y, \ b \in B_{a,c} \,\}.$$

For large n, every such b is greater than Y: indeed $b > n/(2Y^2) = \frac{1}{2}n^{1-2\delta} > Y$. Thus a, c, b are distinct primes and each target acb lies in $(n/2, n]$.

Proposition A.1 (Max-gain domination is false in rank three)**.** *The following finite scored rank-3 capture state has a unique max-gain Maker move, but Breaker can answer that move by deleting one uncaptured vertex and make the scaled potential decrease.*

Proof. Let the already-captured vertices be

$$C = \{z_{ij} : 1 \le i, j \le 3\},$$

and let u_1, u_2, u_3, v be uncaptured. There are no deleted or scored edges. The live edges are

$$f = \{u_1, u_2, u_3\}, \qquad w(f) = 9,$$

and, for $1 \le i, j \le 3$,

$$e_{ij} = \{v, u_i, z_{ij}\}, \qquad w(e_{ij}) = 4.$$

For a live edge e, write $\Phi(e) = 2^{|e \cap C|} w(e)$ and $Q = 8S + \sum \Phi(e)$.

Before Maker moves, $\Phi(f) = 9$ and $\Phi(e_{ij}) = 8$ for all i, j. If Maker plays f, then the score contribution of f changes by $8w(f) - \Phi(f) = 72 - 9 = 63$, and each of the nine edges e_{ij} doubles from 8 to 16. Hence

$$\Delta(f) = 63 + 9 \cdot 8 = 135.$$

After this move, the vertex v is still uncaptured, and deleting v removes all nine edges e_{ij}, now of potential 16 each. The Breaker loss is

$$9 \cdot 16 = 144 > 135.$$

Thus the full round decreases Q.

It remains only to check that f is the unique max-gain move. If Maker plays some e_{ij}, its own score gain is $8 \cdot 4 - 8 = 24$. The edge f doubles from 9 to 18, giving gain 9. The two other edges with the same u_i gain 24 each, and the six edges with the other two u-vertices gain 8 each. Therefore

$$\Delta(e_{ij}) = 24 + 9 + 2 \cdot 24 + 6 \cdot 8 = 129 < 135.$$

So the unique max-gain move is f, and it does not dominate the subsequent vertex deletion at v. □

Proposition A.2 (K_4-fiber refutation in the residual arithmetic game)**.** *For every fixed $0 < \delta < 1/4$ and all sufficiently large n, the residual slot hypergraph $G(n, \delta)$ contains a reachable pre-Maker state at which every legal Maker move admits a legal Breaker reply that strictly decreases the scaled potential Q. Hence the general safe-edge hypothesis is false even inside the arithmetic games $G(n, \delta)$.*

Proof. Here $G(n,\delta)$ is the scored rank-three slot hypergraph with edges

$$e_{a,c,b} := \{b, ab, cb\}, \qquad a < c, \quad a, c \in \mathcal{P}_Y, \quad b \in B_{a,c},$$

each of weight 1.

Choose the four fixed small primes $13, 17, 19, 23$. Their pairwise products are

$$13 \cdot 17 = 221, \quad 13 \cdot 19 = 247, \quad 13 \cdot 23 = 299, \quad 17 \cdot 19 = 323, \quad 17 \cdot 23 = 391, \quad 19 \cdot 23 = 437.$$

All six products lie between 221 and 437, and $2 \cdot 221 = 442 > 437$. For sufficiently large n, since $23 \le n^\delta = Y$, all four primes lie in $\mathcal{P}_Y$. By the prime number theorem, the interval

$$\left(\frac{n}{442}, \frac{n}{437}\right]$$

contains a prime q for all sufficiently large n. Also $q > Y$ for large n, since $\delta < 1/4$.

For every pair $a < c$ among $\{13, 17, 19, 23\}$, we have

$$q > \frac{n}{442} \ge \frac{n}{2ac} \qquad \text{and} \qquad q \le \frac{n}{437} \le \frac{n}{ac},$$

because $ac \ge 221$ and $ac \le 437$. Hence $q \in B_{a,c}$ for all six pairs, and the six edges

$$e_{a,c} := \{q, aq, cq\}$$

all occur in $G(n,\delta)$. They form a K_4-fiber over the common slot q.

We next reach a state in which only these six edges are live and no vertex in them has yet been captured. While any other edge is live, Maker uses the alternate-scoring move to score that exact target without adding slot vertices to C. Any edge outside this six-edge family has at least one slot outside

$$U := \{q, 13q, 17q, 19q, 23q\},$$

so Breaker deletes such an outside slot. This never harms the six desired edges. Repeating this finite operation leaves exactly the six K_4-fiber edges live, with $C = \varnothing$.

From that state, play two legal rounds. First Maker captures

$$e_{13,17} = \{q, 13q, 17q\},$$

so $q, 13q, 17q \in C$, and Breaker deletes the live edge $e_{13,19}$. Next Maker captures

$$e_{19,23} = \{q, 19q, 23q\},$$

so $q, 13q, 17q, 19q, 23q \in C$, and Breaker deletes the live edge $e_{13,23}$.

At the next pre-Maker state, the only remaining live edges from this fiber are

$$e_{17,19} = \{q, 17q, 19q\}, \qquad e_{17,23} = \{q, 17q, 23q\}.$$

Both have all three vertices in C, so $\Phi(e) = 2^3 \cdot 1 = 8$ for each. There are no other live edges by construction.

Maker has only two possible live edges to play. If Maker plays $e_{17,19}$, by ordinary capture or by alternate scoring, then the change in Q is

$$+8 \cdot 1 - 8 = 0,$$

because S increases by 1, the edge of potential 8 is removed, and no new vertex is added to C. The remaining live edge $e_{17,23}$ still contributes 8. Breaker then deletes $e_{17,23}$, decreasing Q by 8. Thus $Q_{\text{after}} = Q_{\text{before}} - 8$. The same argument holds if Maker instead plays $e_{17,23}$.

Therefore this reachable pre-Maker state has the claimed property: every legal Maker move admits a legal Breaker reply that strictly decreases Q. $\square$

Proposition A.3 (Conditional finite scaled capture)**.** *Consider a finite weighted incidence game of rank at most* 3 *with the scaled potential*

$$Q := 8S + \sum_{e \text{ live}} 2^{|e\cap C|} w(e),$$

where C is the current captured-vertex set and S is the total scored weight. In the graph variant, Breaker may delete one unclaimed live edge; in the scored hypergraph variant, Breaker may score one live edge.

The general safe-edge hypothesis stated below is provably false even for $G(n,\delta)$; see Proposition A.2. In this paper the finite-capture proposition is used only under the restricted hypothesis that the same property holds for states actually generated by the activation and residual strategies in Propositions A.6 and A.9. This restricted hypothesis remains open.

Assume the following restricted safe-edge hypothesis: at every pre-Maker state generated by the strategy under consideration with at least one positive-weight live edge, Maker has a legal live edge f such that, after Maker plays f, every legal Breaker reply leaves Q at least as large as it was before Maker's move.

Under this restricted hypothesis, if the game is continued until no positive-weight live edge remains, the final claimed or scored weight is at least one eighth of the initial total edge-weight:

$$S_{\text{fin}} \geq \frac{1}{8}\sum_{e} w(e).$$

Proof. Maker always chooses an edge supplied by the restricted safe-edge hypothesis. By definition of that hypothesis, Q is nondecreasing after every full Maker–Breaker round.

Initially $C = \varnothing$, $S = 0$, and all initial edges are live, so

$$Q_0 = \sum_{e} w(e).$$

At terminal time there is no positive-weight live edge, hence $Q_{\text{fin}} = 8S_{\text{fin}}$. Therefore

$$8S_{\text{fin}} = Q_{\text{fin}} \geq Q_0 = \sum_{e} w(e),$$

which is the claimed one-eighth bound.

For the graph variant, a Breaker edge-deletion reply is included in the restricted safe-edge hypothesis. For the scored hypergraph variant, a Breaker scored-edge reply is harmless whenever it occurs, since it changes Q by $8w(e) - 2^{|e\cap C|}w(e) \geq 0$ for rank at most 3; the only nontrivial part of the hypothesis is therefore the post-Maker vertex-deletion bound. □

Proposition A.4 (Initial target mass)**.** *For every fixed $0 < \delta < 1/4$,*

$$W_0 := \sum_{\substack{a<c \\ a,c\in\mathcal{P}_Y}} |B_{a,c}| \gg_\delta \frac{n(\log\log n)^2}{\log n}.$$

Proof. Since $ac \leq Y^2 = n^{2\delta}$, the interval endpoint $X_{a,c} := n/(ac)$ satisfies $X_{a,c} \geq n^{1-2\delta} \to \infty$. By the prime number theorem, uniformly for all such pairs and all sufficiently large n,

$$\pi(X_{a,c}) - \pi(X_{a,c}/2) \geq \kappa_\delta \frac{X_{a,c}}{\log n} = \kappa_\delta \frac{n}{ac\log n}$$

for some positive constant κ_δ. Therefore

$$W_0 \geq \kappa_\delta \frac{n}{\log n} \sum_{\substack{a<c \\ a,c\in\mathcal{P}_Y}} \frac{1}{ac}.$$

Mertens' theorem for primes gives

$$\sum_{p\in\mathcal{P}_Y}\frac{1}{p}=\log\log Y+O(1)=\log\log n+O_\delta(1),$$

and $\sum_p p^{-2}<\infty$. Hence

$$\sum_{a<c\in\mathcal{P}_Y}\frac{1}{ac}=\frac{1}{2}\left[\left(\sum_{p\in\mathcal{P}_Y}\frac{1}{p}\right)^2-\sum_{p\in\mathcal{P}_Y}\frac{1}{p^2}\right]=\left(\frac{1}{2}+o(1)\right)(\log\log n)^2.$$

Combining the last two displays proves the claim. □

Proposition A.5 (Residual divisibility embedding)**.** *Let $\mathcal{T}_*$ be any subfamily of targets $t=acb\in\mathcal{T}$ such that, for every $t=acb\in\mathcal{T}_*$, the numbers a, c, and ac are already unavailable as legal moves. Attach to t the three slot labels*

$$b,\qquad ab,\qquad cb$$

and let $e_t:=\{b,ab,cb\}$. Then the residual divisibility game on $\mathcal{T}_*$ is at least as favorable to Prolonger as the scored rank-3 hypergraph game on the hyperedges $\{e_t:t\in\mathcal{T}_*\}$:

1. *the only future moves that can make $t=acb$ illegal are b, ab, cb, and t itself;*
2. *playing b, ab, or cb deletes exactly the residual targets whose hyperedges contain that slot;*
3. *playing the exact target t kills no other target in $\mathcal{T}_*$, and is therefore a scored-edge move;*
4. *every live hyperedge corresponds to a legal actual move t.*

Proof. Fix $t=acb\in\mathcal{T}_*$. Because $t>n/2$, no proper multiple of t lies in $\{2,\dots,n\}$. Thus a future move can make t illegal only by being t itself or a proper divisor of t. The positive divisors are

$$1,\ a,\ c,\ ac,\ b,\ ab,\ cb,\ acb.$$

The divisor 1 is not a move, while a, c, and ac are unavailable by hypothesis. This proves the first assertion.

For slot incidence, suppose another target is $t'=a'c'b'$. Since $b,b'>Y\geq a,a',c,c'$, the divisibility $b\mid t'$ holds iff $b=b'$. Thus a move b kills exactly the targets in the b-fiber. Similarly, if $ab\mid a'c'b'$, the large prime factor forces $b=b'$, and after cancellation a must be a' or c'. This is exactly the condition that the slot ab belongs to $e_{t'}$. The proof for cb is identical.

If two targets t,u in $\mathcal{T}_*$ are comparable, say $t\mid u$, then u is a multiple of t lying in $(n/2,n]$. The next positive multiple of t after t is $2t>n$, so $u=t$. Hence distinct targets are pairwise incomparable, and playing an exact target scores only that target.

Finally, if e_t is live in the slot hypergraph, then none of b, ab, cb, or t has been played. The only other proper divisors of t are $1,a,c,ac$, and these are either not moves or already unavailable. There is no proper multiple of t on the board. Therefore no played number is comparable with t, so the actual move t is legal. This proves the comparison. □

Proposition A.6 (Activation-stage potential)**.** *During activation, let each unclaimed pair edge $e=(a,c)$ carry the current token set $B_e(t)$ of primes $b\in B_{a,c}$ for which the target acb has not yet been killed. Write $w_t(e):=|B_e(t)|$.*

Assume the restricted safe-edge hypothesis of Proposition A.3 for the activation graph states generated below. Prolonger plays as follows. In each activation round, among pair edges with positive current token weight, he chooses a safe edge supplied by that hypothesis; if $e=(a,c)$ is chosen, he plays one live target acb with $b\in B_e(t)$. This scores one actual move, claims the pair edge e, captures the vertices a and c, and leaves the remaining live tokens on that secured pair for the residual phase.

Let S_t be the number of activation targets already played by Prolonger, and define

$$Q_t:=S_t+\sum_e\phi_t(e),$$

where

$$\phi_t(e) = \begin{cases} \frac{1}{8}w_t(e), & e \text{ unclaimed and neither endpoint captured}, \\ \frac{1}{4}w_t(e), & e \text{ unclaimed and one endpoint captured}, \\ \frac{1}{2}w_t(e), & e \text{ unclaimed and two endpoints captured}, \\ w_t(e), & e \text{ claimed}, \\ 0, & e \text{ deleted}. \end{cases}$$

Let E be the total number of target tokens deleted during activation by Shortener moves not modeled as graph vertex deletions a or graph edge deletions ac. At the end of activation,

$$Q_{\text{end}} \geq \frac{W_0}{8} - E.$$

Equivalently,

$$S_{\text{end}} + \sum_{e \text{ secured}} w_{\text{end}}(e) \geq \frac{W_0}{8} - E.$$

Proof. First ignore off-model token deletions. Conditional on the restricted safe-edge hypothesis, the activation process is exactly the rank-2 graph game of Proposition A.3, with one harmless bookkeeping change: when Prolonger claims an edge e, he consumes one token from e and adds it immediately to the score S_t. If the coefficient of e before the move is $\gamma \in \{1/8, 1/4, 1/2\}$, the contribution of e changes from $\gamma w_t(e)$ to

$$1 + (w_t(e) - 1) = w_t(e),$$

exactly as if the edge had become claimed with full weight. All other incident edges change as in the graph game. Thus the scaled potential argument gives $Q_{\text{end}} \geq W_0/8$ when every harmful Shortener move is a graph deletion.

Now restore all actual Shortener moves. A token acb is deleted precisely when Shortener plays one of its still-available harmful divisors or the target itself. Deleting one live token can reduce Q_t by at most 1, since every coefficient appearing in the definition of $\phi_t(e)$ is at most 1. Therefore all off-model deletions together reduce the lower bound by at most E, giving

$$Q_{\text{end}} \geq W_0/8 - E.$$

At the end of activation, a positive-weight edge is either secured or deleted, so Q_{end} is exactly the activation score plus the residual live target weight on secured pairs. □

Proposition A.7 (Activation deletion budget)**.** *For fixed $0 < \delta < 1/4$, the off-model deletion count from Proposition A.6 satisfies*

$$E \ll \frac{Y^4}{\log^4 Y} = o\left(\frac{n(\log\log n)^2}{\log n}\right).$$

Proof. There is at most one activation round per small-prime pair, so the number of rounds is

$$R \leq \binom{|\mathcal{P}_Y|}{2} \ll \frac{Y^2}{\log^2 Y}.$$

We bound the number of live tokens an off-model Shortener move can delete.

If Shortener plays a large prime b, then for each pair (a, c) there is at most one token acb using that b. Hence the move deletes at most $O(Y^2/\log^2 Y)$ tokens. If Shortener plays a lateral semiprime pb with $p \leq Y$ and $b > Y$ prime, the killed targets have the form pcb with the third small prime varying, so there are at most $O(Y/\log Y)$ of them. If Shortener plays an exact target acb, exactly one token is deleted.

The worst of these bounds is the large-prime bound. Across R activation rounds,

$$E \ll R\frac{Y^2}{\log^2 Y} \ll \frac{Y^4}{\log^4 Y}.$$

Since $Y = n^\delta$,

$$\frac{Y^4/\log^4 Y}{n(\log\log n)^2/\log n} \ll_\delta \frac{n^{4\delta-1}}{(\log n)^3(\log\log n)^2} \to 0$$

because $4\delta - 1 < 0$. □

Proposition A.8 (Residual mass on secured pairs)**.** *At the end of activation, the secured pairs carry residual live target weight*

$$M := \sum_{e \text{ secured}} w_{\text{end}}(e) \gg_\delta \frac{n(\log\log n)^2}{\log n}.$$

Proof. By Proposition A.6,

$$S_{\text{end}} + M \geq \frac{W_0}{8} - E.$$

The activation score is at most the number of small-prime pairs:

$$S_{\text{end}} \leq R \ll \frac{Y^2}{\log^2 Y} = o\left(\frac{n(\log\log n)^2}{\log n}\right),$$

because $2\delta < 1$. The deletion budget E is negligible by Proposition A.7, while W_0 has the required lower bound by Proposition A.4. Subtracting the two negligible terms gives the claim. □

Proposition A.9 (Conditional T2 lower bound)**.** *Assume the restricted safe-edge hypothesis of Proposition A.3 for the activation graph states and for the residual slot-hypergraph states generated by the construction above. For every fixed $0 < \delta < 1/4$ there is $c_\delta > 0$ such that*

$$L(n) \geq c_\delta \frac{n(\log\log n)^2}{\log n}$$

for all sufficiently large n.

Proof. After activation, restrict to the residual live targets lying over secured pairs. Their total weight is M, as in Proposition A.8. For each secured pair (a, c), the activation move made a, c, and ac unavailable. Therefore Proposition A.5 compares the remaining divisibility game to a scored rank-3 slot hypergraph whose total edge weight is M.

Under the residual restricted safe-edge hypothesis, Proposition A.3 gives a strategy that forces scored residual target weight at least $M/8$ in this hypergraph model. The comparison proposition says that every such scored hypergraph move is a genuine actual game move, and that the actual game gives Prolonger no fewer legal options than the model. Hence the number of residual moves after activation is at least $M/8$. Since $M \gg_\delta n(\log\log n)^2/\log n$, the same lower bound holds for the total game length, conditional on the restricted safe-edge hypothesis.

Choosing any fixed $\delta < 1/4$, for instance $\delta = 1/8$, gives an absolute constant $c > 0$ and proves Theorem 4.6. □

A.2. **Restricted carrier classes.** The next two propositions record unconditional upper bounds for two restricted families of Prolonger carrier moves. They are not used in the main upper bound, but they clarify which carrier configurations are not responsible for the remaining difficulty.

Proposition A.10 (Disjoint small-prime carriers)**.** *Fix $1/3 < \alpha < 1/2$ and put $y = n^\alpha$. Suppose that every composite Prolonger move has all prime factors at most y, and that the prime supports*

of distinct composite Prolonger moves are pairwise disjoint. Against this restricted class, Shortener has a strategy forcing

$$L(n) = O_\alpha\left(\frac{n}{\log n}\right).$$

Proof. Let B be the set of primes which occur in composite Prolonger moves. By the disjoint-support hypothesis, each $p \in B$ belongs to a unique composite carrier $C(p)$.

Shortener uses the following priority list, always skipping moves which have already become illegal.

First, she plays every legal prime. This resolves every prime not belonging to a composite Prolonger carrier, and also every prime which Prolonger has not already made illegal by placing it inside a carrier.

Second, for every $p \in B$ whose carrier has at least two distinct prime factors, write $e(p) = v_p(C(p))$. If $p^{e(p)+1} \le n$, Shortener plays $p^{e(p)+1}$. This move is legal when it is attempted: it is not comparable with $C(p)$, because it has a larger p-adic exponent while $C(p)$ contains another prime factor; it is not comparable with any other composite carrier by disjointness of supports; and previous priority moves either contain different prime supports or have smaller rank.

Third, for every pair $p, q \in B$ belonging to different carriers, Shortener plays the legal squarefree semiprime pq when possible. Such a move is incomparable with every carrier, since no carrier contains both p and q, and neither p nor q alone is available as a legal prime at this stage.

The number of moves in the three phases is

$$O(\pi(n)) + \pi(y) + \pi(y)^2 = O_\alpha\left(\frac{n}{\log n}\right),$$

because $2\alpha < 1$.

We now show that after these phases every future legal move has no prime factor at most y. Let x be a legal move divisible by a prime $p \le y$.

If $p \notin B$, then phase one either played p or p was already played by Prolonger as a prime; in either case any multiple of p is illegal. Thus $p \in B$.

If x contains primes from two distinct carriers, then one of the phase-three semiprimes pq divides x, unless it was already illegal because an earlier played move dividing pq also divides x. Either way x is illegal.

Hence all prime factors of x which are at most y lie in a single carrier support. If the carrier is a pure prime power p^e, then any x divisible by p and supported only on this carrier is another power of p, and is comparable with p^e. If the carrier has at least two distinct prime factors, then either x divides the carrier, in which case x is illegal, or for some p it has exponent exceeding $v_p(C(p))$. In the latter case $p^{e(p)+1} \mid x$, so the phase-two move makes x illegal.

Thus all primes $\le y$ are resolved. Any remaining legal integer has all prime factors greater than y. Since $y^3 > n$, it has at most two prime factors. The number of integers at most n with one or two prime factors, all exceeding $y = n^\alpha$, is $O_\alpha(n/\log n)$ by the prime number theorem and the standard semiprime estimate

$$\sum_{p>y} \pi(n/p) \ll_\alpha \frac{n}{\log n} \sum_{p>y} \frac{1}{p} = O_\alpha\left(\frac{n}{\log n}\right),$$

where the sum is over $p \le n/y$. Adding the phase moves proves the claim. □

Proposition A.11 (Squarefree rank-three small-prime carriers)**.** *Fix $1/3 < \alpha < 1/2$ and put $y = n^\alpha$. Suppose that every composite Prolonger move is squarefree, has all prime factors at most y, and has support size at most 3. Against this restricted class, Shortener has a strategy forcing*

$$L(n) = O_\alpha\left(\frac{n}{\log n}\right).$$

Proof. Let B be the set of primes which occur in Prolonger's composite carriers. Shortener again uses a fixed priority list, skipping illegal moves.

First she plays every legal prime. Second she plays p^2 for every $p \in B$ when legal. Third she plays every legal squarefree semiprime pq with $p, q \in B$. Fourth she plays every legal squarefree triple pqr with $p, q, r \in B$.

The first phase has $O(n/\log n)$ moves, the second has at most $\pi(y)$ moves, and the third has at most $\pi(y)^2 = o(n/\log n)$ moves. It remains to bound the number of triples

$$T_\alpha(n) := \#\{p < q < r \le y : \ pqr \le n\}.$$

Split according to whether $pq \le n^{1-\alpha}$.

If $pq \le n^{1-\alpha}$, then there are $O(n^{1-\alpha} \log\log n/\log n)$ possible pairs $p < q$, and for each pair at most $\pi(y) = O(n^\alpha/\log n)$ choices of r. This contributes $o(n/\log n)$.

If $pq > n^{1-\alpha}$, then $q \le y = n^\alpha$ implies $p > n^{1-2\alpha}$. In this range,

$$\#\{r : \ q < r \le y, \ pqr \le n\} \ll_\alpha \frac{n}{pq \log n}.$$

Therefore the contribution is at most

$$\ll_\alpha \frac{n}{\log n} \sum_{n^{1-2\alpha} < p < q \le n^\alpha} \frac{1}{pq} = O_\alpha\left(\frac{n}{\log n}\right),$$

because the reciprocal prime sum over a fixed-power interval is $O_\alpha(1)$. Thus $T_\alpha(n) = O_\alpha(n/\log n)$.

We prove that the priority list resolves all primes at most y. After the prime and square phases, any remaining legal move supported on B must be squarefree. If it has support size 1, it is comparable with a played or blocked prime. If it has support size 2, then the corresponding semiprime was either played in phase three or was already illegal because an earlier played divisor blocks it. If it has support size 3, the same statement holds for phase four.

If a remaining legal move x had support size at least 4 inside B, choose three primes p, q, r from its support. The triple pqr was addressed in phase four. If it was played, then $pqr \mid x$ and x is illegal. If it was not legal when considered, then some earlier played move comparable with pqr must divide pqr; since all earlier non-prime moves in the priority list have support size at most 3 and are squarefree except the squares p^2, that played divisor also divides x. Again x is illegal. Thus no remaining legal move contains a prime at most y.

As in the proof of Proposition A.10, once all primes at most y are resolved, the residual legal set has size $O_\alpha(n/\log n)$. Combining this residual bound with the phase counts proves the proposition. □

Appendix B. Numerical Computations

This appendix records the numerical computations accompanying the main Bonferroni–4 constant. The computations certify the quoted constants and stress-test the finite combinatorial steps, but they are not substitutes for the analytic proof in Section 7.

B.1. **The Bonferroni–4 constant.**

The main theorem uses

$$\frac{\mathcal{W}_4}{2} = 0.1897112 < 0.19.$$

A direct grid computation of the integrals

$$J_r = \frac{1}{r!} \int_{\substack{u_1 + \cdots + u_r \le 1 \\ 0 < u_i \le 1}} \prod_{i=1}^{r} \rho(u_i) \, du_1 \cdots du_r$$

gives the convergence displayed in Table 1. The theorem only needs a strict margin below 0.19, and the finest-grid value provides a comfortable margin.

Grid	J_1	J_2	J_3	J_4	$\mathcal{W}_4$	$\mathcal{W}_4/2$
2^{13}	0.78844185	0.18676003	0.02008041	0.00122154	0.37945930	0.18972965
2^{14}	0.78849100	0.18679383	0.02008737	0.00122220	0.37943767	0.18971883
2^{15}	0.78851725	0.18681153	0.02009097	0.00122255	0.37942586	0.18971293
2^{16}	0.78852127	0.18681405	0.02009149	0.00122260	0.37942388	0.18971194
2^{17}	0.78852704	0.18681777	0.02009220	0.00122266	0.37942118	0.18971059

TABLE 1. Grid convergence for the Bonferroni–4 truncation.

B.2. **Finite sanity checks.** Four finite checks were used to guard against implementation or formulation errors in the proof.

(1) *Odd-part injection.* Exhaustive verification at $n = 14$ over all 732 antichains found no collision under the odd-part map.
(2) *Monotone replacement.* Exact inclusion–exclusion counts at $n = 10^4, 10^5, 10^6$ for test prime pairs (q_j) and (p_j) with $q_j \le p_j$ confirmed $N(q_1, \dots, q_K) \le N(p_1, \dots, p_K)$.
(3) *Squarefree divisor counts.* For $r \le 4$, the relevant squarefree r-fold divisor counts match the scale $O_r(n(\log\log n)^{r-1}/\log n)$ used in the Bonferroni floor-error bound.
(4) *Controlled prime rounding.* Synthetic monotone comparison prefixes were rounded upward to increasing odd-prime prefixes at $n = 10^4, 10^5, 10^6, 10^8$; the resulting changes in the moment terms were small and consistent with the prime-rounding bridge.

Check	Recorded outcome
Odd-part injection	`(True, None, None, 732)` at $n = 14$
Monotone replacement	Verified at $n = 10^4, 10^5, 10^6$ for listed test prefixes
M_r scaling	Ratified for $r = 1, 2, 3, 4$ up to $n = 10^6$
Prime-rounding bridge	Tested on synthetic prefixes up to $n = 10^8$

TABLE 2. Finite checks for the combinatorial components of the upper-bound proof.

B.3. **Interpretation.** The numerical artifacts support three claims: the constant $\mathcal{W}_4/2 < 0.19$ is reproducible; the finite steps in Theorems 7.4 and 7.7 behave as expected in representative regimes; and the sharper limiting value near 0.18969 is numerically stable. The theorem in this paper is the finite Bonferroni–4 bound below 0.19, not a claim that the full alternating series has been analytically justified.

APPENDIX C. FORMALIZATION ARTIFACT MAP

This appendix records the status of the Lean artifacts corresponding to the paper's principal results. The table distinguishes between fully formalized finite or structural cores and prose components which use standard analytic number theory or conditional game-theoretic hypotheses.

The table should be read conservatively. A zero-sorry finite core is not being advertised as an end-to-end verification of every analytic asymptotic in the same theorem unless the table says so explicitly. Conversely, when the remaining obligations are standard analytic-number-theory inputs, the novelty of the combinatorial argument is already isolated in the formal artifact.

ARTIFACT AVAILABILITY

The public artifact repository for this paper is [Bud26]. It contains the Lean formalizations, Aristotle submission files and outputs, numerical scripts, and supporting research notes used in the preparation of the present manuscript. Repository-relative paths quoted in Appendices B and C are relative to that artifact root.

Result	Lean status	Remaining prose or assumptions
Shield Reduction, Theorem 3.1	Zero-sorry Lean verification.	None beyond the finite definitions of the divisibility poset.
Exact 5/24 cover, Theorem 5.1	The structural cover and packing identities are zero-sorry Lean-verified.	The asymptotic floor-sum count $5n/24 + O(1)$ is an elementary prose calculation.
13/36 upper bound, Theorem 6.2	The odd-part compression and Bonferroni core are zero-sorry Lean-verified.	The paper-level exposition uses standard prime-counting estimates for the prime-prefix setup.
Polynomial shield-weight lower bound, Theorem 4.1	The combinatorial core, including the prime-antichain construction, log-budget argument, and exchange principle, has a Lean artifact.	Remaining sorries are Mertens' theorem, a Chebyshev-type prime-counting bound, a Chebyshev-ϑ asymptotic, and the final assembly using those analytic inputs.
5/16 upper bound, Theorem 6.3	The compression, sieve, and algebraic optimization core has a Lean artifact.	One remaining bundled placeholder covers the Chebyshev prime-prefix estimate and the game-tree wrapper realizing the prescribed prefix.
Conditional T2 lower bound, Theorem 4.6	The finite graph-potential core, scored hypergraph core, and local arithmetic embedding core are Lean-verified.	The restricted safe-edge hypothesis is conditional, and the asymptotic activation wrapper remains in prose.
Main $0.19n$ upper bound, Theorem 7.8	The endgame reduction from sufficiently close first-four moment values to the strict inequality $L(n) < 0.19n$ is zero-sorry Lean-verified.	The local density theorem, monotone envelope and inversion, and prime-rounding bridge are the prose moment-convergence inputs proved in Section 7.

Table 3. Formalization status of the principal results.

Acknowledgments

The author thanks Liam Price, Abhimanyu Adenwalla, StijnC, natso26, Xiao_Hu, and Desmond Weisenberg for the forum refinement chain leading from the first linear bound to the previous public record, and Thomas Bloom for maintaining `erdosproblems.com`. The author also wishes to record substantial gratitude for the research assistance provided by OpenAI's GPT-5.4 Pro, Anthropic's Claude Opus 4.7, and Google's Gemini 3.1 DeepThink, coordinated through a custom multi-round research harness built by the author. GPT-5.4 Pro produced the majority of the exploratory mathematical output used in this project, with Claude and Gemini providing extensive auditing, synthesis, and drafting support. A separate methodological paper and public-facing writeup will describe that workflow in detail.

Declaration of Generative AI and AI-assisted Technologies in the Research and Writing Process

During the preparation of this work, the mathematical content was produced primarily by generative AI systems—OpenAI GPT-5.4 Pro, Anthropic Claude Opus 4.7, and Google Gemini 3.1 DeepThink—coordinated through a custom multi-round research harness designed and operated by the author. GPT-5.4 Pro generated the majority of the novel mathematical arguments, including the shield reduction, the polynomial shield barrier, the Round 15 piecewise-density analysis, and the barrier taxonomy. Claude Opus 4.7 and Gemini 3.1 DeepThink provided adversarial auditing, cross-model verification, synthesis across rounds, and drafting support. The author's contributions were the harness design, problem selection and scoping, round-by-round direction and adversarial

prompting, selection and integration of outputs into a coherent manuscript, Lean/Aristotle formalization work, numerical verification, and final writing. The author takes full responsibility for the correctness of the claims, the validity of the proofs as presented, the accuracy of the citations, and the integrity of the formal artifacts. A description of the harness and workflow is available at [URL].

References

[BHW16] Csaba Biró, Paul Horn, and D. Jacob Wildstrom, *An upper bound on the extremal version of hajnal's triangle-free game*, Discrete Applied Mathematics **198** (2016), 20–28.

[Blo26] Thomas F. Bloom, *Erdős problem #872*, https://www.erdosproblems.com/872, 2026, Accessed 2026-04-20.

[Bud26] Om Buddhdev, *erdos-harness: Artifacts and formalization files for erdős problem #872*, https://github.com/xa8zz/erdos-harness, 2026, Public artifact repository for this paper; accessed 2026-04-20.

[CE90] Peter J. Cameron and Paul Erdős, *On the number of sets of integers with various properties*, Number Theory (Banff, AB, 1988), de Gruyter, Berlin, 1990, pp. 61–79.

[Erd92] Paul Erdős, *Some of my forgotten problems in number theory*, Hardy-Ramanujan Journal **15** (1992), 34–50.

[Erd26] Erdős Problems Forum contributors, *Discussion thread for erdős problem #872*, https://www.erdosproblems.com/forum/thread/872, 2026, Includes refinements by Adenwalla, StijnC, natso26, Xiao_Hu, Desmond Weisenberg, and others, leading to the bound $(419/1008 + o(1))n$; accessed 2026-04-20.

[FS92] Zoltán Füredi and Ákos Seress, *On hajnal's triangle-free game*, Graphs and Combinatorics **8** (1992), no. 1, 75–79.

[Kuc24] Petr Kucheriaviy, *Erdős inequality for primitive sets*, 2024.

[Lic23] Jared Duker Lichtman, *A proof of the erdős primitive set conjecture*, Forum of Mathematics, Pi **11** (2023), e18.

[MP10] Greg Martin and Carl Pomerance, *Primitive sets with large counting functions*, 2010.

[PG26] Liam Price and GPT-5.2 Pro, *First linear upper bound for erdős problem #872*, https://www.erdosproblems.com/forum/thread/872, 2026, Forum contribution on the Erdős Problems discussion thread for problem #872, February 2026.

Independent Researcher
Email address: sensho@sensho.xyz