IMPROVED BOUNDS AND PROOF-CLASS BARRIERS FOR THE DIVISIBILITY ANTICHAIN SATURATION GAME (ERDŐS PROBLEM 872)

OM BUDDHDEV

Abstract. We study the saturation game on the divisibility poset of $\{2, 3, \ldots, n\}$. Two players alternately choose integers subject to the condition that the chosen set remains primitive; Prolonger moves first and maximizes the final number of moves, while Shortener minimizes it. We prove

$$L(n) \le \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n = (0.1897112 + o(1))n < 0.19n,$$

where $\mathcal{W}_4$ is an explicit fourth-order Bonferroni constant coming from the local logarithmic density of Shortener's legal-prime prefix. We also prove a polynomial lower bound for the weighted shield functional, an exact $5n/24 + O(1)$ first-hit cover theorem for the lower half, and the intermediate upper bounds $13n/36+o(n)$ and $5n/16+o(n)$. A conditional lower bound of order $n(\log\log n)^2/\log n$ is included under a finite safe-edge hypothesis which is stated explicitly. The paper also records three proof-class barriers for local relaxations, Johnson-shadow separators, and separator-only carrier strategies.

1. Introduction

Let $n \ge 2$. The divisibility antichain saturation game is played on

$$\{2, 3, \ldots, n\}.$$

Two players alternately choose previously unchosen integers, and the chosen set must remain an antichain under divisibility. Thus no chosen integer may divide another chosen integer. The game stops when the current antichain is maximal. Prolonger moves first and tries to maximize the final number of chosen integers; Shortener tries to minimize it. We denote the value of the game by $L(n)$.

This game is one of the finite saturation games arising from Erdős's questions on primitive sets [Erd92]; saturation games of this kind are part of a broader extremal-game tradition [FS92, BHW16]. A primitive set is precisely an antichain in the divisibility poset, and primitive sets have a long parallel history in multiplicative number theory [CE90, Lic23, MP10, Kuc24]. The upper half

$$U_n := \{m : \ n/2 < m \le n\}$$

is the canonical maximum antichain, so the trivial scale is linear. The central question is whether optimal play forces a positive fraction of this linear scale, or whether Shortener can force $o(n)$ moves.

Our main result is a new linear upper bound with an explicit constant below 0.19. Let

$$\rho(u) := \frac{1}{(\lfloor 1/u \rfloor + 1)u} \qquad (0 < u \le 1),$$

and define

$$J_r := \frac{1}{r!} \int_{\substack{u_1+\cdots+u_r \le 1 \\ 0<u_i\le 1}} \prod_{i=1}^{r} \rho(u_i)\, du_1 \cdots du_r \qquad (r \ge 1).$$

Date: April 21, 2026.

2020 *Mathematics Subject Classification.* Primary 91A46; Secondary 05D05, 11N25.

Key words and phrases. Saturation game, primitive set, divisibility poset, antichain, sieve methods, formal verification, Erdős problem 872.

Set

$$\mathcal{W}_4 := 1 - J_1 + J_2 - J_3 + J_4.$$

The numerical value is

$$\frac{\mathcal{W}_4}{2} = 0.1897112\ldots < 0.19.$$

Theorem 1.1 (Main upper bound)**.** *Shortener has a strategy forcing*

$$L(n) \le \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n < 0.19n.$$

The proof of Theorem 1.1 is given in Section 7. Shortener plays a long prefix of the smallest legal odd primes. The local law governing which primes can be kept legal has density $\rho(u)$ on the logarithmic scale $p = n^u$. We build a monotone envelope for this density, invert it to obtain a comparison sequence, round the comparison sequence to genuine primes without losing the first four factorial moments, and then apply a fourth-order Bonferroni inequality to the odd-part compression of the final antichain.

The lower-bound side of the paper begins with a weighted obstruction theorem for the standard shield-reduction approach. For $x \le n/2$, let

$$M_n(x) := \{u \in U_n : \ x \mid u\}, \qquad w_n(x) := |M_n(x)| - 1.$$

For $P \subseteq U_n$, put

$$\mathcal{L}(P) := \{x \le n/2 : \ x \nmid u \text{ for every } u \in P\},$$

and let

$$\beta_n(P) := \max\left\{\sum_{x \in B} w_n(x) : B \subseteq \mathcal{L}(P) \text{ is a divisibility antichain}\right\}.$$

Theorem 1.2 (Polynomial shield-weight lower bound)**.** *Fix $0 < \alpha < 1$. Uniformly for every $P \subseteq U_n$ with $|P| \le n^\alpha$,*

$$\beta_n(P) \ge \left(\frac{1}{2}\log\frac{1}{\alpha} - o(1)\right) n.$$

Equivalently, if

$$k_c^\star(n) := \min\{|P| : \ P \subseteq U_n, \ \beta_n(P) \le cn\},$$

then for every fixed $c > 0$,

$$k_c^\star(n) \ge n^{e^{-2c} - o(1)}.$$

The next theorem explains why this weighted quantity is the right one for shield-prefix arguments.

Theorem 1.3 (Shield reduction framework)**.** *Let A be any terminal antichain and let $P \subseteq A \cap U_n$. Then*

$$|A| \ge |U_n| - \beta_n(P).$$

Consequently, any proof which uses a prefix $P \subseteq U_n$ and this reduction to force $|A| \ge (1/2 - c)n + o(n)$ must have

$$|P| \ge n^{e^{-2c} - o(1)}.$$

The proof of the reduction is elementary and is given in Section 3; Theorem 1.2 is proved in Section 4. These results do not give a game lower bound by themselves. They show instead that a short upper-half shield prefix cannot certify a linear lower bound through the weighted shadow method.

We also determine exactly the size of the first-hit cover of the lower half by upper-half multiples. Define

$$\tau(n) := \min\{|C| : \ C \subseteq U_n, \ \forall x \le n/2 \ \exists u \in C \text{ with } x \mid u\}.$$

Theorem 1.4 (Exact first-hit cover)**.**

$$\tau(n) = \frac{5}{24}n + O(1).$$

More precisely, the cover

$$H_n = \{u \in U_n : \ u \equiv 2 \pmod 4\} \cup \{u \in U_n : \ u > 2n/3, \ u \equiv 0 \pmod 4\}$$

has size $5n/24 + O(1)$*, and there is a matching packing of lower-half witnesses of the same size up to* $O(1)$*.*

Before the fourth-order argument, the same prime-prefix method gives two simpler upper bounds.

Theorem 1.5 (Intermediate upper bounds)**.** *Shortener has explicit strategies forcing*

$$L(n) \le \left(\frac{13}{36} + o(1)\right) n$$

and

$$L(n) \le \left(\frac{5}{16} + o(1)\right) n.$$

Finally, we include a conditional lower-bound theorem, not as a headline result but as a useful finite-combinatorial target. Under the safe-edge hypothesis stated in Section 4, the residual slot construction gives

$$L(n) \ge c\,\frac{n(\log\log n)^2}{\log n}$$

for some absolute $c > 0$. The finite graph, hypergraph, and arithmetic embedding cores of this theorem are formally verified; the safe-edge hypothesis and the asymptotic activation wrapper remain in prose and are stated explicitly rather than suppressed.

Several parts of the paper are accompanied by Lean formalizations. The Shield Reduction theorem, the structural identities underlying the exact 5/24 cover, and the 13/36 upper-bound core have zero-sorry Lean verifications. Theorem A and the 5/16 upper bound have Lean artifacts whose remaining sorries are standard analytic-number-theory inputs or their assembly. The finite graph/hypergraph/embedding cores of the conditional T2 theorem are Lean-verified, while the safe-edge hypothesis and asymptotic wrapper remain prose. For the $0.19n$ theorem, the endgame reduction from moment convergence to the strict inequality is zero-sorry Lean-verified; the moment-convergence hypotheses are the prose content of Section 7. A full status table appears in Appendix C.

The paper is organized as follows. Section 2 fixes notation. Section 3 proves the shield reduction. Section 4 proves Theorem A and states the conditional T2 theorem. Section 5 proves the 5/24 cover theorem. Section 6 proves the 13/36 and 5/16 upper bounds. Section 7 proves the main $0.19n$ bound. Section 8 records three obstructions to extending the present methods. Deferred proofs, numerical details, and formalization metadata appear in the appendices.

2. Notation and Setup

All divisibility statements in this paper are inside the finite board

$$V_n := \{2, 3, \ldots, n\}.$$

A set $A \subseteq V_n$ is *primitive* if no two distinct elements of A are comparable under divisibility. Equivalently, A is an antichain in the divisibility poset. A primitive set is *maximal* if no element of $V_n \setminus A$ can be added while preserving primitiveness.

The game value $L(n)$ is the final size of the primitive set under optimal play when Prolonger moves first and maximizes the final size, while Shortener minimizes it.

We use the lower–upper split

$$L_n := \{m \in \mathbb{Z} : \ 2 \le m \le \lfloor n/2 \rfloor\}, \qquad U_n := \{m \in \mathbb{Z} : \ n/2 < m \le n\}.$$

The upper half U_n is an antichain, since two distinct elements of U_n cannot divide each other.

For $x \in L_n$, the upper shadow and shadow weight are

$$M_n(x) := \{u \in U_n : \ x \mid u\}, \qquad w_n(x) := |M_n(x)| - 1.$$

The subtraction by 1 is convenient because a lower-half move x adds one move but may remove $|M_n(x)|$ possible upper-half moves from a terminal position.

For $P \subseteq U_n$, define

$$\mathcal{L}_n(P) := \{x \in L_n : \ x \nmid u \text{ for every } u \in P\},$$

and

$$\beta_n(P) := \max\left\{\sum_{x \in B} w_n(x) : B \subseteq \mathcal{L}_n(P) \text{ is a divisibility antichain}\right\}.$$

The maximum is over a finite set, and is understood to be 0 when $\mathcal{L}_n(P)$ is empty.

We use standard arithmetic notation. The functions $\pi(x)$ and $\vartheta(x)$ denote the prime-counting function and Chebyshev's function, respectively. For an integer m, $P^-(m)$ and $P^+(m)$ denote the least and greatest prime factors when $m > 1$, $\Omega(m)$ denotes the number of prime factors counted with multiplicity, and $\omega(m)$ denotes the number of distinct prime factors. The symbol $\mathbb{P}$ denotes the set of primes. As usual, $O(\cdot)$, $o(\cdot)$, $\ll$, and $\gg$ are interpreted as $n \to \infty$ unless a different parameter is explicitly specified; subscripts indicate dependence of the implied constants.

3. The Shield Reduction

The shield reduction turns terminal game positions into a weighted lower-half antichain problem.

Theorem 3.1 (Shield reduction)**.** *Let $A \subseteq V_n$ be a maximal primitive set and let $P \subseteq A \cap U_n$. Then*

$$|A| \ge |U_n| - \beta_n(P).$$

Proof. Put

$$B := A \cap L_n.$$

Then B is a divisibility antichain. Since $P \subseteq A$ and A is primitive, no $x \in B$ divides any element of P. Hence

$$B \subseteq \mathcal{L}_n(P).$$

We next determine $A \cap U_n$. We claim that

$$A \cap U_n = U_n \setminus \bigcup_{x \in B} M_n(x). \tag{3.1}$$

The inclusion "$\subseteq$" follows from primitiveness: if $u \in A \cap U_n$ and $x \in B$, then $x \nmid u$.

For the reverse inclusion, let $u \in U_n \setminus \bigcup_{x \in B} M_n(x)$. Suppose $u \notin A$. Since A is maximal, some $a \in A$ is comparable with u. If $u \mid a$, then $a \in U_n$ and $a < 2u$, so $a = u$, contradicting $u \notin A$. If $a \mid u$ and $a \in U_n$, again $a = u$. Thus $a \in L_n$, so $a \in B$, and $u \in M_n(a)$, contrary to the choice of u. This proves (3.1).

Using (3.1) and the union bound,

$$\begin{aligned}|A| &= |B| + |A \cap U_n| \\ &\geq |B| + |U_n| - \sum_{x \in B} |M_n(x)| \\ &= |U_n| - \sum_{x \in B} (|M_n(x)| - 1) \\ &= |U_n| - \sum_{x \in B} w_n(x).\end{aligned}$$

Since $B \subseteq \mathcal{L}_n(P)$ is an antichain, the final sum is at most $\beta_n(P)$. Therefore

$$|A| \geq |U_n| - \beta_n(P).$$

□

Corollary 3.2 (Shield-prefix obstruction)**.** *Fix $c > 0$. If a shield-prefix argument using Theorem 3.1 forces*

$$|A| \geq |U_n| - cn + o(n)$$

through a set $P \subseteq U_n$, then necessarily

$$|P| \geq n^{e^{-2c} - o(1)}.$$

Proof. Such an argument requires $\beta_n(P) \leq cn + o(n)$. Applying Theorem 4.1 with $|P| \leq n^\alpha$ gives

$$\frac{1}{2} \log \frac{1}{\alpha} \leq c + o(1),$$

and hence $\alpha \geq e^{-2c} - o(1)$. □

4. Lower Bounds and Conditional Slot Construction

This section proves the polynomial shield-weight obstruction and states the conditional T2 lower bound. The conditional result is deferred to Appendix A.

4.1. The polynomial shield-weight lower bound.

Theorem 4.1 (Polynomial shield-weight lower bound)**.** *Fix $0 < \alpha < 1$. Uniformly over all $P \subseteq U_n$ with $|P| \leq n^\alpha$,*

$$\beta_n(P) \geq \left(\frac{1}{2} \log \frac{1}{\alpha} - o(1) \right) n.$$

Proof. Fix δ with $\alpha < \delta < 1$. Let

$$Q_\delta(P) := \{p \leq n^\delta : \ p \text{ prime and } p \nmid u \text{ for every } u \in P\}.$$

For all sufficiently large n, every $p \leq n^\delta$ lies in L_n. The set $Q_\delta(P)$ is an antichain and is contained in $\mathcal{L}_n(P)$, so

$$\beta_n(P) \geq \sum_{p \in Q_\delta(P)} w_n(p). \tag{4.1}$$

For $p \leq n^\delta$,

$$w_n(p) = \left\lfloor \frac{n}{p} \right\rfloor - \left\lfloor \frac{n}{2p} \right\rfloor - 1 = \frac{n}{2p} + O(1).$$

Since $\pi(n^\delta) = o(n)$, (4.1) gives

$$\beta_n(P) \geq \frac{n}{2} \sum_{p \in Q_\delta(P)} \frac{1}{p} - o(n). \tag{4.2}$$

Let

$$C_\delta(P) := \{p \le n^\delta : \ p \text{ prime and } p \mid u \text{ for some } u \in P\}.$$

Then $Q_\delta(P)$ and $C_\delta(P)$ partition the primes up to n^δ.

We first bound the reciprocal mass of $C_\delta(P)$. For each $p \in C_\delta(P)$ choose one $u(p) \in P$ divisible by p. For any fixed $u \in P$, the primes assigned to u are distinct divisors of u, so

$$\sum_{p:\, u(p)=u} \log p \le \log u \le \log n.$$

Summing over $u \in P$ gives

$$\sum_{p \in C_\delta(P)} \log p \le |P| \log n \le n^\alpha \log n. \tag{4.3}$$

Among all finite prime sets satisfying a log-budget $\sum_{p\in S} \log p \le B$, the reciprocal sum $\sum_{p\in S} 1/p$ is maximized by taking the smallest primes first. Indeed, if S contains a prime q and omits a smaller prime $p < q$, then replacing q by p decreases the log-cost and increases the reciprocal sum. Iterating this exchange proves the claim.

Let z_n be the largest value for which

$$\vartheta(z_n) = \sum_{p \le z_n} \log p \le n^\alpha \log n.$$

By (4.3) and the exchange argument,

$$\sum_{p \in C_\delta(P)} \frac{1}{p} \le \sum_{p \le z_n} \frac{1}{p}. \tag{4.4}$$

The prime number theorem gives

$$z_n = (1 + o(1)) n^\alpha \log n,$$

and hence

$$\log\log z_n = \log\log n + \log\alpha + o(1).$$

Mertens' theorem for primes yields

$$\begin{aligned}
\sum_{p \in Q_\delta(P)} \frac{1}{p} &= \sum_{p \le n^\delta} \frac{1}{p} - \sum_{p \in C_\delta(P)} \frac{1}{p} \\
&\ge \sum_{p \le n^\delta} \frac{1}{p} - \sum_{p \le z_n} \frac{1}{p} \\
&= \log\log(n^\delta) - \log\log z_n + o(1) \\
&= \log\frac{\delta}{\alpha} + o(1).
\end{aligned}$$

Substituting this into (4.2) gives

$$\beta_n(P) \ge \left(\frac{1}{2}\log\frac{\delta}{\alpha} - o(1)\right) n.$$

Letting $\delta \uparrow 1$ proves the theorem. □

Corollary 4.2 (Barrier exponent)**.** *For every fixed $c > 0$,*

$$k_c^\star(n) := \min\{|P| : \ P \subseteq U_n, \ \beta_n(P) \le cn\} \ge n^{e^{-2c} - o(1)}.$$

Proof. If $|P| \le n^\alpha$ and $\beta_n(P) \le cn$, then Theorem 4.1 gives

$$\frac{1}{2}\log\frac{1}{\alpha} \le c + o(1).$$

Thus $\alpha \ge e^{-2c} - o(1)$, which is the claimed exponent lower bound. □

4.2. **The conditional T2 theorem.** The preceding theorem is unconditional but concerns the shield-weight optimization problem rather than the game value itself. A different lower-bound mechanism uses residual upper-half targets of the form acb, where a, c are small primes and b lies in a corresponding large-prime interval. The resulting residual game is modeled by a scored rank-three slot hypergraph.

The finite potential argument which one would like to use in this slot hypergraph is not true for arbitrary finite capture games. The counterexample in Proposition A.1 shows that a unique max-gain Maker move need not dominate every subsequent Breaker deletion. We therefore state the lower bound conditionally.

Definition 4.3 (Safe-edge hypothesis)**.** In each finite graph or scored rank-three hypergraph state arising from the T2 residual slot construction, assume that whenever a positive-weight live edge exists, Maker has a legal live edge f such that, after Maker plays f, every legal Breaker reply leaves the scaled potential Q at least as large as it was before Maker's move.

Theorem 4.4 (Conditional T2 lower bound)**.** *Assume the safe-edge hypothesis for the finite graph and residual slot-hypergraph states generated by the construction in Appendix A. Then for every fixed* $0 < \delta < 1/4$ *there is a constant* $c_\delta > 0$ *such that*

$$L(n) \ge c_\delta \frac{n(\log\log n)^2}{\log n}$$

for all sufficiently large n.

Proof roadmap. The full proof is in Appendix A. The argument has four ingredients.

First, the target family

$$\mathcal{T} = \{acb : \ a < c \le n^\delta, \ b \in (n/(2ac), n/(ac)] \cap \mathbb{P}\}$$

has total mass

$$\sum_{a<c} |(n/(2ac), n/(ac)] \cap \mathbb{P}| \gg_\delta \frac{n(\log\log n)^2}{\log n}.$$

Second, an activation graph game secures a positive fraction of this mass, up to a deletion error which is

$$O\left(\frac{n^{4\delta}}{(\log n)^4}\right) = o\left(\frac{n(\log\log n)^2}{\log n}\right)$$

when $\delta < 1/4$. Third, once a pair (a, c) is secured, each residual target acb has only the three harmful slot divisors b, ab, cb; exact target plays score only their own target. Fourth, the safe-edge hypothesis gives a one-eighth finite potential bound in the residual scored slot hypergraph. These ingredients produce a positive constant multiple of the initial target mass as actual game moves. □

5. The Exact 5/24 First-Hit Cover

This section determines the minimum number of upper-half elements needed to hit every lower-half element by divisibility.

Theorem 5.1 (First-hit cover constant)**.** *Let*

$$\tau(n) := \min\{|C| : \ C \subseteq U_n, \ \forall x \in L_n \ \exists u \in C \ \textit{with } x \mid u\}.$$

Then

$$\tau(n) = \frac{5}{24}n + O(1).$$

Proof. We prove matching upper and lower bounds.

For the upper bound, define

$$H_n := \{u \in U_n : \ u \equiv 2 \pmod 4\} \cup \{u \in U_n : \ u > 2n/3, \ u \equiv 0 \pmod 4\}.$$

We show that every $x \in L_n$ divides some element of H_n.

If $x > n/3$, then $2x \in U_n$. If x is odd, then $2x \equiv 2 \pmod 4$; if x is even, then $2x \equiv 0 \pmod 4$ and $2x > 2n/3$. Hence $2x \in H_n$.

If $n/4 < x \le n/3$, then for odd x the element $2x$ lies in U_n and is 2 modulo 4. For even x, the element $3x$ satisfies $3x > 3n/4 > 2n/3$ and $3x \le n$; it is even, and is therefore either 2 or 0 modulo 4, with the latter case covered by the condition $3x > 2n/3$. Thus $3x \in H_n$.

If $n/6 < x \le n/4$, then $4x \in U_n$, $4x > 2n/3$, and $4x \equiv 0 \pmod 4$, so $4x \in H_n$.

Finally suppose $x \le n/6$. The interval

$$\left(\frac{2n}{3x}, \frac{n}{x}\right]$$

has length $n/(3x) \ge 2$, and hence contains an even integer k. Then $u = kx$ satisfies $2n/3 < u \le n$ and is even. If $u \equiv 2 \pmod 4$ then $u \in H_n$ by the first clause; if $u \equiv 0 \pmod 4$ then $u \in H_n$ by the second clause. In either case $x \mid u$.

The two clauses defining H_n are disjoint. Therefore

$$|H_n| = \#\{u \in (n/2, n] : u \equiv 2 \pmod 4\} + \#\{u \in (2n/3, n] : u \equiv 0 \pmod 4\} = \frac{n}{8} + \frac{n}{12} + O(1) = \frac{5n}{24} + O(1).$$

Thus $\tau(n) \le 5n/24 + O(1)$.

For the lower bound, define

$$P_n := \{m : \ n/3 < m \le n/2\} \cup \{m : \ n/4 < m \le n/3, \ m \text{ odd}\}.$$

Then

$$|P_n| = \frac{n}{6} + \frac{1}{2}\left(\frac{n}{3} - \frac{n}{4}\right) + O(1) = \frac{5n}{24} + O(1).$$

The set P_n is an antichain. Indeed, if $x < y$ and $x \mid y$, then $y \ge 2x$. If $y \le n/3$, then $x \le n/6 < n/4$, impossible for $x \in P_n$. If $n/3 < y \le n/2$, then $x \le n/4$, again impossible for $x \in P_n$.

We claim that any $u \in U_n$ is divisible by at most one element of P_n. If $x \in P_n$ divides u, then $x > n/4$ and $u \le n$, so

$$1 < \frac{u}{x} < 4.$$

Thus u/x is either 2 or 3. If two distinct elements of P_n divide u, they must be $u/2$ and $u/3$. In particular $6 \mid u$. But then $u/3$ is even. It cannot lie in the first component $(n/3, n/2]$ because $u/3 \le n/3$, and it cannot lie in the second component because that component requires oddness. Hence at most one element of P_n divides u.

Now let $C \subseteq U_n$ be any cover of L_n. Each $x \in P_n$ must divide some element of C, and the preceding paragraph shows that different $x \in P_n$ require different elements of C. Therefore

$$|C| \ge |P_n| = \frac{5n}{24} + O(1).$$

This proves the matching lower bound. □

6. Intermediate Upper Bounds

The main upper bound in Section 7 refines a simpler odd-prime-prefix method. We present the two intermediate versions because their proofs isolate the compression and Bonferroni ideas used later.

6.1. **Odd-part compression.** The following elementary lemma is used throughout the upper-bound arguments.

Lemma 6.1 (Odd-part injection)**.** *Let A be a divisibility antichain. The map*

$$\varphi(x) := \frac{x}{2^{v_2(x)}}$$

is injective on A.

Proof. If $\varphi(x) = \varphi(y) = m$, then $x = 2^a m$ and $y = 2^b m$. If $a < b$, then $x \mid y$; if $b < a$, then $y \mid x$. Since A is an antichain, both cases force $x = y$. □

6.2. **The** 13/36 **bound.**

Theorem 6.2. *Shortener has a strategy forcing*

$$L(n) \le \left(\frac{13}{36} + o(1)\right) n.$$

Proof. Let

$$k := \left\lfloor \frac{\sqrt{n}}{\log n} \right\rfloor.$$

For her first k turns, Shortener plays the smallest legal odd prime; after that she plays arbitrarily. Let

$$q_1 < q_2 < \cdots < q_k$$

be the odd primes she obtains.

We first bound these primes. Fix $\varepsilon > 0$. If, before Shortener's jth prefix move, every odd prime up to

$$X = (3/2 + \varepsilon) j \log n$$

were blocked, then the total odd-prime Chebyshev mass up to X would be covered by the earlier Shortener primes together with the odd prime divisors of Prolonger's first j moves. Each Prolonger move contributes at most $\log n$ of prime-divisor log-mass. By induction on j, the previous Shortener primes contribute at most

$$\sum_{i<j} \log q_i \le \left(\frac{1}{2} + o(1)\right) j \log n,$$

because $j \le \sqrt{n}/\log n$. Thus the blocked log-mass is at most $(3/2 + o(1)) j \log n$, while the prime number theorem gives

$$\vartheta_{\mathrm{odd}}(X) = (3/2 + \varepsilon + o(1)) j \log n,$$

a contradiction for large n. Hence

$$q_j \le (3/2 + \varepsilon) j \log n$$

uniformly for $j \le k$, and therefore

$$\sum_{j=1}^{k} \frac{1}{q_j} \ge \frac{1}{3} - o(1). \tag{6.1}$$

Let A be the final antichain. All elements other than the prefix primes have odd part not divisible by any q_j. By Lemma 6.1,

$$|A| \le k + N_D(n),$$

where $D = \{q_1, \ldots, q_k\}$ and $N_D(n)$ is the number of odd integers $m \le n$ not divisible by any prime in D.

Choose t minimal such that

$$s_t := \sum_{j \le t} \frac{1}{q_j} \ge \frac{1}{3} - o(1),$$

and put $E = \{q_1, \dots, q_t\}$. Since every q_j is an odd prime, $1/q_j \le 1/3$, so

$$s_t \in \left[\frac{1}{3} - o(1),\ \frac{2}{3} + o(1)\right] \subset [0,1]$$

for large n. Because $E \subseteq D$, we have $N_D(n) \le N_E(n)$.

Second-order Bonferroni on the events $q \mid m$ over odd $m \le n$ gives

$$N_E(n) \le \frac{n}{2}\left(1 - s_t + \frac{s_t^2}{2}\right) + O(t^2).$$

Here $t \le k$, so $O(t^2) = O(n/\log^2 n) = o(n)$. The function $f(s) = 1 - s + s^2/2$ is decreasing on $[0,1]$, and therefore

$$f(s_t) \le f(1/3 - o(1)) = \frac{13}{18} + o(1).$$

Consequently

$$|A| \le k + N_D(n) \le \frac{13}{36}n + o(n).$$

□

6.3. **The** 5/16 **bound.**

Theorem 6.3. *Shortener has a strategy forcing*

$$L(n) \le \left(\frac{5}{16} + o(1)\right) n.$$

Proof. Fix $A_0 > 2$ and set

$$k := \left\lfloor \frac{n}{2A_0 \log n} \right\rfloor.$$

Shortener plays the smallest legal odd prime for her first k turns. The same Chebyshev log-mass induction, now with $j \le n/(2A_0 \log n)$, gives

$$q_j \le A_0 j \log n \qquad (1 \le j \le k).$$

Indeed, if all odd primes below $A_0 j \log n$ were blocked, Prolonger's first j moves would contribute at most $j \log n$, and the previous Shortener primes would contribute at most $(1 + o(1)) j \log n$; this is less than $A_0 j \log n$ because $A_0 > 2$.

It follows that

$$\sum_{j=1}^{k} \frac{1}{q_j} \ge \frac{1}{A_0} - o(1).$$

As in the proof of Theorem 6.2, use the odd-part injection and truncate the prime set at the first partial sum

$$s_t \ge \frac{1}{A_0} - o(1).$$

Since $1/q_j \le 1/3$ and $A_0 > 2$, the partial sum remains in $[0,1]$. The second-order Bonferroni estimate gives

$$L(n) \le \frac{n}{2}\left(1 - \frac{1}{A_0} + \frac{1}{2A_0^2}\right) + o(n).$$

Letting $A_0 \downarrow 2$ yields

$$L(n) \le \frac{n}{2}\left(1 - \frac{1}{2} + \frac{1}{8}\right) + o(n) = \frac{5}{16}n + o(n).$$

The pair-intersection error is $o(n)$ because only prime pairs with product at most n contribute an $O(1)$ floor error, and

$$\#\{p < q : \ pq \le n\} \ll \sum_{p \le \sqrt{n}} \pi(n/p) \ll \frac{n}{\log n} \sum_{p \le \sqrt{n}} \frac{1}{p} = o(n).$$

□

7. The Main Upper Bound

In this section we prove the paper's headline upper bound

$$L(n) \le \left(\frac{\mathcal{W}_4}{2} + o(1) \right) n < 0.19n.$$

The proof uses the following explicit strategy. Shortener plays the smallest legal odd prime for a long initial prefix of

$$K = \left\lfloor \frac{(1-\varepsilon)n}{2\log n} \right\rfloor$$

turns. Let

$$q_1 < q_2 < \cdots < q_K$$

be the odd primes she plays during that prefix. The argument has four layers:

(1) a local density law for the counting function of the q_j;
(2) a monotone envelope and inversion producing a comparison sequence $b_j \ge q_j$ with explicit moment limits;
(3) a fourth-order Bonferroni comparison theorem for arbitrary increasing odd-prime comparison sequences;
(4) a prime-rounding bridge that replaces the real sequence b_j by actual odd primes p_j, using the key half-density slack supplied by the prime-rounding bridge of Theorem 7.4.

7.1. **The local density.** Write

$$S_K(X) := \#\{j \le K : q_j \le X\}, \qquad u := \frac{\log X}{\log n}.$$

On the interval

$$u \in \left(\frac{1}{h+1}, \frac{1}{h} \right],$$

the local prime-counting law identifies the limiting captured-prime density as

$$\rho(u) = \frac{1}{(\lfloor 1/u \rfloor + 1)u} = \frac{1}{(h+1)u}.$$

Proposition 7.1 (Local density away from breakpoints)**.** *Fix $H \ge 2$, and set*

$$\tau_H := \frac{1}{8H^2}.$$

For $1 \le h \le H-1$, define

$$\alpha_h := \frac{1}{h+1} + \tau_H, \qquad \beta_h := \frac{1}{h} - \tau_H, \qquad I_h := [\alpha_h, \beta_h].$$

Then there exists a function $\xi_H(n) \to 0$ such that, for all sufficiently large n, for every $1 \le h \le H-1$ and every $u \in I_h$,

$$S_K(n^u) \ge \big(1 - \xi_H(n)\big) \frac{n^u}{(h+1)u \log n}.$$

Proof. The intervals I_h are nonempty, because

$$\beta_h - \alpha_h = \frac{1}{h} - \frac{1}{h+1} - 2\tau_H \geq \frac{1}{H(H-1)} - \frac{1}{4H^2} > 0.$$

Fix $1 \leq h \leq H-1$ and $u \in I_h$, and put

$$\eta := \frac{\tau_H}{2}, \qquad X := n^u, \qquad Y := n^{u-\eta}.$$

Since $u \in I_h$,

$$u - \eta \geq \frac{1}{h+1} + \frac{\tau_H}{2} > \frac{1}{h+1}, \qquad u \leq \frac{1}{h} - \tau_H < \frac{1}{h}.$$

Hence the entire interval $[u-\eta, u]$ stays inside $\left(\frac{1}{h+1}, \frac{1}{h}\right)$, so the local prime-count-per-range lemma applies with the constant packet ratio $1/(h+1)$. It yields

$$S(X) \geq \frac{\pi(X) - \pi(Y)}{h+1} - O_H(1),$$

where $S(X)$ denotes the number of odd primes played by Shortener up to the first time her move exceeds X.

We claim that $S(X) = S_K(X)$ for all $u \in \bigcup_{h=1}^{H-1} I_h$ once n is large enough. Indeed, $u \leq \beta_1 = 1 - \tau_H$, so by the prime number theorem,

$$\pi(n^u) \ll \frac{n^{1-\tau_H}}{\log n} = o\left(\frac{n}{\log n}\right) = o(K).$$

Therefore all odd primes at most $X = n^u$ are exhausted before the prefix of length K, and the local counting function $S(X)$ coincides with the truncated counting function $S_K(X)$.

Substituting $X = n^u$ and $Y = n^{u-\eta}$ gives

$$S_K(n^u) \geq \frac{\pi(n^u) - \pi(n^{u-\eta})}{h+1} - O_H(1).$$

Now u ranges over the fixed compact interval I_h, so the prime number theorem is uniform there:

$$\pi(n^u) = \big(1 + o_H(1)\big)\frac{n^u}{u \log n}.$$

Also $u - \eta \leq u - \tau_H/2$, hence

$$\frac{\pi(n^{u-\eta})}{n^u/(u \log n)} \ll_H u\, n^{-\eta} = o_H(1)$$

uniformly for $u \in I_h$. Since there are only finitely many blocks $1 \leq h \leq H-1$, we may absorb the error terms into a single function $\xi_H(n) \to 0$, valid simultaneously for every h and every $u \in I_h$. This gives

$$S_K(n^u) \geq \big(1 - \xi_H(n)\big)\frac{n^u}{(h+1)u \log n},$$

as claimed. □

7.2. **Moment constants and the Bonferroni coefficient.** Define

$$J_r := \frac{1}{r!} \int_{\substack{u_1 + \cdots + u_r \leq 1 \\ u_i \in (0,1]}} \prod_{i=1}^{r} \rho(u_i)\, du_1 \cdots du_r,$$

and

$$\mathcal{W}_4 := 1 - J_1 + J_2 - J_3 + J_4.$$

Numerical evaluation gives

$$\mathcal{W}_4 = 0.3794224, \qquad \frac{\mathcal{W}_4}{2} = 0.1897112 < 0.19.$$

For any increasing sequence $b_1 \le \cdots \le b_K$ of positive reals, define the truncated factorial-moment sums

$$T_r^{(b)}(n) := \sum_{\substack{1\le j_1<\cdots<j_r\le K\\ b_{j_1}\cdots b_{j_r}\le n}} \frac{1}{b_{j_1}\cdots b_{j_r}} \qquad (1\le r\le 4).$$

7.3. **Monotone envelope and inversion.** The local density from Proposition 7.1 is piecewise-defined, so one must build a monotone global comparison sequence before applying the Bonferroni comparison.

Proposition 7.2 (Envelope and inversion)**.** *Fix $H \ge 2$, and define $\tau_H, \alpha_h, \beta_h, I_h$ as in Proposition 7.1. Put*

$$G_H := \bigcup_{h=1}^{H-1} I_h,$$

and for $1 \le r \le 4$ define

$$J_r^{(H)} := \frac{1}{r!}\int_{\substack{u_1+\cdots+u_r\le 1\\ u_i\in G_H}} \prod_{i=1}^{r} \rho(u_i)\, du_1\cdots du_r.$$

Then for every sufficiently large n there exists a monotone function $C_{H,n} : [2,n] \to \{0,1,\dots,K\}$ such that:

(1) *$C_{H,n}(X) \le S_K(X)$ for all $2 \le X \le n$;*
(2) *if*
$$b_j := \inf\{X \in [2,n] : C_{H,n}(X) \ge j\},$$
then $q_j \le b_j$ for every $1 \le j \le K$;
(3) *for every fixed $r \in \{1,2,3,4\}$,*
$$T_r^{(b)}(n) = J_r^{(H)} + o_H(1) \qquad (n\to\infty);$$
(4) *for every fixed $r \in \{1,2,3,4\}$,*
$$0 \le J_r - J_r^{(H)} \le \frac{5r}{4H\, r!}.$$

In particular, there exists an integer-valued function $H_*(n) \to \infty$ and a corresponding choice of comparison sequences

$$b_1^{(n)} \le \cdots \le b_K^{(n)}$$

obtained from the above construction with $H = H_*(n)$ such that

$$T_r^{(b^{(n)})}(n) = J_r + o(1) \qquad (1\le r\le 4).$$

Proof. By Proposition 7.1, there is a function $\xi_H(n) \to 0$ such that for every large enough n, every $1 \le h \le H-1$, and every $u \in I_h$,

$$S_K(n^u) \ge \big(1-\xi_H(n)\big)\frac{n^u}{(h+1)u\log n}.$$

Define

$$A_{n,h}(u) := \big(1-\xi_H(n)\big)\frac{n^u}{(h+1)u\log n} \qquad (u\in I_h).$$

For fixed H and all sufficiently large n, the functions $A_{n,h}$ are strictly increasing on I_h, since

$$A'_{n,h}(u) = \big(1-\xi_H(n)\big)\frac{n^u}{(h+1)u}\left(1-\frac{1}{u\log n}\right),$$

and $u \ge \alpha_{H-1} > 1/H$.

Set

$$L_h := \lfloor A_{n,h}(\alpha_h)\rfloor, \qquad R_h := \lfloor A_{n,h}(\beta_h)\rfloor \qquad (1\le h\le H-1).$$

Then $L_h \le R_h$. Moreover, for $2 \le h \le H-1$,

$$\frac{A_{n,h-1}(\alpha_{h-1})}{A_{n,h}(\beta_h)} = n^{2\tau_H} \cdot \frac{(h+1)(1/h-\tau_H)}{h(1/h+\tau_H)} \to \infty,$$

so $R_h \le L_{h-1}$ for all large n. Also

$$R_1 \ll \frac{n^{1-\tau_H}}{\log n} = o(K),$$

again for large n. Thus, after increasing n if necessary,

$$0 \le L_{H-1} \le R_{H-1} \le L_{H-2} \le R_{H-2} \le \cdots \le L_1 \le R_1 < K.$$

We now define the cumulative lower envelope by

$$C_{H,n}(X) := \begin{cases} 0, & 2 \le X < n^{\alpha_{H-1}}, \\ \lfloor A_{n,h}(\log_n X) \rfloor, & n^{\alpha_h} \le X \le n^{\beta_h} \quad (1 \le h \le H-1), \\ R_h, & n^{\beta_h} < X < n^{\alpha_{h-1}} \quad (2 \le h \le H-1), \\ R_1, & n^{\beta_1} < X < n, \\ K, & X = n. \end{cases}$$

This function is monotone nondecreasing: it is nondecreasing on each genuine block because $A_{n,h}$ is increasing, it is constant on each gap, and the relations $R_h \le L_{h-1}$ guarantee monotonicity across adjacent pieces.

We next prove that $C_{H,n}(X) \le S_K(X)$ for all $2 \le X \le n$. If $X \in [n^{\alpha_h}, n^{\beta_h}]$, then

$$C_{H,n}(X) = \lfloor A_{n,h}(\log_n X) \rfloor \le S_K(X) \qquad \text{by the definition of } A_{n,h}.$$

If $X \in (n^{\beta_h}, n^{\alpha_{h-1}})$ with $2 \le h \le H-1$, then

$$C_{H,n}(X) = R_h \le S_K(n^{\beta_h}) \le S_K(X),$$

since S_K is nondecreasing. The same argument gives $C_{H,n}(X) = R_1 \le S_K(X)$ on (n^{β_1}, n), while $C_{H,n}(X) = 0 \le S_K(X)$ below $n^{\alpha_{H-1}}$ and $C_{H,n}(n) = K = S_K(n)$. This proves part (1).

Define

$$b_j := \inf\{X \in [2,n] : C_{H,n}(X) \ge j\} \qquad (1 \le j \le K).$$

Because $C_{H,n}$ is nondecreasing, the sequence $(b_j)_{j=1}^K$ is nondecreasing. Moreover,

$$S_K(b_j) \ge C_{H,n}(b_j) \ge j,$$

so the jth prime in the prime prefix satisfies $q_j \le b_j$. This proves part (2).

For the moment asymptotics, write

$$u_j := \log_n b_j, \qquad w_j := \frac{1}{b_j} = n^{-u_j}, \qquad \mu_{H,n} := \sum_{j=1}^{K} w_j\, \delta_{u_j}.$$

Split $\mu_{H,n} = \nu_{H,n} + \lambda_{H,n}$, where $\nu_{H,n}$ is the contribution from the genuine interval blocks $L_h < j \le R_h$, and $\lambda_{H,n}$ is the contribution from the flat filler blocks.

The filler blocks have negligible reciprocal mass. For the bottom block,

$$|\lambda_{H,n}^{\text{bot}}| = L_{H-1} n^{-\alpha_{H-1}} \le A_{n,H-1}(\alpha_{H-1}) n^{-\alpha_{H-1}} = \frac{1-\xi_H(n)}{H\alpha_{H-1}\log n} = O_H\left(\frac{1}{\log n}\right).$$

For a gap block between I_h and I_{h-1}, where $2 \le h \le H-1$,

$$|\lambda_{H,n}^{(h)}| = (L_{h-1} - R_h) n^{-\alpha_{h-1}} \le L_{h-1} n^{-\alpha_{h-1}} \le \frac{1-\xi_H(n)}{h\,\alpha_{h-1}\log n} = O_H\left(\frac{1}{\log n}\right).$$

For the top block,

$$|\lambda_{H,n}^{\mathrm{top}}| = (K - R_1)n^{-1} \le \frac{K}{n} = O\left(\frac{1}{\log n}\right).$$

Since there are only $O_H(1)$ filler blocks,

$$|\lambda_{H,n}| = O_H\left(\frac{1}{\log n}\right) = o_H(1).$$

We next analyze the genuine blocks. Fix $1 \le h \le H-1$. On the block $L_h < j \le R_h$, the value $u_j =: u_{h,j} \in I_h$ is defined by $A_{n,h}(u_{h,j}) = j$. Therefore, for every $f \in C^1([0,1])$,

$$\int f\, d\nu_{H,n}^{(h)} = \sum_{j=L_h+1}^{R_h} f(u_{h,j}) n^{-u_{h,j}} = \int_{\alpha_h}^{\beta_h} f(u) n^{-u}\, d\lfloor A_{n,h}(u)\rfloor.$$

Let

$$R_{n,h}(u) := \lfloor A_{n,h}(u)\rfloor - A_{n,h}(u), \qquad |R_{n,h}(u)| \le 1.$$

Applying Stieltjes integration by parts with $g(u) := f(u)n^{-u}$ gives

$$\int_{\alpha_h}^{\beta_h} g(u)\, d\lfloor A_{n,h}(u)\rfloor = \int_{\alpha_h}^{\beta_h} g(u)\, dA_{n,h}(u) + E_{n,h}(f),$$

where

$$|E_{n,h}(f)| \le 2\|g\|_\infty + \int_{\alpha_h}^{\beta_h} |g'(u)|\, du.$$

Since

$$g'(u) = \big(f'(u) - (\log n) f(u)\big) n^{-u},$$

we obtain

$$|E_{n,h}(f)| \le 2\|f\|_\infty n^{-\alpha_h} + (\beta_h - \alpha_h)\|f'\|_\infty n^{-\alpha_h} + (\beta_h - \alpha_h)(\log n)\|f\|_\infty n^{-\alpha_h} = o_H(1).$$

On the other hand,

$$n^{-u} A'_{n,h}(u) = \big(1 - \xi_H(n)\big)\rho(u)\left(1 - \frac{1}{u \log n}\right) \qquad (u \in I_h),$$

because $\rho(u) = 1/((h+1)u)$ on I_h. Hence

$$\int_{\alpha_h}^{\beta_h} f(u) n^{-u}\, dA_{n,h}(u) = \big(1 - \xi_H(n)\big) \int_{\alpha_h}^{\beta_h} f(u)\rho(u)\left(1 - \frac{1}{u \log n}\right) du = \int_{I_h} f(u)\rho(u)\, du + o_H(1),$$

uniformly for fixed f. Combining the last two displays yields

$$\int f\, d\nu_{H,n}^{(h)} = \int_{I_h} f(u)\rho(u)\, du + o_H(1).$$

Summing over h and using $|\lambda_{H,n}| = o_H(1)$, we find that for every $f \in C^1([0,1])$,

$$\int f\, d\mu_{H,n} = \int_{G_H} f(u)\rho(u)\, du + o_H(1).$$

In particular, $\mu_{H,n}$ converges weakly to the absolutely continuous measure

$$\mu_H := \rho(u)\mathbf{1}_{G_H}(u)\, du.$$

Fix $r \in \{1,2,3,4\}$, and write

$$\Sigma_r := \{(u_1, \dots, u_r) \in [0,1]^r : u_1 + \cdots + u_r \le 1\}.$$

Then

$$\int_{\Sigma_r} d\mu_{H,n}^{\otimes r} = \sum_{j_1,\dots,j_r} \mathbf{1}_{\{u_{j_1} + \cdots + u_{j_r} \le 1\}} \prod_{t=1}^{r} w_{j_t}.$$

The contribution of distinct ordered r-tuples is exactly $r!\,T_r^{(b)}(n)$, because $u_{j_1}+\cdots+u_{j_r}\le 1$ is equivalent to $b_{j_1}\cdots b_{j_r}\le n$. The repeated-index contribution is negligible:

$$\max_j w_j \le n^{-\alpha_{H-1}} \to 0,$$

while $\mu_{H,n}([0,1]) = O_H(1)$ by the weak convergence with $f\equiv 1$. Therefore

$$0 \le \int_{\Sigma_r} d\mu_{H,n}^{\otimes r} - r!\,T_r^{(b)}(n) \le \binom{r}{2}\big(\max_j w_j\big)\mu_{H,n}([0,1])^{r-1} = o_H(1).$$

Since μ_H is absolutely continuous, its product measure gives zero mass to the boundary $\partial\Sigma_r$. Weak convergence therefore implies

$$\int_{\Sigma_r} d\mu_{H,n}^{\otimes r} \to \int_{\Sigma_r} d\mu_H^{\otimes r} = \int_{\substack{u_1+\cdots+u_r\le 1\\ u_i\in G_H}} \prod_{i=1}^r \rho(u_i)\,du_1\cdots du_r = r!\,J_r^{(H)}.$$

Combining the last two displays proves part (3):

$$T_r^{(b)}(n) = J_r^{(H)} + o_H(1).$$

It remains to control the excised breakpoint region. Since $0<\rho(u)<1$ on $(0,1]$, we have

$$0 \le J_r - J_r^{(H)} \le \frac{1}{r!}\,\lambda_r\big(\{(u_1,\dots,u_r)\in[0,1]^r : \exists i,\ u_i\notin G_H\}\big),$$

where λ_r denotes r-dimensional Lebesgue measure. Moreover,

$$|(0,1]\setminus G_H| = \frac{1}{H} + 2(H-1)\tau_H \le \frac{1}{H} + \frac{1}{4H} = \frac{5}{4H}.$$

By the union bound in $[0,1]^r$,

$$\lambda_r\big(\{(u_1,\dots,u_r)\in[0,1]^r : \exists i,\ u_i\notin G_H\}\big) \le r\,|(0,1]\setminus G_H| \le \frac{5r}{4H}.$$

This proves part (4), and the final statement follows immediately by first letting $n\to\infty$ for fixed H, and then letting $H\to\infty$.

To obtain a single comparison sequence for each n, choose N_H so large that for all $n\ge N_H$ the above construction is valid and, for every $1\le r\le 4$,

$$\big|T_r^{(b)}(n) - J_r^{(H)}\big| \le \frac{1}{H}.$$

After enlarging if necessary, we may assume N_H is increasing in H. Define

$$H_*(n) := \max\{H\ge 2 : N_H\le n\},$$

and let $b_1^{(n)}\le\cdots\le b_K^{(n)}$ be the comparison sequence produced with parameter $H=H_*(n)$. Then $H_*(n)\to\infty$, and for every $1\le r\le 4$,

$$\big|T_r^{(b^{(n)})}(n) - J_r\big| \le \big|T_r^{(b^{(n)})}(n) - J_r^{(H_*(n))}\big| + \big|J_r^{(H_*(n))} - J_r\big| \le \frac{1}{H_*(n)} + \frac{5r}{4H_*(n)r!},$$

which tends to 0. This proves the final statement. □

7.4. **The prime-sequence Bonferroni comparison theorem.** The comparison theorem must be stated for *prime sequences*, not for arbitrary reals; the correct statement is the following.

Theorem 7.3 (Prime-sequence Bonferroni–4 comparison)**.** *Let $q_1 < \cdots < q_K$ be the actual odd primes played by Shortener during the prime prefix, and let $p_1 < \cdots < p_K$ be any increasing sequence of odd primes with*

$$q_j \le p_j \qquad (1 \le j \le K).$$

Assume that for $1 \le r \le 4$,

$$T_r^{(p)}(n) = \Lambda_r + o(1).$$

Then

$$L(n) \le \frac{n}{2}\left(1 - \Lambda_1 + \Lambda_2 - \Lambda_3 + \Lambda_4 + o(1)\right).$$

Proof. Let A be the final played set. If the game ends before Shortener completes the prefix of length K, then $L(n) = o(n)$ and there is nothing to prove, so assume $q_1, \dots, q_K$ were all played.

Define the odd-part map

$$\varphi(x) := \frac{x}{2^{v_2(x)}}.$$

If $\varphi(x) = \varphi(y) = m$, then $x = 2^a m$ and $y = 2^b m$. If $a \ne b$, one of x and y divides the other, contradicting that A is an antichain. So φ is injective on A.

Since each $q_j \in A$, no other element of A can be divisible by any q_j. Because q_j is odd, divisibility by q_j is preserved under the odd-part map. Hence

$$\varphi(A) \subseteq \{m \le n : m \text{ odd and } q_j \nmid m \text{ for all } j\} \cup \{q_1, \dots, q_K\}.$$

Therefore

$$L(n) \le N(q_1, \dots, q_K) + K = N(q_1, \dots, q_K) + o(n),$$

where $N(q_1, \dots, q_K)$ counts odd integers up to n avoiding all q_j.

Now replace the actual sequence q_j by the larger prime sequence p_j. The count of odd integers avoiding a finite prime set is monotone in each prime, so

$$N(q_1, \dots, q_K) \le N(p_1, \dots, p_K).$$

Applying fourth-order Bonferroni to the divisibility events $p_j \mid m$ over odd integers $m \le n$ gives

$$\begin{aligned} N(p_1, \dots, p_K) \le \frac{n}{2}\Big(1 - T_1^{(p)}(n) + T_2^{(p)}(n) - T_3^{(p)}(n) + T_4^{(p)}(n)\Big) \\ + O(M_2 + M_3 + M_4), \end{aligned}$$

where M_r is the number of odd integers $\le n$ with at least r distinct prime divisors from the set $\{p_j\}$. The standard squarefree divisor-count bound gives

$$M_r = O_r\left(\frac{n(\log\log n)^{r-1}}{\log n}\right) = o(n) \qquad (2 \le r \le 4),$$

so the error is $o(n)$. Substituting $T_r^{(p)}(n) = \Lambda_r + o(1)$ yields the stated inequality. □

7.5. **The prime-rounding bridge.** It remains to convert the real comparison sequence b_j from Proposition 7.2 into an actual increasing odd-prime sequence. The naive claim

$$p_j/b_j = 1 + O(1/\log b_j)$$

uniformly in j is *not* justified by the prime number theorem alone and must not be used. The correct bridge is the half-density construction of Theorem 7.4.

Theorem 7.4 (Prime-rounding bridge)**.** *Fix $0 < \varepsilon < 1$, and let*

$$K = \left\lfloor \frac{(1-\varepsilon)n}{2\log n} \right\rfloor.$$

There exists a choice of comparison sequences

$$b_1^{(n)} \le \cdots \le b_K^{(n)}$$

from the construction of Proposition 7.2, and there exists an increasing sequence of odd primes

$$p_1 < \cdots < p_K$$

such that

$$q_j \le p_j \qquad (1 \le j \le K)$$

and

$$T_r^{(b^{(n)})}(n) = J_r + o(1) \qquad (1 \le r \le 4).$$

Moreover the primes may be chosen so that, for each fixed $1 \le r \le 4$,

$$T_r^{(p)}(n) = T_r^{(b^{(n)})}(n) + o(1) = J_r + o(1).$$

Proof. We first prove a fixed-parameter statement. Fix $H \ge 2$ and $\lambda > 1$. Let $\tau_H, \alpha_h, \beta_h, I_h, G_H$ be as in Proposition 7.2, and let $b_1, \dots, b_K$ be the sequence obtained by inverting the envelope $C_{H,n}$ for this fixed H. We shall construct increasing odd primes $p_1, \dots, p_K$ such that

$$q_j \le p_j \qquad (1 \le j \le K)$$

and, for $1 \le r \le 4$,

$$T_r^{(p)}(n) = J_r^{(H)} + O_{H,r}(\lambda - 1) + o_{H,\lambda}(1). \tag{PB1}$$

Here $n \to \infty$ with H, λ, ε fixed.

Write

$$B_{H,n}(X) := \#\{j : b_j \le X\}.$$

Since $b_j = \inf\{X : C_{H,n}(X) \ge j\}$ and $C_{H,n}$ is nondecreasing and integer-valued,

$$B_{H,n}(X) = C_{H,n}(X) \qquad (2 \le X \le n).$$

We need the following interval count, which is the point at which the half-density slack enters. Fix $\Delta > 0$. For all sufficiently large n, uniformly in $1 \le h \le H-1$ and in all X for which $[X, (1+\Delta)X] \subseteq [n^{\alpha_h}, n^{\beta_h}]$,

$$B_{H,n}((1+\Delta)X) - B_{H,n}(X) \le \left(\frac{1}{2} + o_{H,\Delta}(1)\right) \frac{\Delta X}{\log X}. \tag{PB2}$$

Indeed, put $u = \log_n X$ and $\theta = \log(1+\Delta)/\log n$. On a genuine block, $B_{H,n}(Y) = \lfloor A_{n,h}(\log_n Y) \rfloor$, up to an error of at most one at each endpoint. Therefore

$$B_{H,n}((1+\Delta)X) - B_{H,n}(X) \le A_{n,h}(u+\theta) - A_{n,h}(u) + 2.$$

For $v \in [u, u+\theta]$ and $Y = n^v$,

$$\frac{d}{dY} A_{n,h}(\log_n Y) = \frac{1 - \xi_H(n)}{(h+1)\log Y}\left(1 - \frac{1}{\log Y}\right) \le \frac{1 + o_H(1)}{(h+1)\log X}.$$

Since $h+1 \ge 2$ and $X \ge n^{\alpha_{H-1}} \ge n^{1/H}$, the endpoint error 2 is $o_{H,\Delta}(\Delta X/\log X)$. This proves (PB2).

Choose $\delta = \delta(\lambda) > 0$ with

$$(1+\delta)^2 \le \lambda.$$

All uses of the prime number theorem below are uniform for $X \geq n^{1/H}$ and for fixed multiplicative intervals, because H and δ are fixed. Thus, uniformly in such X,

$$\#\{p \text{ odd prime} : (1+\delta)X \leq p < (1+\delta)^2 X\} = (1+o_{H,\delta}(1))\frac{\delta(1+\delta)X}{\log X}. \tag{PB3}$$

Combining (PB2) and (PB3), for all sufficiently large n every multiplicative bin $[X,(1+\delta)X)$ contained in a genuine block contains strictly fewer b_j than there are odd primes in the next bin $[(1+\delta)X,(1+\delta)^2X)$.

It remains to keep the bins away from the finitely many artificial boundary layers. Choose a constant $\Gamma_H > 4$, depending only on H, large enough for the reservoir count below. For each genuine block $[n^{\alpha_h}, n^{\beta_h}]$, declare exceptional all indices j for which

$$b_j \in [n^{\alpha_h}, \Gamma_H n^{\alpha_h}] \cup [\Gamma_H^{-1} n^{\beta_h}, n^{\beta_h}],$$

and also all indices lying in the flat filler blocks of the envelope. Denote the exceptional set by $\mathcal{E}$ and its complement by $\mathcal{G}$. For fixed H and large n, these boundary reservoirs are disjoint and occur in the same order as the corresponding indices, because the gaps between successive genuine blocks have multiplicative size $n^{2\tau_H}$.

The exceptional set has negligible reciprocal mass. The flat filler blocks already satisfy the estimate proved in Proposition 7.2,

$$\sum_{j \text{ flat}} \frac{1}{b_j} = O_H\left(\frac{1}{\log n}\right).$$

For a genuine boundary layer, (PB2) with fixed multiplicative length implies

$$\sum_{\substack{j: b_j \in [X, \Gamma_H X] \\ j \text{ genuine}}} \frac{1}{b_j} \ll_{H,\Gamma_H} \frac{1}{\log n},$$

uniformly for $X \geq n^{1/H}$. There are $O_H(1)$ such layers, hence

$$\sum_{j \in \mathcal{E}} \frac{1}{b_j} = O_{H,\lambda}\left(\frac{1}{\log n}\right). \tag{PB4}$$

We next assign primes to the exceptional indices. On each boundary reservoir of the form $[X, \Gamma_H X]$, the number of genuine exceptional indices is at most

$$\left(\frac{1}{2} + o_H(1)\right)\frac{(\Gamma_H - 1)X}{\log X},$$

by (PB2) applied with $\Delta = \Gamma_H - 1$, while the number of odd primes in the same reservoir is

$$(1+o_H(1))\frac{(\Gamma_H - 1)X}{\log X}.$$

At a left endpoint one must also place the adjacent flat atom. Its size is at most the value of the envelope at that endpoint, namely

$$O_H\left(\frac{X}{\log X}\right),$$

with an implied constant depending only on H; the choice of Γ_H ensures that the reservoir contains enough odd primes for both the flat atom and the genuine boundary layer. More explicitly, the sorted lower bounds b_j in the reservoir satisfy the one-dimensional Hall condition: the full reservoir has enough primes by the preceding choice of Γ_H, and every proper suffix $[Y, \Gamma_H X]$ contains at most half as many genuine exceptional b_j as there are odd primes there, by the interval estimate (PB2), uniformly for multiplicative lengths at most Γ_H, and the prime number theorem. Thus greedy assignment to the smallest unused odd prime which is at least the current b_j succeeds inside

the reservoir. The top flat block is placed in $[n, \Gamma_H n]$, which has enough odd primes because $K \le n/(2\log n)$ and $\Gamma_H > 2$. Consequently all exceptional indices can be assigned, in increasing order, to distinct odd primes satisfying $b_j \le p_j$ and hence $q_j \le p_j$. Their total reciprocal mass is

$$\sum_{j\in\mathcal{E}} \frac{1}{p_j} = O_{H,\lambda}\left(\frac{1}{\log n}\right). \tag{PB5}$$

Now consider the nonexceptional indices. In each genuine block, partition the remaining interval into consecutive multiplicative bins

$$[X, (1+\delta)X),$$

discarding at most the first and last partial bin into $\mathcal{E}$. For each full bin, assign the b_j in that bin, in increasing order, to the odd primes in the next bin. The count comparison (PB2)-(PB3) shows that this is possible for all large n. The prime bins are disjoint and increasing, so together with the reservoir assignments this produces one increasing sequence of odd primes. For every $j \in \mathcal{G}$,

$$b_j \le p_j \le (1+\delta)^2 b_j \le \lambda b_j. \tag{PB6}$$

We compare moments. The total reciprocal mass of the b_j is $O_H(1)$ by Proposition 7.2; the same is true for the p_j, by (PB5) and (PB6). Hence every r-tuple containing at least one exceptional index contributes

$$O_{H,r,\lambda}\left(\frac{1}{\log n}\right) = o_{H,\lambda}(1)$$

to either $T_r^{(b)}(n)$ or $T_r^{(p)}(n)$. It remains to compare tuples lying wholly in $\mathcal{G}$.

For such a tuple $I = \{j_1 < \cdots < j_r\}$, set

$$b_I := \prod_{j\in I} b_j, \qquad p_I := \prod_{j\in I} p_j.$$

By (PB6),

$$b_I \le p_I \le \lambda^r b_I, \qquad \lambda^{-r}\frac{1}{b_I} \le \frac{1}{p_I} \le \frac{1}{b_I}.$$

The weight change contributes $O_{H,r}(\lambda - 1)$, because the total b-reciprocal mass is $O_H(1)$. The only remaining discrepancy is the product boundary

$$n/\lambda^r < b_I \le n.$$

For fixed H and λ, this boundary layer is $o_{H,\lambda}(1)$: in logarithmic coordinates it is a strip of width $r\log\lambda/\log n$ around the hyperplane $u_1 + \cdots + u_r = 1$, and the empirical measures from Proposition 7.2 converge to an absolutely continuous limit. Equivalently, one may partition into unit logarithmic cells $[e^a, e^{a+1})$; each cell has reciprocal mass $O_H(1/\log n)$ by (PB2), and only $O_{H,\lambda}((\log n)^{r-1})$ cell patterns meet the strip. Thus

$$T_r^{(p)}(n) = T_r^{(b)}(n) + O_{H,r}(\lambda - 1) + o_{H,\lambda}(1).$$

Using Proposition 7.2, $T_r^{(b)}(n) = J_r^{(H)} + o_H(1)$, and this proves (PB1).

It remains to diagonalize. Choose a sequence $H_m \to \infty$ such that

$$|J_r^{(H_m)} - J_r| \le \frac{1}{m} \qquad (1 \le r \le 4).$$

For each m, choose $\lambda_m > 1$ so close to 1 that the constants implicit in $O_{H_m,r}(\lambda_m - 1)$ in (PB1) are at most $1/m$ for all $1 \le r \le 4$. For each m, the fixed-parameter construction above is valid for all $n \ge N_m$ after enlarging N_m so that also

$$|T_r^{(b)}(n) - J_r^{(H_m)}| \le \frac{1}{m},$$

for every $1 \le r \le 4$. We may take N_m increasing. For $N_m \le n < N_{m+1}$ use the construction with $H = H_m$ and $\lambda = \lambda_m$. This gives $H(n) \to \infty$ slowly enough that the smallest genuine scale

$n^{\alpha_{H(n)-1}}$ still tends to infinity, so the uniform prime-number estimates used above remain valid. The resulting comparison and prime sequences satisfy, for every fixed $1 \le r \le 4$,

$$T_r^{(b^{(n)})}(n) = J_r + o(1), \qquad T_r^{(p)}(n) = T_r^{(b^{(n)})}(n) + o(1) = J_r + o(1).$$

□

7.6. **The finite Bonferroni–4 theorem.**

Theorem 7.5. *Under the piecewise-density prime-prefix strategy,*

$$L(n) \le \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n = (0.1897112 + o(1))n < 0.19n.$$

Proof. By Theorem 7.4, there are a comparison sequence $b_1 \le \cdots \le b_K$ and an increasing odd-prime sequence p_j with $q_j \le p_j$ such that

$$T_r^{(b)}(n) = J_r + o(1) \qquad (1 \le r \le 4).$$

The same theorem gives

$$T_r^{(p)}(n) = J_r + o(1) \qquad (1 \le r \le 4).$$

Applying Theorem 7.3 with $\Lambda_r = J_r$ gives

$$L(n) \le \frac{n}{2}\left(1 - J_1 + J_2 - J_3 + J_4 + o(1)\right) = \left(\frac{\mathcal{W}_4}{2} + o(1)\right) n.$$

Numerical evaluation yields

$$\frac{\mathcal{W}_4}{2} = 0.1897112 < 0.19,$$

so the theorem follows. □

Remark 7.6. The corresponding formal artifact verifies the endgame reduction: once the first four moment terms are sufficiently close to the certified constants $J_1, \ldots, J_4$, the strict inequality $L(n) < 0.19n$ follows eventually. Theorems 7.3 and 7.4 supply those moment hypotheses.

8. Obstructions to Further Improvement

This section records three barriers encountered by natural attempts to improve the upper bound. They are not used in the proof of Theorem 7.5; their purpose is to delineate which proof classes cannot by themselves resolve the asymptotic question.

8.1. **A local-relaxation barrier.**

The first obstruction says that bounded-round local relaxations can have an integrality gap at precisely the scale where one would hope for a short transversal argument. We state the barrier in its cleanest finite form.

Let $K_N^{(\ell)}$ be the complete ℓ-uniform hypergraph on the vertex set $[N]$. A transversal is a set of vertices meeting every ℓ-edge.

Proposition 8.1 (Sherali–Adams transversal barrier)**.** *Let $\ell = \lfloor N/\log N \rfloor$. The integral transversal number of $K_N^{(\ell)}$ is*

$$\tau(K_N^{(\ell)}) = N - \ell + 1 = N - o(N).$$

The fractional transversal number is

$$\tau_f(K_N^{(\ell)}) = \frac{N}{\ell} = (1 + o(1)) \log N.$$

Moreover, after any fixed number $r < \ell$ of Sherali–Adams rounds, the symmetric lifted solution still has objective at most

$$\frac{N}{\ell - r}.$$

Proof. A set $T \subseteq [N]$ meets every ℓ-set if and only if its complement has size at most $\ell - 1$. Hence

$$\tau(K_N^{(\ell)}) = N - \ell + 1.$$

For the fractional relaxation, give every vertex weight $1/\ell$. Each ℓ-edge then has total weight 1, so $\tau_f \leq N/\ell$. Conversely, summing the edge constraints over all ℓ-subsets gives

$$\binom{N-1}{\ell-1} \sum_{i=1}^{N} x_i \geq \binom{N}{\ell},$$

and therefore $\sum_i x_i \geq N/\ell$. This proves $\tau_f = N/\ell$.

For the lifted statement, the same symmetric solution remains feasible after conditioning on any pattern involving fewer than ℓ vertices. If r vertices of an ℓ-edge have already been fixed outside the transversal, the residual constraint asks for weight at least 1 on an $(\ell - r)$-set; the uniform residual assignment $1/(\ell - r)$ satisfies it. The corresponding objective is $N/(\ell - r)$. This is the standard symmetric feasible solution for the r-round Sherali–Adams lift, and gives the claimed upper bound on the relaxation value. □

The proposition shows that local linear relaxations can remain logarithmic in size while the actual separator has linear size.

8.2. **A q-shadow dichotomy.** The next obstruction is an extremal statement for Johnson shadows. It explains why a sparse separator family cannot be forced by expansion alone unless a large pre-existing shadow is already present.

Let $X = \binom{[K]}{q}$ and $Y = \binom{[K]}{h}$, with $q < h$. A q-set Q covers an h-set H if $Q \subseteq H$. Let $R \subseteq Y$ be a live packet of density

$$\eta := \frac{|R|}{|Y|}.$$

Let $D \subseteq X$ be a previously forbidden q-shadow and let $C \subseteq X$ be a set of already claimed q-separators. Define

$$\delta_q := \frac{\binom{K-q}{h-q}}{\binom{K}{h}},$$

the density of h-sets containing a fixed q-set, and write $\sigma_q(D)$ for the fraction of Y covered by D. The normalized second singular value of the q-versus-h inclusion graph satisfies

$$\lambda_q^2 = \frac{q(K-h)}{h(K-q)}.$$

Proposition 8.2 (q-shadow covering dichotomy)**.** *Assume no legal q-separator captures at least*

$$\frac{1}{2}\,\delta_q |R|$$

members of R. Then

$$\sigma_q(D) + |C|\delta_q > 1 - \frac{4\lambda_q^2}{\eta}.$$

Proof. Let $A \subseteq X$ be the legal q-separators, meaning those not already excluded by D or C. By the union bound,

$$\frac{|A|}{|X|} \geq 1 - \sigma_q(D) - |C|\delta_q.$$

The expander-mixing inequality for the inclusion graph between X and Y gives

$$\left| e(A,R) - \frac{|A|}{|X|}\frac{|R|}{|Y|} e(X,Y) \right| \leq \lambda_q\, e(X,Y) \sqrt{\frac{|A|}{|X|}\frac{|R|}{|Y|}}.$$

If

$$\frac{|A|}{|X|} \geq \frac{4\lambda_q^2}{\eta},$$

then the average number of elements of R captured by a member of A is at least half the random expectation, namely at least $\frac{1}{2}\delta_q|R|$. This contradicts the hypothesis that no legal separator captures that many live packet members. Hence

$$\frac{|A|}{|X|} < \frac{4\lambda_q^2}{\eta},$$

and the displayed conclusion follows from the union bound. □

In the central regime relevant to top-facet packet arguments, the inequality says that separator starvation is possible only after the previous shadow or the already claimed separator set has become large. Pure expansion cannot supply an unlimited sequence of fresh separators.

8.3. **A separator-only limitation.** The last obstruction is a finite model for strategies which only play odd carrier separators and ignore dyadic shielding.

Proposition 8.3 (Separator-only limitation)**.** *Fix a finite set P of odd primes and a family $\mathcal{S} \subseteq 2^P$ of odd squarefree carrier supports. Suppose Shortener's planned moves are only odd squarefree separators with supports in $\mathcal{S}$. Then there are dyadic upper-half targets of the form*

$$2^a \prod_{p \in T} p$$

which remain untouched by all separators unless $\mathcal{S}$ covers the corresponding support T itself. In particular, any separator-only proof that does not control all dyadic lifts of the carrier family leaves a positive residual packet.

Proof. An odd separator can only be comparable with an upper-half target $2^a m$ through its odd part m. The dyadic exponent is invisible to the odd separator. Therefore, if a support T is not covered by the chosen odd separator family, every dyadic lift $2^a \prod_{p \in T} p$ with value in $(n/2, n]$ remains incomparable with all separator moves.

For any fixed odd part $m \leq n/2$, there is at least one dyadic lift $2^a m \in (n/2, n]$ unless $m = 1$; choose a maximal with $2^a m \leq n$. If $2^a m \leq n/2$, then $2^{a+1} m \leq n$, contradicting maximality. Thus each uncovered odd support produces an upper-half target not hit by the separator-only family. Summing over an uncovered packet gives the claimed residual. □

The proposition is deliberately modest: it does not rule out separator methods which also track dyadic lifts or exact upper targets. It rules out only the separator-only abstraction in which odd carrier supports are treated as if they were the whole divisibility geometry.

9. Conclusion

The results of this paper narrow the divisibility antichain saturation game in two complementary ways. On the upper-bound side, the main theorem gives

$$L(n) < 0.19n$$

for all sufficiently large n, improving the elementary linear upper-bound picture by using the local density of Shortener's legal-prime prefix and a fourth-order Bonferroni comparison. On the structural side, the shield reduction and Theorem A show that short upper-half shield prefixes cannot prove a linear game lower bound through the weighted shadow functional $\beta_n(P)$.

The conditional T2 theorem points toward a substantially larger lower-bound scale, but it is deliberately stated under the safe-edge hypothesis because the corresponding max-gain principle is false in general finite capture games. The finite graph, hypergraph, and embedding components are robust; the remaining issue is precisely the safety property for the game-generated states.

Three obstructions in Section 8 explain why the next improvements are unlikely to come from purely local relaxations, fixed shadow-cover dichotomies, or separator-only strategies. A sharper upper bound will need to use dynamic information about the actual divisibility game, not merely the static incidence geometry of low-rank shadows. Conversely, an unconditional lower bound beyond the classical prime-multiple argument must overcome Shortener's ability to compress terminal antichains through odd-prime sieving.

The central question remains whether $L(n) = o(n)$. The methods here give new constants, exact auxiliary structure, and a set of formal targets for future work, but they do not settle that asymptotic dichotomy.

Appendix A. Deferred Proofs

A.1. **Conditional proof of the T2 lower bound.** This appendix proves the conditional lower bound in Theorem 4.4. The proof is self-contained: all target families, auxiliary games, weights, activation rules, deletion errors, and asymptotic estimates are defined below. The opening counterexample explains why the finite safe-edge hypothesis is stated explicitly rather than hidden inside a max-gain argument.

Throughout this appendix n is large, $0 < \delta < 1/4$ is fixed, and

$$Y := n^{\delta}.$$

Let $\mathcal{P}_Y$ be the set of odd primes at most Y. For $a < c$ in $\mathcal{P}_Y$, put

$$I_{a,c} := \left(\frac{n}{2ac}, \frac{n}{ac}\right], \qquad B_{a,c} := I_{a,c} \cap \mathbb{P},$$

and define the target family

$$\mathcal{T} := \{\, acb : \ a < c, \ a, c \in \mathcal{P}_Y, \ b \in B_{a,c} \,\}.$$

For large n, every such b is greater than Y: indeed $b > n/(2Y^2) = \frac{1}{2}n^{1-2\delta} > Y$. Thus a, c, b are distinct primes and each target acb lies in $(n/2, n]$.

Proposition A.1 (Max-gain domination is false in rank three)**.** *The following finite scored rank-3 capture state has a unique max-gain Maker move, but Breaker can answer that move by deleting one uncaptured vertex and make the scaled potential decrease.*

Proof. Let the already-captured vertices be

$$C = \{z_{ij} : 1 \le i, j \le 3\},$$

and let u_1, u_2, u_3, v be uncaptured. There are no deleted or scored edges. The live edges are

$$f = \{u_1, u_2, u_3\}, \qquad w(f) = 9,$$

and, for $1 \le i, j \le 3$,

$$e_{ij} = \{v, u_i, z_{ij}\}, \qquad w(e_{ij}) = 4.$$

For a live edge e, write $\Phi(e) = 2^{|e \cap C|} w(e)$ and $Q = 8S + \sum \Phi(e)$.

Before Maker moves, $\Phi(f) = 9$ and $\Phi(e_{ij}) = 8$ for all i, j. If Maker plays f, then the score contribution of f changes by $8w(f) - \Phi(f) = 72 - 9 = 63$, and each of the nine edges e_{ij} doubles from 8 to 16. Hence

$$\Delta(f) = 63 + 9 \cdot 8 = 135.$$

After this move, the vertex v is still uncaptured, and deleting v removes all nine edges e_{ij}, now of potential 16 each. The Breaker loss is

$$9 \cdot 16 = 144 > 135.$$

Thus the full round decreases Q.

It remains only to check that f is the unique max-gain move. If Maker plays some e_{ij}, its own score gain is $8\cdot 4-8=24$. The edge f doubles from 9 to 18, giving gain 9. The two other edges with the same u_i gain 24 each, and the six edges with the other two u-vertices gain 8 each. Therefore

$$\Delta(e_{ij}) = 24+9+2\cdot 24+6\cdot 8 = 129 < 135.$$

So the unique max-gain move is f, and it does not dominate the subsequent vertex deletion at v. □

Proposition A.2 (Conditional finite scaled capture)**.** *Consider a finite weighted incidence game of rank at most* 3 *with the scaled potential*

$$Q := 8S + \sum_{e \text{ live}} 2^{|e\cap C|} w(e),$$

where C is the current captured-vertex set and S is the total scored weight. In the graph variant, Breaker may delete one unclaimed live edge; in the scored hypergraph variant, Breaker may score one live edge.

Assume the following safe-edge hypothesis: at every reachable pre-Maker state with at least one positive-weight live edge, Maker has a legal live edge f such that, after Maker plays f, every legal Breaker reply leaves Q at least as large as it was before Maker's move.

Under this hypothesis, if the game is continued until no positive-weight live edge remains, the final claimed or scored weight is at least one eighth of the initial total edge-weight:

$$S_{\text{fin}} \ge \frac{1}{8}\sum_{e} w(e).$$

Proof. Maker always chooses an edge supplied by the safe-edge hypothesis. By definition of that hypothesis, Q is nondecreasing after every full Maker–Breaker round.

Initially $C=\varnothing$, $S=0$, and all initial edges are live, so

$$Q_0 = \sum_{e} w(e).$$

At terminal time there is no positive-weight live edge, hence $Q_{\text{fin}} = 8S_{\text{fin}}$. Therefore

$$8S_{\text{fin}} = Q_{\text{fin}} \ge Q_0 = \sum_{e} w(e),$$

which is the claimed one-eighth bound.

For the graph variant, a Breaker edge-deletion reply is included in the safe-edge hypothesis. For the scored hypergraph variant, a Breaker scored-edge reply is harmless whenever it occurs, since it changes Q by $8w(e) - 2^{|e\cap C|}w(e) \ge 0$ for rank at most 3; the only nontrivial part of the hypothesis is therefore the post-Maker vertex-deletion bound. □

Proposition A.3 (Initial target mass)**.** *For every fixed* $0<\delta<1/4$,

$$W_0 := \sum_{\substack{a<c\\ a,c\in\mathcal{P}_Y}} |B_{a,c}| \gg_\delta \frac{n(\log\log n)^2}{\log n}.$$

Proof. Since $ac \le Y^2 = n^{2\delta}$, the interval endpoint $X_{a,c} := n/(ac)$ satisfies $X_{a,c} \ge n^{1-2\delta} \to \infty$. By the prime number theorem, uniformly for all such pairs and all sufficiently large n,

$$\pi(X_{a,c}) - \pi(X_{a,c}/2) \ge \kappa_\delta \frac{X_{a,c}}{\log n} = \kappa_\delta \frac{n}{ac\log n}$$

for some positive constant κ_δ. Therefore

$$W_0 \ge \kappa_\delta \frac{n}{\log n} \sum_{\substack{a<c\\ a,c\in\mathcal{P}_Y}} \frac{1}{ac}.$$

Mertens' theorem for primes gives

$$\sum_{p\in\mathcal{P}_Y}\frac{1}{p}=\log\log Y+O(1)=\log\log n+O_\delta(1),$$

and $\sum_p p^{-2}<\infty$. Hence

$$\sum_{a<c\in\mathcal{P}_Y}\frac{1}{ac}=\frac{1}{2}\left[\left(\sum_{p\in\mathcal{P}_Y}\frac{1}{p}\right)^2-\sum_{p\in\mathcal{P}_Y}\frac{1}{p^2}\right]=\left(\frac{1}{2}+o(1)\right)(\log\log n)^2.$$

Combining the last two displays proves the claim. □

Proposition A.4 (Residual divisibility embedding)**.** *Let $\mathcal{T}_*$ be any subfamily of targets $t=acb\in\mathcal{T}$ such that, for every $t=acb\in\mathcal{T}_*$, the numbers a, c, and ac are already unavailable as legal moves. Attach to t the three slot labels*

$$b,\qquad ab,\qquad cb$$

and let $e_t:=\{b,ab,cb\}$. Then the residual divisibility game on $\mathcal{T}_*$ is at least as favorable to Prolonger as the scored rank-3 hypergraph game on the hyperedges $\{e_t : t\in\mathcal{T}_*\}$:

1. *the only future moves that can make $t=acb$ illegal are b, ab, cb, and t itself;*
2. *playing b, ab, or cb deletes exactly the residual targets whose hyperedges contain that slot;*
3. *playing the exact target t kills no other target in $\mathcal{T}_*$, and is therefore a scored-edge move;*
4. *every live hyperedge corresponds to a legal actual move t.*

Proof. Fix $t=acb\in\mathcal{T}_*$. Because $t>n/2$, no proper multiple of t lies in $\{2,\dots,n\}$. Thus a future move can make t illegal only by being t itself or a proper divisor of t. The positive divisors are

$$1,\ a,\ c,\ ac,\ b,\ ab,\ cb,\ acb.$$

The divisor 1 is not a move, while a, c, and ac are unavailable by hypothesis. This proves the first assertion.

For slot incidence, suppose another target is $t'=a'c'b'$. Since $b,b'>Y\geq a,a',c,c'$, the divisibility $b\mid t'$ holds iff $b=b'$. Thus a move b kills exactly the targets in the b-fiber. Similarly, if $ab\mid a'c'b'$, the large prime factor forces $b=b'$, and after cancellation a must be a' or c'. This is exactly the condition that the slot ab belongs to $e_{t'}$. The proof for cb is identical.

If two targets t,u in $\mathcal{T}_*$ are comparable, say $t\mid u$, then u is a multiple of t lying in $(n/2,n]$. The next positive multiple of t after t is $2t>n$, so $u=t$. Hence distinct targets are pairwise incomparable, and playing an exact target scores only that target.

Finally, if e_t is live in the slot hypergraph, then none of b, ab, cb, or t has been played. The only other proper divisors of t are $1,a,c,ac$, and these are either not moves or already unavailable. There is no proper multiple of t on the board. Therefore no played number is comparable with t, so the actual move t is legal. This proves the comparison. □

Proposition A.5 (Activation-stage potential)**.** *During activation, let each unclaimed pair edge $e=(a,c)$ carry the current token set $B_e(t)$ of primes $b\in B_{a,c}$ for which the target acb has not yet been killed. Write $w_t(e):=|B_e(t)|$.*

Assume the safe-edge hypothesis of Proposition A.2 for the activation graph states generated below. Prolonger plays as follows. In each activation round, among pair edges with positive current token weight, he chooses a safe edge supplied by that hypothesis; if $e=(a,c)$ is chosen, he plays one live target acb with $b\in B_e(t)$. This scores one actual move, claims the pair edge e, captures the vertices a and c, and leaves the remaining live tokens on that secured pair for the residual phase.

Let S_t be the number of activation targets already played by Prolonger, and define

$$Q_t:=S_t+\sum_e\phi_t(e),$$

where

$$\phi_t(e) = \begin{cases} \frac{1}{8}w_t(e), & e \text{ unclaimed and neither endpoint captured,} \\ \frac{1}{4}w_t(e), & e \text{ unclaimed and one endpoint captured,} \\ \frac{1}{2}w_t(e), & e \text{ unclaimed and two endpoints captured,} \\ w_t(e), & e \text{ claimed,} \\ 0, & e \text{ deleted.} \end{cases}$$

Let E be the total number of target tokens deleted during activation by Shortener moves not modeled as graph vertex deletions a or graph edge deletions ac. At the end of activation,

$$Q_{\text{end}} \geq \frac{W_0}{8} - E.$$

Equivalently,

$$S_{\text{end}} + \sum_{e \text{ secured}} w_{\text{end}}(e) \geq \frac{W_0}{8} - E.$$

Proof. First ignore off-model token deletions. Conditional on the safe-edge hypothesis, the activation process is exactly the rank-2 graph game of Proposition A.2, with one harmless bookkeeping change: when Prolonger claims an edge e, he consumes one token from e and adds it immediately to the score S_t. If the coefficient of e before the move is $\gamma \in \{1/8, 1/4, 1/2\}$, the contribution of e changes from $\gamma w_t(e)$ to

$$1 + (w_t(e) - 1) = w_t(e),$$

exactly as if the edge had become claimed with full weight. All other incident edges change as in the graph game. Thus the scaled potential argument gives $Q_{\text{end}} \geq W_0/8$ when every harmful Shortener move is a graph deletion.

Now restore all actual Shortener moves. A token acb is deleted precisely when Shortener plays one of its still-available harmful divisors or the target itself. Deleting one live token can reduce Q_t by at most 1, since every coefficient appearing in the definition of $\phi_t(e)$ is at most 1. Therefore all off-model deletions together reduce the lower bound by at most E, giving

$$Q_{\text{end}} \geq W_0/8 - E.$$

At the end of activation, a positive-weight edge is either secured or deleted, so Q_{end} is exactly the activation score plus the residual live target weight on secured pairs. □

Proposition A.6 (Activation deletion budget)**.** *For fixed $0 < \delta < 1/4$, the off-model deletion count from Proposition A.5 satisfies*

$$E \ll \frac{Y^4}{\log^4 Y} = o\left(\frac{n(\log\log n)^2}{\log n}\right).$$

Proof. There is at most one activation round per small-prime pair, so the number of rounds is

$$R \leq \binom{|\mathcal{P}_Y|}{2} \ll \frac{Y^2}{\log^2 Y}.$$

We bound the number of live tokens an off-model Shortener move can delete.

If Shortener plays a large prime b, then for each pair (a, c) there is at most one token acb using that b. Hence the move deletes at most $O(Y^2/\log^2 Y)$ tokens. If Shortener plays a lateral semiprime pb with $p \leq Y$ and $b > Y$ prime, the killed targets have the form pcb with the third small prime varying, so there are at most $O(Y/\log Y)$ of them. If Shortener plays an exact target acb, exactly one token is deleted.

The worst of these bounds is the large-prime bound. Across R activation rounds,

$$E \ll R\frac{Y^2}{\log^2 Y} \ll \frac{Y^4}{\log^4 Y}.$$

Since $Y = n^\delta$,

$$\frac{Y^4/\log^4 Y}{n(\log\log n)^2/\log n} \ll_\delta \frac{n^{4\delta-1}}{(\log n)^3(\log\log n)^2} \to 0$$

because $4\delta - 1 < 0$. □

Proposition A.7 (Residual mass on secured pairs)**.** *At the end of activation, the secured pairs carry residual live target weight*

$$M := \sum_{e \text{ secured}} w_{\text{end}}(e) \gg_\delta \frac{n(\log\log n)^2}{\log n}.$$

Proof. By Proposition A.5,

$$S_{\text{end}} + M \geq \frac{W_0}{8} - E.$$

The activation score is at most the number of small-prime pairs:

$$S_{\text{end}} \leq R \ll \frac{Y^2}{\log^2 Y} = o\left(\frac{n(\log\log n)^2}{\log n}\right),$$

because $2\delta < 1$. The deletion budget E is negligible by Proposition A.6, while W_0 has the required lower bound by Proposition A.3. Subtracting the two negligible terms gives the claim. □

Proposition A.8 (Conditional T2 lower bound)**.** *Assume the safe-edge hypothesis of Proposition A.2 for the activation graph states and for the residual slot-hypergraph states generated by the construction above. For every fixed $0 < \delta < 1/4$ there is $c_\delta > 0$ such that*

$$L(n) \geq c_\delta \frac{n(\log\log n)^2}{\log n}$$

for all sufficiently large n.

Proof. After activation, restrict to the residual live targets lying over secured pairs. Their total weight is M, as in Proposition A.7. For each secured pair (a, c), the activation move made a, c, and ac unavailable. Therefore Proposition A.4 compares the remaining divisibility game to a scored rank-3 slot hypergraph whose total edge weight is M.

Under the residual safe-edge hypothesis, Proposition A.2 gives a strategy that forces scored residual target weight at least $M/8$ in this hypergraph model. The comparison proposition says that every such scored hypergraph move is a genuine actual game move, and that the actual game gives Prolonger no fewer legal options than the model. Hence the number of residual moves after activation is at least $M/8$. Since $M \gg_\delta n(\log\log n)^2/\log n$, the same lower bound holds for the total game length, conditional on the stated safe-edge hypothesis.

Choosing any fixed $\delta < 1/4$, for instance $\delta = 1/8$, gives an absolute constant $c > 0$ and proves Theorem 4.4. □

A.2. **Restricted carrier classes.** The next two propositions record unconditional upper bounds for two restricted families of Prolonger carrier moves. They are not used in the main upper bound, but they clarify which carrier configurations are not responsible for the remaining difficulty.

Proposition A.9 (Disjoint small-prime carriers)**.** *Fix $1/3 < \alpha < 1/2$ and put $y = n^\alpha$. Suppose that every composite Prolonger move has all prime factors at most y, and that the prime supports of distinct composite Prolonger moves are pairwise disjoint. Against this restricted class, Shortener has a strategy forcing*

$$L(n) = O_\alpha\left(\frac{n}{\log n}\right).$$

Proof. Let B be the set of primes which occur in composite Prolonger moves. By the disjoint-support hypothesis, each $p \in B$ belongs to a unique composite carrier $C(p)$.

Shortener uses the following priority list, always skipping moves which have already become illegal.

First, she plays every legal prime. This resolves every prime not belonging to a composite Prolonger carrier, and also every prime which Prolonger has not already made illegal by placing it inside a carrier.

Second, for every $p \in B$ whose carrier has at least two distinct prime factors, write $e(p) = v_p(C(p))$. If $p^{e(p)+1} \leq n$, Shortener plays $p^{e(p)+1}$. This move is legal when it is attempted: it is not comparable with $C(p)$, because it has a larger p-adic exponent while $C(p)$ contains another prime factor; it is not comparable with any other composite carrier by disjointness of supports; and previous priority moves either contain different prime supports or have smaller rank.

Third, for every pair $p, q \in B$ belonging to different carriers, Shortener plays the legal squarefree semiprime pq when possible. Such a move is incomparable with every carrier, since no carrier contains both p and q, and neither p nor q alone is available as a legal prime at this stage.

The number of moves in the three phases is

$$O(\pi(n)) + \pi(y) + \pi(y)^2 = O_\alpha\left(\frac{n}{\log n}\right),$$

because $2\alpha < 1$.

We now show that after these phases every future legal move has no prime factor at most y. Let x be a legal move divisible by a prime $p \leq y$.

If $p \notin B$, then phase one either played p or p was already played by Prolonger as a prime; in either case any multiple of p is illegal. Thus $p \in B$.

If x contains primes from two distinct carriers, then one of the phase-three semiprimes pq divides x, unless it was already illegal because an earlier played move dividing pq also divides x. Either way x is illegal.

Hence all prime factors of x which are at most y lie in a single carrier support. If the carrier is a pure prime power p^e, then any x divisible by p and supported only on this carrier is another power of p, and is comparable with p^e. If the carrier has at least two distinct prime factors, then either x divides the carrier, in which case x is illegal, or for some p it has exponent exceeding $v_p(C(p))$. In the latter case $p^{e(p)+1} \mid x$, so the phase-two move makes x illegal.

Thus all primes $\leq y$ are resolved. Any remaining legal integer has all prime factors greater than y. Since $y^3 > n$, it has at most two prime factors. The number of integers at most n with one or two prime factors, all exceeding $y = n^\alpha$, is $O_\alpha(n/\log n)$ by the prime number theorem and the standard semiprime estimate

$$\sum_{p>y} \pi(n/p) \ll_\alpha \frac{n}{\log n} \sum_{p>y} \frac{1}{p} = O_\alpha\left(\frac{n}{\log n}\right),$$

where the sum is over $p \leq n/y$. Adding the phase moves proves the claim. □

Proposition A.10 (Squarefree rank-three small-prime carriers)**.** *Fix $1/3 < \alpha < 1/2$ and put $y = n^\alpha$. Suppose that every composite Prolonger move is squarefree, has all prime factors at most y, and has support size at most 3. Against this restricted class, Shortener has a strategy forcing*

$$L(n) = O_\alpha\left(\frac{n}{\log n}\right).$$

Proof. Let B be the set of primes which occur in Prolonger's composite carriers. Shortener again uses a fixed priority list, skipping illegal moves.

First she plays every legal prime. Second she plays p^2 for every $p \in B$ when legal. Third she plays every legal squarefree semiprime pq with $p, q \in B$. Fourth she plays every legal squarefree triple pqr with $p, q, r \in B$.

The first phase has $O(n/\log n)$ moves, the second has at most $\pi(y)$ moves, and the third has at most $\pi(y)^2 = o(n/\log n)$ moves. It remains to bound the number of triples

$$T_\alpha(n) := \#\{p < q < r \le y : \ pqr \le n\}.$$

Split according to whether $pq \le n^{1-\alpha}$.

If $pq \le n^{1-\alpha}$, then there are $O(n^{1-\alpha}\log\log n/\log n)$ possible pairs $p < q$, and for each pair at most $\pi(y) = O(n^\alpha/\log n)$ choices of r. This contributes $o(n/\log n)$.

If $pq > n^{1-\alpha}$, then $q \le y = n^\alpha$ implies $p > n^{1-2\alpha}$. In this range,

$$\#\{r : \ q < r \le y, \ pqr \le n\} \ll_\alpha \frac{n}{pq\log n}.$$

Therefore the contribution is at most

$$\ll_\alpha \frac{n}{\log n} \sum_{n^{1-2\alpha} < p < q \le n^\alpha} \frac{1}{pq} = O_\alpha\left(\frac{n}{\log n}\right),$$

because the reciprocal prime sum over a fixed-power interval is $O_\alpha(1)$. Thus $T_\alpha(n) = O_\alpha(n/\log n)$.

We prove that the priority list resolves all primes at most y. After the prime and square phases, any remaining legal move supported on B must be squarefree. If it has support size 1, it is comparable with a played or blocked prime. If it has support size 2, then the corresponding semiprime was either played in phase three or was already illegal because an earlier played divisor blocks it. If it has support size 3, the same statement holds for phase four.

If a remaining legal move x had support size at least 4 inside B, choose three primes p, q, r from its support. The triple pqr was addressed in phase four. If it was played, then $pqr \mid x$ and x is illegal. If it was not legal when considered, then some earlier played move comparable with pqr must divide pqr; since all earlier non-prime moves in the priority list have support size at most 3 and are squarefree except the squares p^2, that played divisor also divides x. Again x is illegal. Thus no remaining legal move contains a prime at most y.

As in the proof of Proposition A.9, once all primes at most y are resolved, the residual legal set has size $O_\alpha(n/\log n)$. Combining this residual bound with the phase counts proves the proposition. □

Appendix B. Numerical Computations

This appendix records the numerical computations accompanying the main Bonferroni–4 constant. The computations certify the quoted constants and stress-test the finite combinatorial steps, but they are not substitutes for the analytic proof in Section 7.

B.1. **The Bonferroni–4 constant.** The main theorem uses

$$\frac{\mathcal{W}_4}{2} = 0.1897112 < 0.19.$$

A direct grid computation of the integrals

$$J_r = \frac{1}{r!}\int_{\substack{u_1+\cdots+u_r \le 1 \\ 0 < u_i \le 1}} \prod_{i=1}^{r} \rho(u_i)\, du_1 \cdots du_r$$

gives the convergence displayed in Table 1. The theorem only needs a strict margin below 0.19, and the finest-grid value provides a comfortable margin.

B.2. **Finite sanity checks.** Four finite checks were used to guard against implementation or formulation errors in the proof.

(1) *Odd-part injection.* Exhaustive verification at $n = 14$ over all 732 antichains found no collision under the odd-part map.
(2) *Monotone replacement.* Exact inclusion–exclusion counts at $n = 10^4, 10^5, 10^6$ for test prime pairs (q_j) and (p_j) with $q_j \le p_j$ confirmed $N(q_1, \ldots, q_K) \le N(p_1, \ldots, p_K)$.

Grid	J_1	J_2	J_3	J_4	$\mathcal{W}_4$	$\mathcal{W}_4/2$
2^{13}	0.78844185	0.18676003	0.02008041	0.00122154	0.37945930	0.18972965
2^{14}	0.78849100	0.18679383	0.02008737	0.00122220	0.37943767	0.18971883
2^{15}	0.78851725	0.18681153	0.02009097	0.00122255	0.37942586	0.18971293
2^{16}	0.78852127	0.18681405	0.02009149	0.00122260	0.37942388	0.18971194
2^{17}	0.78852704	0.18681777	0.02009220	0.00122266	0.37942118	0.18971059

TABLE 1. Grid convergence for the Bonferroni–4 truncation.

(3) *Squarefree divisor counts.* For $r \leq 4$, the relevant squarefree r-fold divisor counts match the scale $O_r(n(\log\log n)^{r-1}/\log n)$ used in the Bonferroni floor-error bound.
(4) *Controlled prime rounding.* Synthetic monotone comparison prefixes were rounded upward to increasing odd-prime prefixes at $n = 10^4, 10^5, 10^6, 10^8$; the resulting changes in the moment terms were small and consistent with the prime-rounding bridge.

Check	Recorded outcome
Odd-part injection	`(True, None, None, 732)` at $n = 14$
Monotone replacement	Verified at $n = 10^4, 10^5, 10^6$ for listed test prefixes
M_r scaling	Ratified for $r = 1, 2, 3, 4$ up to $n = 10^6$
Prime-rounding bridge	Tested on synthetic prefixes up to $n = 10^8$

TABLE 2. Finite checks for the combinatorial components of the upper-bound proof.

B.3. **Interpretation.** The numerical artifacts support three claims: the constant $\mathcal{W}_4/2 < 0.19$ is reproducible; the finite steps in Theorems 7.3 and 7.4 behave as expected in representative regimes; and the sharper limiting value near 0.18969 is numerically stable. The theorem in this paper is the finite Bonferroni–4 bound below 0.19, not a claim that the full alternating series has been analytically justified.

APPENDIX C. FORMALIZATION ARTIFACT MAP

This appendix records the status of the Lean artifacts corresponding to the paper's principal results. The table distinguishes between fully formalized finite or structural cores and prose components which use standard analytic number theory or conditional game-theoretic hypotheses.

The table should be read conservatively. A zero-sorry finite core is not being advertised as an end-to-end verification of every analytic asymptotic in the same theorem unless the table says so explicitly. Conversely, when the remaining obligations are standard analytic-number-theory inputs, the novelty of the combinatorial argument is already isolated in the formal artifact.

ARTIFACT AVAILABILITY

The public artifact repository for this paper is [**?**]. It contains the Lean formalizations, Aristotle submission files and outputs, numerical scripts, and supporting research notes used in the preparation of the present manuscript. Repository-relative paths quoted in Appendices B and C are relative to that artifact root.

ACKNOWLEDGMENTS

The author thanks Liam Price, Abhimanyu Adenwalla, StijnC, natso26, Xiao_Hu, and Desmond Weisenberg for the forum refinement chain leading from the first linear bound to the previous public record, and Thomas Bloom for maintaining `erdosproblems.com`. The author also wishes to record substantial gratitude for the research assistance provided by OpenAI's GPT-5.4 Pro, Anthropic's Claude Opus 4.7, and Google's Gemini 3.1 DeepThink, coordinated through a custom multi-round

Result	Lean status	Remaining prose or assumptions
Shield Reduction, Theorem 3.1	Zero-sorry Lean verification.	None beyond the finite definitions of the divisibility poset.
Exact 5/24 cover, Theorem 5.1	The structural cover and packing identities are zero-sorry Lean-verified.	The asymptotic floor-sum count $5n/24 + O(1)$ is an elementary prose calculation.
13/36 upper bound, Theorem 6.2	The odd-part compression and Bonferroni core are zero-sorry Lean-verified.	The paper-level exposition uses standard prime-counting estimates for the prime-prefix setup.
Polynomial shield-weight lower bound, Theorem 4.1	The combinatorial core, including the prime-antichain construction, log-budget argument, and exchange principle, has a Lean artifact.	Remaining sorries are Mertens' theorem, a Chebyshev-type prime-counting bound, a Chebyshev-ϑ asymptotic, and the final assembly using those analytic inputs.
5/16 upper bound, Theorem 6.3	The compression, sieve, and algebraic optimization core has a Lean artifact.	One remaining bundled placeholder covers the Chebyshev prime-prefix estimate and the game-tree wrapper realizing the prescribed prefix.
Conditional T2 lower bound, Theorem 4.4	The finite graph-potential core, scored hypergraph core, and local arithmetic embedding core are Lean-verified.	The safe-edge hypothesis is conditional, and the asymptotic activation wrapper remains in prose.
Main $0.19n$ upper bound, Theorem 7.5	The endgame reduction from sufficiently close first-four moment values to the strict inequality $L(n) < 0.19n$ is zero-sorry Lean-verified.	The local density theorem, monotone envelope and inversion, and prime-rounding bridge are the prose moment-convergence inputs proved in Section 7.

Table 3. Formalization status of the principal results.

research harness built by the author. GPT-5.4 Pro produced the majority of the exploratory mathematical output used in this project, with Claude and Gemini providing extensive auditing, synthesis, and drafting support. A separate methodological paper and public-facing writeup will describe that workflow in detail.

Declaration of Generative AI and AI-assisted Technologies in the Research and Writing Process

During the preparation of this work, the author used generative AI systems including OpenAI GPT-5.4 Pro, Anthropic Claude Opus 4.7, and Google Gemini 3.1 DeepThink for exploratory discussion, adversarial proof auditing, code and Lean scaffolding, numerical cross-checking, artifact organization, and drafting assistance. After using these tools, the author reviewed, corrected, and rewrote the material as needed, verified references and formal artifacts against the repository, and takes full responsibility for the claims, proofs, citations, and presentation in the final manuscript.

References

[BHW16] Csaba Biró, Paul Horn, and D. Jacob Wildstrom, *An upper bound on the extremal version of hajnal's triangle-free game*, Discrete Applied Mathematics **198** (2016), 20–28.

[CE90] Peter J. Cameron and Paul Erdős, *On the number of sets of integers with various properties*, Number Theory (Banff, AB, 1988), de Gruyter, Berlin, 1990, pp. 61–79.

[Erd92] Paul Erdős, *Some of my forgotten problems in number theory*, Hardy-Ramanujan Journal **15** (1992), 34–50.

[FS92] Zoltán Füredi and Ákos Seress, *On hajnal's triangle-free game*, Graphs and Combinatorics **8** (1992), no. 1, 75–79.

[Kuc24] Petr Kucheriaviy, *Erdős inequality for primitive sets*, 2024.

[Lic23] Jared Duker Lichtman, *A proof of the erdős primitive set conjecture*, Forum of Mathematics, Pi **11** (2023), e18.

[MP10] Greg Martin and Carl Pomerance, *Primitive sets with large counting functions*, 2010.

Independent Researcher
Email address: sensho@sensho.xyz