



Review Article

Sleep in children with autistic spectrum disorder

Flavia Cortesi^{a,*}, Flavia Giannotti^a, Anna Ivanenko^b, Kyle Johnson^c^a Center of Pediatric Sleep Disorders, Department of Developmental Neurology and Psychiatry, University of Rome "La Sapienza" Italy, Via dei Sabelli 108, 00185 Rome, Italy^b Division of Child and Adolescent Psychiatry, Children's Memorial Hospital, Chicago, IL, USA^c Division of Child and Adolescent Psychiatry, Oregon Health and Science University, USA

ARTICLE INFO

Article history:

Received 2 September 2009

Received in revised form 31 December 2009

Accepted 7 January 2010

Keywords:

Autism

Sleep

Sleep disorders

Autistic regression

Treatment

Asperger syndrome

Melatonin

ABSTRACT

Children and adolescents with autistic spectrum disorders (ASD) suffer from sleep problems, particularly insomnia, at a higher rate than typically developing children, ranging from 40% to 80%. Sleep problems in ASD might occur as a result of complex interactions between biological, psychological, social/environmental, and family factors, including child rearing practices that are not conducive to good sleep. Interestingly, children with a history of developmental regression have a more disturbed sleep pattern than children without regression. Even though regulation of sleep in children with ASD is still poorly understood, circadian abnormalities in autism might be the result of genetic abnormalities related to melatonin synthesis and melatonin's role in modulating synaptic transmission. Recently a bifurcation of the sleep/wake cycle with increased sensitivity to external noise and short sleep duration causing irregular sleep onset and wake up times has been suggested. Identifying and treating sleep disorders may result not only in improved sleep, but also impact favorably on daytime behavior and family functioning. Several studies have also demonstrated effectiveness of behavioral interventions for sleep onset and maintenance problems in these populations. When behavioral interventions are not effective or lead only to a partial response, pharmacological treatment options should be considered. Studies of melatonin use in children with ASD provide evidence for its effectiveness and safety in the long run. The clinician assessing a child with an ASD should screen carefully for sleep disorders and make referrals as indicated.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Autism spectrum disorders (ASD) are life-long neurodevelopmental disorders that affect all areas of child development, characterized by markedly abnormal or impaired social interaction and communication, restricted interests, and stereotypical behaviors [1]. Numerous research studies indicate that children with ASD experience more sleep problems than the general population. The main objective of this review is to present research data on the sleep problems in children with ASD with emphasis on their prevalence, clinical characteristics, sleep architecture, potential etiology, influence on daytime behavior, assessment and treatment.

2. Prevalence of sleep problems in ASD

Sleep problems are highly prevalent in children with ASD ranging from 40% to 80% [2–12]. A recent study showed that sleep problems rank as one of the most common concurrent clinical disorders

among children with ASD [13]. Since the majority of research studies include children with ASD and intellectual deficits, the prevalence rates of sleep problems need to be evaluated in the context of comorbid intellectual disability. Some reports revealed an increased proportion of severely developmentally delayed children with ASD who had sleep problems [14,15]. Given this association between intellectual deficits and sleep problems, attempts have been made to control for mental retardation as a confounding variable. In a study of highly functioning children with ASD, sleep problems appeared to be relatively specific to children with ASD compared to both typically developing children and children with intellectual disabilities without autism [16]. Based on parental reports, Krakowiak and colleagues [17] found that 53% of children (2–5 years of age) with ASD had at least one frequently experienced sleep problem compared to 46% of children with non-ASD developmental delays and 32% of typically developing (TD) children.

These studies and others demonstrate that children with ASD are particularly vulnerable to sleep difficulties [16–20]. However, the majority of these studies are based on parental reports. A recent study, using a multimethod approach, found that pre-school children with autism and developmentally delayed children without autism showed more sleep problems as reported by their

* Corresponding author. Tel.: +39 0644712219; fax: +39 4957857.

E-mail address: flavia.cortesi@uniroma1.it (F. Cortesi).

parents than TD children, but these problems were not directly supported by more objective actigraphic data. Moreover, they found that parents of children with ASD slightly over reported sleep problems. Thus, the authors suggested that objective and subjective measures of sleep problems in pre-school-age children may produce different results [21].

Sleep problems often last for many years and sometimes continue into adulthood [22,23]. In one study, 63% of children with ASD and sleep problems experienced persistence of sleep difficulties over time [24]. However, Goodlin-Jones and colleagues recently used objective and subjective measures to assess sleep in pre-school children with autism, children with developmental delays only, and typically developing children; objectively measured sleep problems were rarely persistent over a 6-month period of time [25].

3. Clinical characteristics of sleep in children with ASD

Sleep onset and maintenance problems resulting in reduced sleep duration are the most common concerns expressed by the parents of children with ASD. Several studies have reported settling problems with some children with ASD taking more than an hour to fall asleep [10,17,26]. According to parental reports, night wakings also commonly occur in children with ASD. Periods of nocturnal awakening lasting for up to 2–3 h have been reported when the child may simply laugh, talk, scream, or get up and play with toys or various objects in the room [9,15,26,27]. As noted by Richdale and Schreck [20], these night wakings are typical of children with autism. Studies utilizing actigraphy largely confirm increased sleep latency, frequent night waking, and shorter sleep duration [21,24,28]. It is worth noting that sleep onset and night-waking problems are often associated with poor sleep hygiene or maladaptive sleep associations [24]. Thus, as in the general population, behavioral insomnia is frequently experienced by children with ASD.

Although some sleep problems may have a strong behavioral component, circadian rhythm disturbances have also been reported in children with ASD [24]. Irregular sleep–wake patterns, free running sleep/wake rhythms, sleep onset delay and early morning awakening have been reported in children with autism [29–31]. Infants with autism tend to show early waking in the morning or later sleep onset in the evening [29]. Inanuma [32] reported significant day to day variability in sleep onset or waking times in children with ASD, especially in those less than 4 years of age.

Regulation of sleep in children with ASD is still poorly understood. In their recent study, Matsuura and colleagues [33] suggested a bifurcation of the sleep/wake cycle with increased sensitivity to external noise and short sleep duration causing irregular sleep onset and wake up times. The authors speculated that children with ASD have an altered balance between homeostatic and circadian forces leading to a heightened sensitivity of the sleep/wake cycle to environmental stimuli. They also concluded that sleep regulation tended to improve with age. The latter conclusion is supported by a study showing that more than 70% of children with ASD experience delays in circadian rhythm development by at least 5 months compared to typically developing children [31]. Interestingly, in a recent study comparing the sleep patterns and problems of pre-school age children with carefully diagnosed autism, developmental delay and TD, Goodlin-Jones and colleagues [21] found that children with autism sleep for shorter periods of time during a 24-h day. This suggests that sleep patterns in children with autism are organized differently than other children.

Moreover, seasonal changes in sleep patterns have been described among children with ASD. In one case, the sleep problems of an adolescent male with autism decreased from January to June and disappeared by July and August. His sleep disturbances re-

curred in October and were sustained through November and December [30]. In a recent study by Giannotti and colleagues [15], more than 10% of children with autism had sleep problems that varied by season. Sleep problems rapidly increased during the fall and spring, particularly in those individuals with a history of developmental regression.

Studies investigating variables potentially related to sleep problems (age, IQ, severity of autistic symptoms) in children with autism have yielded mixed results. Regarding the relationship between IQ and sleep problems, Richdale and Prior [18] reported more frequent sleep problems in children with moderate to severe intellectual disability compared to autistic children with mild degrees of disability. Another study reported only more night wakings [34] in children with mental retardation compared to children with autism and no retardation. In contrast, the majority of studies suggested that sleep problems can occur at all IQ levels, including in those who do not have mental retardation [2,16,17,35,36]. Regarding the relationship between age and sleep problems in children with autism, several studies have found increasing age to be associated with decreasing sleep problems [8,17,18]. In contrast, Honomichl [6] found that school aged children were reported to present more sleep problems than pre-schoolers. However, other studies reported no significant relationship between age and sleep problems in children with ASD [24,34–36]. Finally, regarding the association between sleep problems and severity of autistic symptomatology, some studies have reported a significant positive association between autism severity and parent reported sleep problems [27,36,37]. These findings suggest that sleep problems are part of the autism symptom complex and increase with increasing autism severity.

In considering prevalence and clinical characteristics of sleep problems in ASD, it should be noted that the majority of studies have been carried out on heterogeneous samples including children of various ages, intellectual levels, and types of autistic spectrum disorders. There have not been detailed analyses of associations between sleep patterns and particular medical conditions, which may themselves alter sleep or their treatments. Few studies have been carried out on more homogeneous groups.

4. Sleep architecture in children with ASD

Several PSG studies have confirmed the presence of disrupted sleep architecture in children with ASD [14,38–41]. PSG abnormalities include increased REM density [39], reduction of REM sleep, and an increase of undifferentiated sleep with a clear abundance of spindle activity not only during stage 2 sleep but also during SWS and REM sleep [38]. Longer sleep latency, increased night wakings, lower sleep efficiency, increased stage 1 sleep and decreased SWS as well as decreased density of spindle activity [41] have been described in children with ASD. One study indicated reduced total sleep time and shorter REM latency [14]. In this study, Miano and colleagues investigated sleep NREM microstructure in children with ASD and intellectual disability using cyclic alternating pattern (CAP) analysis [14,42]. They found a lower CAP rate associated with SWS sleep suggesting a subtle alteration of the arousal level during NREM. Whether these microstructural alterations of sleep relate to autism or intellectual disability is not known. A recent PSG investigation found that among children with autism, those with a history of autistic regression had longer sleep latency, less sleep efficiency, increased wake after sleep onset and longer REM latency than those without regression. Moreover, children with autistic regression showed lower CAP rates and A1 index during light sleep than children without history of regression and than TD children [43]. In contrast, a PSG study of a selected group of children with autism and sleep problems but without mental

retardation found lower sleep efficiency, prolonged sleep latency, decreased REM sleep and increased stages 3 and 4 sleep compared to children with ASD without sleep problems and typically developing children [10]. Interestingly, a PSG study on sleep architecture and NREM alteration found that children with Asperger syndrome showed lower CAP rates in stages 1 and 2, but not in SWS, and an increased percentage of A1 and decreased percentage of A2 compared with TD children. Moreover, with respect to children with ASD, Asperger subjects showed an increased CAP rate in SWS and decreased rate in sleep stage 2 suggesting that these different microstructural alterations of sleep might be related to the degree of mental ability [44].

Inconsistent results on sleep architecture in ASD can be explained by the wide range of patient population and methodology used across different studies. Future research in this area should emphasize more stringent diagnostic and selection criteria for subjects and more unified definitions of sleep disorders.

5. Pathophysiological mechanisms of sleep disorders in ASD

Richdale and Schreck [20] proposed that sleep problems in ASD might occur as a result of complex interactions between biological, psychological, social/environmental and family factors, including child rearing practices that are not conducive to good sleep. Any one or combination of these factors might contribute to sleep problems in children with ASD.

6. Neurobiological abnormalities

Intrinsic biological or genetic abnormalities that alter neuronal pathways in the brain might contribute to sleep problems in ASD. Most autistic children exhibit abnormal EEG and sleep/wake patterns, suggesting that the balance of excitation–inhibition is disrupted. Proposed hypotheses of sleep dysregulation in autism include abnormalities in the hypothalamic–pituitary–adrenal axis regulating circadian rhythms and alteration in hormone/neurotransmitter (melatonin/serotonin) production [9].

Several neurotransmitter systems, including gamma-aminobutyric acid (GABA), serotonin and melatonin, are implicated in promoting sleep and establishing a regular sleep/wake cycle. Aberration in these systems may be responsible for sleep disturbance in ASD [45]. Failure to develop a circadian cycle due to failure to entrain to the day/night cycle has been suggested in ASD [46]. This implicates involvement of five HT neurons among ascending brainstem monoaminergic neurons that modulate development of the circadian sleep/wake cycle [47].

The preoptic area of the hypothalamus is a major sleep-promoting system that uses GABA as a neurotransmitter. Sleep-active neurons in the preoptic area project to brainstem regions that contain neurons involved in arousal from sleep; inhibiting these regions in turn promotes sleep. These regions include pedunculopontine and laterodorsal tegmental nuclei (PPT/LDT), locus coeruleus, and dorsal raphe [48]. In autism, GABAergic interneurons migration and maturation could be affected [45]. A region of genetic susceptibility has been identified on chromosome 15q that contains GABA-related genes [49]. The expression of the autism susceptibility genes may affect sleep regulation by interfering with the normal inhibitory function of GABA via the preoptic area neurons. Therefore, the reduction of sleep time might be a consequence of dysfunction of the brainstem aminergic pathways [39].

7. Circadian abnormalities

Bourgeron [50] suggested that clock genes associated with low melatonin may be involved in abnormal circadian timing in aut-

ism. He hypothesized that circadian abnormalities in autism might be the result of genetic abnormalities related to melatonin synthesis and melatonin's role in modulating synaptic transmission. An association between clock gene polymorphisms and autism has been reported [51]. These findings support the hypothesis of circadian abnormalities underlying sleep disturbances in autism.

At least four independent studies have demonstrated abnormal melatonin regulation in individuals with ASD compared with controls, including elevated daytime melatonin and significantly lower nocturnal melatonin [47,52–54]. Among those, one study shows significantly lower excretion rates of urinary 6-sulphatoxymelatonin (the major metabolite of melatonin) in children and adolescents with autism compared to age-and-gender matched controls, with more marked differences in pre-pubertal subjects [47]. Interestingly, the low level of melatonin correlated with severity of autistic symptomatology. In a more recent study, Melke [55] found abnormal melatonin levels in the unaffected parents of children with autism, suggesting a possible genetic origin. In this study, the melatonin deficit was associated with low *ASMT* activity. The *ASMT* gene, located on pseudo-autosomal region 1 of the sex chromosomes and deleted in several individuals with ASD, encodes the last enzyme involved in melatonin synthesis, indicating that variations in this gene could explain the melatonin deficit.

Abnormalities in serotonin synthesis have been demonstrated in children with ASD, including an increased synthesis and asymmetry in serotonin production [56,57]. Recently, Makkonen [58] found reduced serotonin transporter binding capacity in children with ASD. It has been hypothesized that a primary deficit of melatonin synthesis could indirectly lead to accumulation of upstream substrates (*N*-acetylserotonin) and serotonin. Thus, a single defect within the melatonin synthesis pathway could lead to both low melatonin and high serotonin levels in autism [50]. Taken together, these data indicate that the deficit in melatonin may cause abnormal sleep/wake cycles in individuals with ASD. Given these low melatonin levels, exogenous melatonin administration may be therapeutic.

8. Psychological or behavioral factors

Children with ASD have a high rate of psychiatric comorbidities that may interfere with the regulation of sleep. Anxiety disorders, mood disorders, and attention-deficit/hyperactivity disorder frequently occur in patients with ASD [59], resulting in complicated presentations of sleep disturbances [8,9,37,60].

Behavioral sleep disorders such as bedtime resistance, sleep-onset association and limit-setting types of behavioral insomnia are common among children with ASD and can be successfully addressed with behavioral interventions [61]. Maladaptive bedtime routines may lead to settling difficulties at bedtime with inconsistent sleep onset time. The adherence to positive and consistent bedtime routines is associated with a reduction in sleep problems in children with ASD [61,62].

Children's sleep problems can significantly impact the quality of sleep of other family members. Sleep problems were reported by almost 70% of 210 Hong Kong families with a young child with ASD. Parental stress predicted the severity of sleep problems in these children [63]. Sleep maintenance difficulties in children with ASD have a significant impact on parental sleep [64]. A recent study by Meltzer [65] reported poorer sleep quality and shorter total sleep time among parents of children with autism compared to parents of typically developing children.

9. Developmental regression and sleep problems

In a minority of patients, symptoms of autism become apparent after a period of normal or mildly delayed development followed

by the loss of previously acquired skills [66,67]. Autistic regression typically occurs between 1 and 3 years of age, with a peak at around 24 months. This regression occurs in about 15–40% of children with autism [66–70]. Even though the mechanisms underlying autistic regression are still unknown, it has been postulated that rapid synaptic growth and pruning during the second year of life is impaired in autism due to genetic influences [71] resulting in a particular developmental phenotype.

Interestingly, Giannotti and co-authors found that over 50% of children with autism had at least one sleep problem, with a peak onset during the second year of life which coincides with the time of autistic regression [72]. This temporal overlap might be coincidental, but could also be interpreted as the expression of a biologically determined mechanism which influences a particular developmental trajectory during a critical phase of child development. More recent data from parental surveys and nocturnal polysomnography suggest that children with ASD and a history of regression have a more disturbed sleep pattern than children without regression [15]. It is noteworthy that approximately one-third of children with autism develop epilepsy [73] with prevalence estimates between 12% and 39% [74,75]. Epileptiform EEG abnormalities in children with autism have been reported to range between 8% and 15% [68] and 60% [76,77]. Although epilepsy cannot be considered a causal factor, in some cases it arises in the context of abnormal brain systems or networks which seems to be implicated in the autistic disorder as well.

The coexistence of disrupted sleep patterns and significantly higher presence of epilepsy in regressed children suggests a disruption of neuronal circuitry, which might be a reflection of the underlying pathology. In addition, seizures and sleep problems could possibly be caused by the underlying pathological process and have a compounding effect on the evolving neurocognitive dysfunction in autism. Although the biological basis and possible causal relationships of these associations remain unknown, this may point to different subgroups of patients with autism.

10. Impact of sleep problems on daytime behavior in children with ASD

An important reason, if not the most important, for intervening when children with ASD have sleep problems is to improve daytime functioning. Parents of poorly sleeping children with ASD routinely report worse daytime behavior following nights of fragmented or insufficient sleep. Although several studies have pointed out the negative effect of sleep disorders on daytime behavior in children with intellectual disabilities [22,78–80], few studies have evaluated the effect of sleep disorders exclusively in autism. Schreck and colleagues [27] demonstrated that short sleep duration was associated with social skills deficits, stereotypic behavior, and increased overall autism scores. Children with autism described by their parents as “poor sleepers” as opposed to “good sleepers,” had more difficulties with reciprocal social interaction on the Autism Diagnostic Observation Schedule and higher scores related to affective problems on the Child Behavior Checklist [10]. In a case report, a girl with ASD and obstructive sleep apnea showed improvement in daytime behaviors measured before and after treatment with adenotonsillectomy [81]. Her ability to interact socially improved as did her ability to focus and she demonstrated less repetitive behavior and auditory sensitivity.

11. Non-pharmacological treatment

Sleep hygiene and behavioral therapy have been shown to be effective interventions for both typically developing children and for children with ASD. Several studies have demonstrated effective-

ness of behavioral interventions for sleep onset and maintenance problems in children with ASD [61,62]. Richdale and Wiggs [82] have written a comprehensive review of behavioral interventions in children with developmental disabilities and autism. Basic principles of sleep hygiene include the selection of an appropriate bedtime, establishment of a positive consistent bedtime routine, minimizing television watching and playing computer or video games, and reducing emotional/behavioral stimulation at night. It is equally important to establish a consistent bedtime and wake time to promote a regular sleep/wake cycle.

Children with ASD frequently have hypersensitivity to environmental stimuli, including noises or tactile sensitivity to bedclothes or blankets. Addressing sensory hypersensitivity especially associated with the sleep environment may help improve quality of sleep. For example, parents have reported weighted blankets being helpful in children with tactile sensitivities. However, there are no evidence-based research studies supporting their consistent use for children with ASD. A parental survey conducted by Williams and colleagues [83] indicated that wrapping a child with ASD in a blanket helped with sleep onset.

As a general rule the sleep environment should provide the child with the most comfortable temperature, preferred texture of the pajamas and bedding, and optimal control of noise and light. Some children may benefit from white noise machines or other soothing sounds during sleep.

Other behavioral interventions like graduated extinction, scheduled nocturnal awakenings, and bedroom passes can be utilized to set appropriate behavioral limits at bedtime and to promote the development of more adaptive self-soothing skills [84]. Limiting naps may help to improve sleep continuity and reduce sleep onset latency in children who tend to nap extensively during the day. Light therapy is effective in advancing or delaying the sleep phase in patients with circadian sleep disorders and can be considered for children with ASD who present with circadian rhythms dysfunction.

12. Pharmacological treatment

When behavioral interventions are not effective or lead only to a partial response, pharmacological treatment options should be considered in as an adjuvant to ongoing behavioral therapy for sleep disorders. When considering pharmacological options, it is important to realize that there are no medications approved by the Food and Drug Administration (FDA) for the treatment of pediatric insomnia. Selection of a sleep-promoting agent should be based on the individual patient's characteristics, pharmacological properties of the drug, previous history of child's response to sedative/hypnotics, psychiatric and medical comorbidities and drug-drug interactions.

Melatonin has been more extensively studied in children with insomnia than any other sleep-promoting medication. Initial research in children with neurodevelopmental disorders demonstrated reductions in sleep latency and improvements in total sleep time and sleep efficiency when melatonin was administered close to bedtime [85–88]. A more recent large retrospective study that included over 100 children with ASD documented minimal adverse effects from melatonin with improved quality of sleep in 85% of participants [89]. Melatonin was shown to be effective in a dose range from 3 to 6 mg in children with ASD [90–92]. Controlled-release melatonin was found to be superior to behavioral intervention in a controlled study of children with ASD using actigraphy. Melatonin in this study was highly effective in reducing sleep onset latency, night-to-night variability in bedtime, waking after sleep onset, and in increasing sleep duration [92]. There is evidence to support use of extended release formulations of melatonin for

sleep maintenance insomnia in children [93]. Although, the studies of melatonin use in children with ASD provide evidence for its effectiveness and safety they frequently lack controls and are small in sample sizes.

There is limited research on the use of prescription medications with sedative properties for children with autism. One study demonstrated the effectiveness of niaprazine, a histamine H1-receptor antagonist with sedative properties which is available in Europe, to treat insomnia in this population [94]. For more detailed information on the pharmacological treatment of pediatric insomnia, please refer to a recent review by Owens [95].

Successful pharmacological treatment of comorbid psychiatric disorders like ADHD, mood and/or anxiety disorders as well as neurological conditions like epilepsy and movement disorders may help to alleviate symptoms of insomnia and markedly improve outcome.

13. Conclusion

Children and adolescents with an ASD are at substantial risk for experiencing sleep problems, particularly insomnia. The clinician assessing a child with an ASD should screen carefully for sleep disorders and make referrals as indicated. Identifying and treating sleep disorders may result not only in improved sleep, but also impact favorably on daytime behavior and family functioning. Fortunately, there are a variety of treatments available, including behavioral interventions and pharmacotherapy. It is imperative to establish standards of care for the evaluation and treatment of sleep disorders in this population. Additionally, clinical trials targeting insomnia are needed using interventions such as behavioral, pharmacological, and light therapy.

References

- [1] American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th ed. Text Revision. Washington, DC: American Psychiatric Association; 2000.
- [2] Richdale AL. Sleep problems in autism: prevalence, cause and intervention. *Dev Med Child Neurol* 1999;41(1):60–6.
- [3] Hering E, Epstein R, Elroy S, Iancu DR, Zelnik N. Sleep patterns in autistic children. *J Autism Dev Disord* 1999;29:143–7.
- [4] Schreck KA, Mulick JA. Parental report of sleep problems in children with autism. *J Autism Dev Disord* 2000;30(2):127–35.
- [5] Wiggs L. Sleep problems in children with developmental disorders. *J R Soc Med* 2001;94:177–9.
- [6] Honomichi RD, Goodlin-Jones BL, Burnham M, Gaylor E, Anders TF. Sleep patterns of children with pervasive developmental disorders. *J Autism Dev Disord* 2002;32(6):553–61.
- [7] Polimeni MA, Richdale AL, Francis AJP. A survey of sleep problems in autism, Asperger's disorder and typically developing children. *J Intellect Disabil Res* 2005;49(4):260–8.
- [8] Liu X, Hubbard JA, Fabes RA, Adam JB. Sleep disturbances and correlates of children with autism spectrum disorders. *Child Psychiatry Hum Dev* 2006;37:179–91.
- [9] Malow BA, McGrew S. Sleep and autism spectrum disorders. In: Tuchman R, Rapin I, editors. *Autism: a neurological disorders of early brain development*. London: Mac Keith Press; 2006. p. 188–201.
- [10] Malow BA, Marzec ML, McGrew SG, Wang L, Henderson LM, Stone WL. Characterizing sleep in children with autism spectrum disorders: a multidimensional approach. *Sleep* 2006;29(12):1563–71.
- [11] Souders M, Mason T, Valadares MS, Bucan M, Levy SE, Mandell D, et al. Sleep behaviors and sleep quality in children with autism spectrum disorders. *Sleep* 2009;32(12):1566–78.
- [12] Paavonen EJ, Vehkalahti K, Vanhala R, Von Wendt L, Nieminen-von Wendt T, Aronen ET. Sleep in children with Asperger syndrome. *J Autism Dev Disord* 2008;38:41–51.
- [13] Ming X, Brimacombe M, Chaaban J, Zimmerman-Bier B, Wagner GC. Autism spectrum disorders: concurrent clinical disorders. *J Child Neurol* 2008;23(1):6–13.
- [14] Miano S, Bruni O, Elia M, Trovato A, Smerieri A, Verrillo E. Sleep in children with autistic spectrum disorder: a questionnaire and polysomnographic study. *Sleep Med* 2007;9:64–70.
- [15] Giannotti F, Cortesi F, Cerquiglini A, Miraglia D, Vagnoni C, Sebastiani T, et al. An investigation of sleep characteristics, EEG abnormalities and epilepsy of developmentally regressed and non-regressed autistic children. *J Autism Dev Disord* 2008;38(10):1888–97.
- [16] Couturier JL, Speechley KN, Steele M, Norman R, Stringer B, Nicolson R. Parental perception of sleep problems in children of normal intelligence with pervasive developmental disorders: prevalence, severity, and pattern. *J Am Acad Child Adolesc Psychiatry* 2005;44(8):815–22.
- [17] Krakowiak P, Goodlin-Jones B, Hertz-Picciotto I, Croen LA, Hansen RL. Sleep problems in children with autism spectrum disorders, developmental delays, and typical development: a population-based study. *J Sleep Res* 2008;17:197–206.
- [18] Richdale AL, Prior MR. The sleep/wake rhythm in children with autism. *Eur Child Adolesc Psychiatry* 1995;4(3):175–86.
- [19] Bradley EA, Summers JA, Wood HL, Bryson SE. Comparing rates of psychiatric and behavior disorders in adolescents and young adults with severe intellectual disability with and without autism. *J Autism Dev Disord* 2004;34(2):151–61.
- [20] Richdale AL, Schreck KA. Sleep problems in autism spectrum disorders: prevalence, nature and possible biopsychosocial etiologies. *Sleep Med Rev* 2009; doi:10.1016/j.smrv.2009.02.003.
- [21] Goodlin-Jones BL, Tang K, Liu J, Anders TF. Sleep patterns in preschool children with autism, developmental delay, and typical development. *J Am Acad Child Adolesc Psychiatry* 2008;47(8):932–40.
- [22] Robinson AM, Richdale AL. Sleep problems in children with an intellectual disability: parental perception of sleep problems, and views of treatment effectiveness. *Child Care Health Dev* 2004;30(2):139–50.
- [23] Bramble D. Consumer opinion concerning the treatment of a common sleep problem. *Child Care Health Dev* 1996;22:355–66.
- [24] Wiggs L, Stores G. Sleep patterns and sleep disorders in children with autistic spectrum disorders: insights using parent report and actigraphy. *Dev Med Child Neurol* 2004;46:372–80.
- [25] Goodlin-Jones B, Schwichtenberg AJ, Iosif AM, Tang K, Liu J, Anders TF. Six month persistence of sleep problems in young children with autism, developmental delay, and typical development. *J Am Acad Child Adolesc Psychiatry* 2009;48(8):847–54.
- [26] Richdale A. Sleep disorders in autism and Asperger's syndrome. In: Stores G, Wiggs L, editors. *Sleep disturbances in disorders of development: its significance and management*. London: Mac Keith Press; 2001. p. 181–91.
- [27] Schreck KA, Mulick JA, Smith A. Sleep problems as possible predictors of intensified symptoms of autism. *Res Dev Disabil* 2004;24:57–66.
- [28] Oyane NM, Bjorvatn B. Sleep disturbances in adolescents and young adults with autism and Asperger syndrome. *Autism* 2005;9(1):83–94.
- [29] Takase M, Taira M, Sasaki H. Sleep–wake rhythm of autistic children. *Psychiatry Clin Neurosci* 1998;52:181–2.
- [30] Hayashi E. Seasonal changes in sleep and behavioral problems in a pubescent case with autism. *Psychiatry Clin Neurosci* 2001;55:223–4.
- [31] Segawa M. Epochs of development of the sleep–wake cycle reflect the modulation of the higher cortical function particular for each epoch. *Sleep Biol Rhythms* 2006;4:4–15.
- [32] Inanuma K. Sleep–wake patterns in autistic children. *Jpn J Child Adolesc Psychiatry* 1984;25(4):205–17.
- [33] Matsuura H, Tateno K, Aou S. Dynamical properties of the two-process model for sleep–wake cycles in infantile autism. *Cogn Neurodyn* 2008;2:221–8.
- [34] Gail Williams P, Sears LL, Allard A. Sleep problems in children with autism. *J Sleep Res* 2004;13:265–8.
- [35] Patzold LM, Richdale AL, Tonge BJ. An investigation into sleep characteristics of children with autism and Asperger's disorder. *J Paediatr Child Health* 1998;34:528–33.
- [36] Dickerson Mayes S, Calhoun SL. Variables related to sleep problems in children with autism. *Res Autism Spectr Dis* 2009;3:931–41.
- [37] DeVincent CJ, Gadow KD, Delosh D, Geller L. Sleep disturbance and its relation to DSM-IV psychiatric symptoms in preschool-age children with pervasive developmental disorder and community controls. *J Child Neurol* 2007;22:161–9.
- [38] Diomedes M, Curatolo P, Scalise A, Placidi F, Caretto F, Gigli GL. Sleep abnormalities in mentally retarded autistic subjects: Down's syndrome with mental retardation and normal subjects. *Brain Dev* 1999;21:548–53.
- [39] Elia M, Ferri R, Musumeci SA, Del Gracco S, Bottitta M, Scuderì C, et al. Sleep in subjects with autistic disorders: a neurophysiological and psychological study. *Brain Dev* 2000;22:88–92.
- [40] Thirumalai SS, Shubin RA, Robinson R. Rapid eye movement sleep behavior disorder in children with autism. *J Child Neurol* 2002;17:173–8.
- [41] Limoges E, Mottron L, Bolduc C, Berthiaume C, Godbout R. Atypical sleep architecture and the autism phenotype. *Brain* 2005;128:1049–61.
- [42] Terzano MG, Mancia D, Salati MR, Costani G, Decembrino A, Parrino L. The cyclic alternating pattern as a physiologic component of normal NREM sleep. *Sleep* 1985;8:137–45.
- [43] Giannotti F, Cortesi F, Vagnoni C, Sebastiani T, Cerquiglini A, Bernabei P. Polysomnographic characteristics in regressed and nonregressed autistic children. *Sleep* 2007;30:A73.
- [44] Bruni O, Ferri R, Vittori E, Novelli L, Vignati M, Porfirio MC, et al. Sleep architecture and NREM alterations in children and adolescents with Asperger syndrome. *Sleep* 2007;30(11):1577–85.
- [45] Levitt P, Eagleson KL, Powell EM. Regulation of neocortical interneuron development and the implications for neurodevelopmental disorders. *Trends Neurosci* 2004;27:400–6.
- [46] Segawa M. Circadian rhythm in early infantile autism. *Shinkei Kenkyu No Shimpo* 1985;29:140–53.

- [47] Tordjman S, Anderson GM, Pichard N, Charbuy H, Touitou Y. Nocturnal excretion of 6-sulphatoxymelatonin in children and adolescents with autistic disorder. *Biol Psychiatry* 2005;57:134–8.
- [48] Saper CB, Chou T, Scammell TE. The sleep switch: hypothalamic control of sleep and wakefulness. *Trends Neurosci* 2001;24:726–31.
- [49] McCauley JL, Olson LM, Delahanty R, Amin T, Nurmi EL, Organ EL, et al. A linkage disequilibrium map of the 1-Mb 15q12 GABAA receptor subunit cluster and association to autism. *Am J Med Genet* 2004;131:51–9.
- [50] Bougeron T. The possible interplay of synaptic and clock genes in autism spectrum disorders. *Cold Spring Harb Symp Biol* 2007;LXXII:1–10.
- [51] Nicholas B, Rudrasingham V, Nash S, Kirov G, Owen MJ, Wimpory DC. Association of *per1* and *Npas2* with autistic disorder: support for the clock genes/social timing hypothesis. *Mol Psychiatry* 2007;12:581–92.
- [52] Ritvo ER, Ritvo R, Yuwiler A, Brothers A, Freeman BJ, Plotkin S. Elevated daytime melatonin concentrations in autism: a pilot study. *Eur Child Adolesc Psychiatry* 1993;2(2):75–8.
- [53] Nir I, Meir D, Zilber N, Knobler H, Hadjez J, Lerner Y. Brief report: circadian melatonin, thyroid-stimulating hormone, prolactin, and cortisol levels in serum of young adults with autism. *J Autism Dev Disord* 1995;25:641–54.
- [54] Kulman G, Lissoni P, Rovelli F, Roselli MG, Brivio F, Sequeri P. Evidence of pineal endocrine hypofunction in autistic children. *Neuroendocrinol Lett* 2000;20:31–4.
- [55] Melke J, Goubran Botros H, Chaste P, Betancur C, Nygren G, Anckarsäter H, et al. Abnormal melatonin synthesis in autism spectrum disorders. *Mol Psychiatry* 2008;13(1):90–8.
- [56] Chugani DC, Muzik O, Behen M, Rothermel R, Janisse JJ, Lee J, et al. Developmental changes in brain serotonin synthesis capacity in autistic and nonautistic children. *Ann Neurol* 1999;45:287–95.
- [57] Chandana SR, Behen ME, Juhász C, Juhász C, Muzik O, Fothermel RD, et al. Significance of abnormalities in developmental trajectory and asymmetry of cortical serotonin synthesis in autism. *Int J Dev Neurosci* 2005;23:171–82.
- [58] Makkonen I, Riikonen R, Kokki H, Airaksinen MM, Kuikka JT. Serotonin and dopamine transporter binding in children with autism determined by SPECT. *Dev Med Child Neurol* 2008;50:593–7.
- [59] de Bruin EL, Ferdinand RF, Meester S, de Nijs FA, Verheyij F. High rates of psychiatric co-morbidity in PPD-NOS. *J Autism Dev Disord* 2007;37:877–86.
- [60] Leyfer OT, Folstein SE, Bacalman S, Davis NO, Dinh E, Morgan J. Comorbid psychiatric disorders in children with autism: interview development and rates of disorders. *J Autism Dev Dis* 2006;36:849–61.
- [61] Weiskop S, Richdale A, Matthews J. Behavioral treatment to reduce sleep problems in children with autism or fragile X syndrome. *Dev Med Child Neurol* 2005;47:94–104.
- [62] Wiggs L, France K. Behavioural treatments for sleep problems in children and adolescents with physical illness, psychological problems or intellectual disabilities. *Sleep Med Rev* 2000;4(3):299–314.
- [63] Doo S, Wing YK. Sleep problems of children with pervasive developmental disorders: correlation with parental stress. *Dev Med Child Neurol* 2006;48:650–5.
- [64] Cotton S, Richdale A. Brief report: parental description of sleep problems in children with autism, Down syndrome and Prader Will syndrome. *Res Dev Disabil* 2006;27:151–61.
- [65] Meltzer LJ. Brief report: sleep in parents of children with autism spectrum disorders. *J Pediatr Psychol* 2008;33(4):380–6.
- [66] Davidovitch M, Glick L, Holtzman G, Tirosh E, Safir MP. Developmental regression in autism: maternal perception. *J Autism Dev Disord* 2000;30:113–9.
- [67] Goldberg WA, Osann K, Filipek PA, Lulhere T, Jarvis K, Modahl C, et al. Language and other regression: assessment and timing. *J Autism Dev Disord* 2003;33:607–16.
- [68] Tuchman RF, Rapin I. Regression in pervasive developmental disorders: seizures and epileptiform electroencephalogram correlates. *Pediatrics* 1997;99:560–6.
- [69] Lord C, Shulman C, DiLavore P. Regression and word loss in autistic spectrum disorders. *J Child Psychol Psychiatr* 2004;45:936–55.
- [70] Baird G, Charman T, Pickles A, Chandler S, Loucas T, Meldrum D, et al. Regression, developmental trajectory and associated problems in disorders in the autism spectrum: the SNAP study. *J Autism Dev Disord* 2008;38(10):1827–36.
- [71] Carper RA, Courchesne E. Localized enlargement of the frontal cortex in early autism. *Biol Psychiatry* 2005;57:126–33.
- [72] Giannotti F, Cortesi F, Cerquiglini A, Bernabei P. An open-label study of controlled release melatonin in treatment of sleep disorders in children with autism. *J Autism Dev Disord* 2006;36(6):741–52.
- [73] Steffenburg S, Steffenburg U, Gillberg C. Autism spectrum disorders in children with active epilepsy and learning disability: comorbidity, pre- and perinatal background, and seizure characteristics. *Dev Med Child Neurol* 2003;45(11):724–30.
- [74] Volkmar FR, Nelson DS. Seizure disorders in autism. *J Am Acad Child Adolesc Psychiatr* 1990;29:127–9.
- [75] Hara H. Autism and epilepsy: a retrospective follow-up study. *Brain & Dev* 2007;29:486–90.
- [76] Lewine JD, Andrews R, Chez M, Patil AA, Devinski O, Smith M, et al. Magnetoencephalographic patterns of epileptiform activity in children with regressive autistic spectrum disorders. *Pediatrics* 1999;104:405–18.
- [77] Hrdlicka M, Komarek V, Propper L, Kulisek R, Zumrova A, Faladova L, et al. Not EEG abnormalities but epilepsy is associated with autistic regression and mental functioning in childhood autism. *Eur Child Adolesc Psychiatr* 2004;13:209–13.
- [78] Wiggs L, Stores G. Severe sleep disturbance and daytime challenging behaviour in children with severe learning disabilities. *J Intellect Disabil Res* 1996;40(6):518–28.
- [79] Didden R, Korzilius H, van Apele B, van Overloop C, de Vries M. Sleep problems, daytime behaviours in children with intellectual disability. *J Intellect Disabil Res* 2002;46(7):537–47.
- [80] Segawa M, Katoh M, Katoh J, Nomura Y. Early modulation of sleep parameters and its importance in later behavior. *Brain Dysfunct* 1992;5:211–23.
- [81] Malow GA, McGrew SG, Harvey M, Henderson LM, Stone WL. Impact of treating sleep apnea in a child with autism spectrum disorder. *Pediatr Neurol* 2006;34(4):325–8.
- [82] Richdale A, Wiggs L. Behavioral approaches to the treatment of sleep problems in children with developmental disorders. What is the state of the art? *Int J Behav Consult Ther* 2005;1:165–89.
- [83] Williams G, Sears L, Allard A. Parent perceptions of efficacy for strategies used to facilitate sleep in children with autism. *J Dev Phys Disabil* 2006;18(1):25–33.
- [84] Kuhn BR, Floress MT. Nonpharmacological interventions for sleep disorders in children. In: Ivanenko Anna, editor. *Sleep and psychiatric disorders in children and adolescents*. New York, London: Informa Healthcare; 2008. p. 261–78.
- [85] Jan JE, Esperzedl H, Appleton RE. The treatment of sleep disorders with melatonin. *Dev Med Child Neurol* 1994;36:97–107.
- [86] Ross C, Davies P, Whitehouse W. Melatonin treatment for sleep disorders in children with neurodevelopmental disorders: an observational study. *Dev Med Child Neurol* 2002;44:339–44.
- [87] Phillips L, Appleton R. Systematic review of melatonin treatment in children with neurodevelopmental disabilities and sleep impairment. *Dev Med Child Neurol* 2004;46:771–5.
- [88] Dodge NN, Wilson GA. Melatonin for treatment of sleep disorders in children with developmental disabilities. *J Child Neurol* 2001;16:581–4.
- [89] Andersen IM, Kaczmarek J, McGrew SG, Malow BA. Melatonin for insomnia in children with autism spectrum disorders. *J Child Neurol* 2008;23(5):482–5.
- [90] Paavonen EJ, Nieminen-von Wendt T, Vanhala R, Aronen ET, von Wendt L. Effectiveness of melatonin in the treatment of sleep disturbances in children with Asperger disorder. *J Child Adolesc Psychopharmacol* 2003;13:83–95.
- [91] Garstang J, Wallis M. Randomized controlled trial of melatonin for children with autistic spectrum disorders and sleep problems. *Child Care Health Dev* 2006;32(5):585–9.
- [92] Giannotti F, Cortesi F, Cerquiglini A, Sebastiani T, Vagnoni C, Bernabei P. The treatment of sleep disorders in childhood autism with melatonin or behavioral therapy: a randomized waiting list controlled study. *Sleep* 2008;A58 (31).
- [93] Jan JE, Hamilton D, Seward N, Fast DK, Freeman RD, Laudon M. Clinical trials of controlled-release melatonin in children with sleep–wake cycle disorders. *J Pineal Res* 2000;29(1):34–9.
- [94] Rossi PG, Posar A, Parmeggiani A, Pipitone E, D'Agata M. Niazepam in the treatment of autistic disorder. *J Child Neurol* 1999;14(8):547–50.
- [95] Owens JA. Pharmacotherapy of pediatric insomnia. *J Am Acad Child Adolesc Psychiatry* 2009;48(2):99–107.