

Signorini's Problem


Max Stuthmann

March 17, 2026

- 1 Introduction – What's the Problem?
- 2 Geometrical and Physical Setup
 - Boundary Decomposition
 - Elasticity and Forces
 - Admissible Displacements
- 3 Energy Minimization
- 4 Weak Formulations
- 5 Strong Formulations
 - A Hybrid Formulation
 - Classical Strong Formulation


Rough Problem Setup

- *Body:* $\Omega \subset \mathbb{R}^d$, $d \in \{2, 3\}$ open, bounded, connected with Lipschitz boundary $\Gamma := \partial\Omega$.




Rough Problem Setup

- ▶ *Body:* $\Omega \subset \mathbb{R}^d$, $d \in \{2, 3\}$ open, bounded, connected with Lipschitz boundary $\Gamma := \partial\Omega$.
- ▶ Ω is sitting on a *rigid foundation*.




Rough Problem Setup

- ▶ *Body:* $\Omega \subset \mathbb{R}^d$, $d \in \{2, 3\}$ open, bounded, connected with Lipschitz boundary $\Gamma := \partial\Omega$.
- ▶ Ω is sitting on a *rigid foundation*.
- ▶ No frictions assumed between Ω and foundation.




Rough Problem Setup

- ▶ *Body:* $\Omega \subset \mathbb{R}^d$, $d \in \{2, 3\}$ open, bounded, connected with Lipschitz boundary $\Gamma := \partial\Omega$.
- ▶ Ω is sitting on a *rigid foundation*.
- ▶ No frictions assumed between Ω and foundation.
- ▶ Ω may be subject to elasticity properties and internal/external forces.





Rough Problem Setup

- ▶ *Body:* $\Omega \subset \mathbb{R}^d$, $d \in \{2, 3\}$ open, bounded, connected with Lipschitz boundary $\Gamma := \partial\Omega$.
- ▶ Ω is sitting on a *rigid foundation*.
- ▶ No frictions assumed between Ω and foundation.
- ▶ Ω may be subject to elasticity properties and internal/external forces.
- ▶ **Not in equilibrium configuration.**



What is to be found?



- ▶ Let $\varphi(\Omega)$ denote Ω in equilibrium configuration.



What is to be found?

- ▶ Let $\varphi(\Omega)$ denote Ω in equilibrium configuration.
- ▶ $\varphi : \bar{\Omega} \rightarrow \mathbb{R}^d$ maps $\bar{\Omega}$ onto $\varphi(\bar{\Omega})$ and can be written as

$$\varphi(x) = x + u(x), \quad u : \bar{\Omega} \rightarrow \mathbb{R}^d.$$





What is to be found?

- ▶ Let $\varphi(\Omega)$ denote Ω in equilibrium configuration.
- ▶ $\varphi : \bar{\Omega} \rightarrow \mathbb{R}^d$ maps $\bar{\Omega}$ onto $\varphi(\bar{\Omega})$ and can be written as

$$\varphi(x) = x + u(x), \quad u : \bar{\Omega} \rightarrow \mathbb{R}^d.$$

- ▶ We aim to find the *displacement function* u .





Content

- 1 Introduction – What's the Problem?
- 2 Geometrical and Physical Setup
 - Boundary Decomposition
 - Elasticity and Forces
 - Admissible Displacements
- 3 Energy Minimization
- 4 Weak Formulations
- 5 Strong Formulations



Boundary Decomposition and Preview on the Boundary Conditions

- Decompose $\Gamma = \partial\Omega$ into three relatively open and pairwise disjoint parts $\Gamma = \overline{\Gamma_D} \dot{\cup} \overline{\Gamma_N} \dot{\cup} \overline{\Gamma_C}$ with $|\Gamma_D|, |\Gamma_C| > 0$ and $|\Gamma_N| \geq 0$.





Boundary Decomposition and Preview on the Boundary Conditions

- ▶ Decompose $\Gamma = \partial\Omega$ into three relatively open and pairwise disjoint parts $\Gamma = \overline{\Gamma_D} \dot{\cup} \overline{\Gamma_N} \dot{\cup} \overline{\Gamma_C}$ with $|\Gamma_D|, |\Gamma_C| > 0$ and $|\Gamma_N| \geq 0$.
- ▶ Each part has different boundary conditions:





Boundary Decomposition and Preview on the Boundary Conditions

- ▶ Decompose $\Gamma = \partial\Omega$ into three relatively open and pairwise disjoint parts $\Gamma = \overline{\Gamma_D} \dot{\cup} \overline{\Gamma_N} \dot{\cup} \overline{\Gamma_C}$ with $|\Gamma_D|, |\Gamma_C| > 0$ and $|\Gamma_N| \geq 0$.
- ▶ Each part has different boundary conditions:
 - ▷ Γ_D : Homogeneous Dirichlet. (Clamped boundary part).





Boundary Decomposition and Preview on the Boundary Conditions

- ▶ Decompose $\Gamma = \partial\Omega$ into three relatively open and pairwise disjoint parts $\Gamma = \overline{\Gamma_D} \dot{\cup} \overline{\Gamma_N} \dot{\cup} \overline{\Gamma_C}$ with $|\Gamma_D|, |\Gamma_C| > 0$ and $|\Gamma_N| \geq 0$.
- ▶ Each part has different boundary conditions:
 - ▷ Γ_D : Homogeneous Dirichlet. (Clamped boundary part).
 - ▷ Γ_N : Neumann conditions. (We impose known surface force density).





Boundary Decomposition and Preview on the Boundary Conditions

- ▶ Decompose $\Gamma = \partial\Omega$ into three relatively open and pairwise disjoint parts $\Gamma = \overline{\Gamma_D} \dot{\cup} \overline{\Gamma_N} \dot{\cup} \overline{\Gamma_C}$ with $|\Gamma_D|, |\Gamma_C| > 0$ and $|\Gamma_N| \geq 0$.
- ▶ Each part has different boundary conditions:
 - ▷ Γ_D : Homogeneous Dirichlet. (Clamped boundary part).
 - ▷ Γ_N : Neumann conditions. (We impose known surface force density).
 - ▷ Γ_C is the *potential* contact boundary, i.e. Γ_C is prescribed a priori but the active contact zone $\Gamma_{\text{act}}(u) \subset \Gamma_C$ is unknown. On Γ_C : Contact conditions. (Nonpenetration, Complementarity, ...)



Boundary Decomposition and Preview on the Boundary Conditions

- ▶ Decompose $\Gamma = \partial\Omega$ into three relatively open and pairwise disjoint parts $\Gamma = \overline{\Gamma_D} \dot{\cup} \overline{\Gamma_N} \dot{\cup} \overline{\Gamma_C}$ with $|\Gamma_D|, |\Gamma_C| > 0$ and $|\Gamma_N| \geq 0$.
- ▶ Each part has different boundary conditions:
 - ▷ Γ_D : Homogeneous Dirichlet. (Clamped boundary part).
 - ▷ Γ_N : Neumann conditions. (We impose known surface force density).
 - ▷ Γ_C is the *potential* contact boundary, i.e. Γ_C is prescribed a priori but the active contact zone $\Gamma_{\text{act}}(u) \subset \Gamma_C$ is unknown. On Γ_C : Contact conditions. (Nonpenetration, Complementarity, ...)
- ▶ Rigid foundation can be describe by gap-function $g : \Gamma_C \rightarrow \mathbb{R}$. We assume $g = 0$.



Content

- 1 Introduction – What's the Problem?
- 2 Geometrical and Physical Setup
 - Boundary Decomposition
 - Elasticity and Forces
 - Admissible Displacements
- 3 Energy Minimization
- 4 Weak Formulations
- 5 Strong Formulations

Basic Notations on Sobolev Spaces

- ▶ On $H^s(\Omega; \mathbb{R}^d)$ for $s = k + \theta$ with $k \in \mathbb{N}_0$ and $\theta \in [0, 1)$, we use $(\cdot, \cdot)_{s, \Omega}$ and $\|\cdot\|_{s, \Omega}$

Basic Notations on Sobolev Spaces

- ▶ On $H^s(\Omega; \mathbb{R}^d)$ for $s = k + \theta$ with $k \in \mathbb{N}_0$ and $\theta \in [0, 1)$, we use $(\cdot, \cdot)_{s, \Omega}$ and $\|\cdot\|_{s, \Omega}$
- ▶ $H^0(\Omega; \mathbb{R}^d) = L^2(\Omega; \mathbb{R}^d)$.

Basic Notations on Sobolev Spaces

- ▶ On $H^s(\Omega; \mathbb{R}^d)$ for $s = k + \theta$ with $k \in \mathbb{N}_0$ and $\theta \in [0, 1)$, we use $(\cdot, \cdot)_{s, \Omega}$ and $\|\cdot\|_{s, \Omega}$
- ▶ $H^0(\Omega; \mathbb{R}^d) = L^2(\Omega; \mathbb{R}^d)$.
- ▶ *Trace Theorem*: Restriction of smooth functions $\overline{\Omega} \rightarrow \mathbb{R}^d$ onto the boundary $\Gamma = \partial\Omega \in \mathcal{C}^{m,1}$ extends uniquely to a continuous surjective linear trace operator

$$\gamma : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1/2}(\Gamma; \mathbb{R}^d), \quad \frac{1}{2} < s \leq m + 1.$$

Basic Notations on Sobolev Spaces

- ▶ On $H^s(\Omega; \mathbb{R}^d)$ for $s = k + \theta$ with $k \in \mathbb{N}_0$ and $\theta \in [0, 1)$, we use $(\cdot, \cdot)_{s, \Omega}$ and $\|\cdot\|_{s, \Omega}$
- ▶ $H^0(\Omega; \mathbb{R}^d) = L^2(\Omega; \mathbb{R}^d)$.
- ▶ *Trace Theorem*: Restriction of smooth functions $\overline{\Omega} \rightarrow \mathbb{R}^d$ onto the boundary $\Gamma = \partial\Omega \in \mathcal{C}^{m,1}$ extends uniquely to a continuous surjective linear trace operator

$$\gamma : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1/2}(\Gamma; \mathbb{R}^d), \quad \frac{1}{2} < s \leq m + 1.$$

- ▶ For now, Ω is Lipschitz, i.e. $m = 0$ and we need $s = 1$, so

$$\gamma : H^1(\Omega; \mathbb{R}^d) \rightarrow H^{1/2}(\Gamma; \mathbb{R}^d)$$

Small Strain Elasticity

- *Symmetric second order tensors:*

$$\mathbb{S}^d := \{s \in \mathbb{R}^d \otimes \mathbb{R}^d \mid s^\top = s\}$$

Small Strain Elasticity

- ▶ *Symmetric second order tensors:*

$$\mathbb{S}^d := \{s \in \mathbb{R}^d \otimes \mathbb{R}^d \mid s^\top = s\}$$

- ▶ *Stresses* are described by fields $\tau : \Omega \rightarrow \mathbb{S}^d$

Small Strain Elasticity

- ▶ *Symmetric second order tensors:*

$$\mathbb{S}^d := \{s \in \mathbb{R}^d \otimes \mathbb{R}^d \mid s^\top = s\}$$

- ▶ *Stresses* are described by fields $\tau : \Omega \rightarrow \mathbb{S}^d$
- ▶ We identify $\mathbb{S}^d$ with symmetric $d \times d$ matrices.

Small Strain Elasticity

- *Symmetric second order tensors:*

$$\mathbb{S}^d := \{s \in \mathbb{R}^d \otimes \mathbb{R}^d \mid s^\top = s\}$$

- *Stresses* are described by fields $\tau : \Omega \rightarrow \mathbb{S}^d$
- We identify $\mathbb{S}^d$ with symmetric $d \times d$ matrices.
- On $\mathbb{R}^{d \times d}$, we use the *Frobenius product*

$$A : B := \sum_{j=1}^d \sum_{k=1}^d A_{jk} B_{jk}, \quad |A| := \sqrt{A : A}.$$

Small Strain Elasticity

- *Symmetric second order tensors:*

$$\mathbb{S}^d := \{s \in \mathbb{R}^d \otimes \mathbb{R}^d \mid s^\top = s\}$$

- *Stresses* are described by fields $\tau : \Omega \rightarrow \mathbb{S}^d$
- We identify $\mathbb{S}^d$ with symmetric $d \times d$ matrices.
- On $\mathbb{R}^{d \times d}$, we use the *Frobenius product*

$$A : B := \sum_{j=1}^d \sum_{k=1}^d A_{jk} B_{jk}, \quad |A| := \sqrt{A : A}.$$

- If $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$ is a fourth order tensor and $s \in \mathbb{S}^d$, we use the *double contraction*

$$s : \mathcal{A}(x) : s := \sum_{i=1}^d \sum_{j=1}^d \sum_{k=1}^d \sum_{l=1}^d \mathcal{A}_{ijkl}(x) s_{ij} s_{kl}.$$

Small Strain Elasticity

- ▶ Let $s \geq 1$.

Small Strain Elasticity

- ▶ Let $s \geq 1$.
- ▶ *Small strain tensor:*

$$\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \varepsilon(u) := \frac{1}{2} (\nabla u + \nabla u^\top).$$

Small Strain Elasticity

- ▶ Let $s \geq 1$.
- ▶ *Small strain tensor:*

$$\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \varepsilon(u) := \frac{1}{2} (\nabla u + \nabla u^\top).$$

- ▶ *Elasticity of Ω is describes by a fourth order tensor $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$.*

Small Strain Elasticity

- ▶ Let $s \geq 1$.
- ▶ *Small strain tensor:*

$$\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \varepsilon(u) := \frac{1}{2} (\nabla u + \nabla u^\top).$$

- ▶ *Elasticity of Ω is describes by a fourth order tensor $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$.*
- ▶ *$\mathcal{A}$ is assumed symmetric and uniformly elliptic and bounded.*

Small Strain Elasticity

- ▶ Let $s \geq 1$.
- ▶ *Small strain tensor:*

$$\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \varepsilon(u) := \frac{1}{2} (\nabla u + \nabla u^\top).$$

- ▶ Elasticity of Ω is describes by a *fourth order tensor* $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$.
- ▶ $\mathcal{A}$ is assumed *symmetric* and *uniformly elliptic* and *bounded*.
- ▶ *(Cauchy) stress tensor:*

$$\sigma : H^1(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \mathcal{A} : \varepsilon(u).$$

Small Strain Elasticity

- ▶ Let $s \geq 1$.
- ▶ *Small strain tensor:*

$$\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \varepsilon(u) := \frac{1}{2} (\nabla u + \nabla u^\top).$$

- ▶ Elasticity of Ω is describes by a *fourth order tensor* $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$.
- ▶ $\mathcal{A}$ is assumed *symmetric* and *uniformly elliptic* and *bounded*.
- ▶ *(Cauchy) stress tensor:*

$$\sigma : H^1(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \mathcal{A} : \varepsilon(u).$$

- ▶ $\mathcal{A}$ has to be *sufficiently smooth* for this.

Small Strain Elasticity

- ▶ Let $s \geq 1$.
- ▶ *Small strain tensor:*

$$\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \varepsilon(u) := \frac{1}{2} (\nabla u + \nabla u^\top).$$

- ▶ Elasticity of Ω is describes by a *fourth order tensor* $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$.
- ▶ $\mathcal{A}$ is assumed *symmetric* and *uniformly elliptic* and *bounded*.
- ▶ *(Cauchy) stress tensor:*

$$\sigma : H^1(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \mathcal{A} : \varepsilon(u).$$

- ▶ $\mathcal{A}$ has to be *sufficiently smooth* for this.
- ▶ For now, we need $s = 1$, i.e. $\varepsilon(u) \in L^2(\Omega; \mathbb{S}^d)$ and $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$.

Small Strain Elasticity

- ▶ Let $s \geq 1$.
- ▶ *Small strain tensor:*

$$\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \varepsilon(u) := \frac{1}{2} (\nabla u + \nabla u^\top).$$

- ▶ Elasticity of Ω is describes by a *fourth order tensor* $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$.
- ▶ $\mathcal{A}$ is assumed *symmetric* and *uniformly elliptic* and *bounded*.
- ▶ *(Cauchy) stress tensor:*

$$\sigma : H^1(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d), \quad u \mapsto \mathcal{A} : \varepsilon(u).$$

- ▶ $\mathcal{A}$ has to be *sufficiently smooth* for this.
- ▶ For now, we need $s = 1$, i.e. $\varepsilon(u) \in L^2(\Omega; \mathbb{S}^d)$ and $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$.
- ▶ $\mathcal{A}_{ijkl} \in L^\infty(\Omega)$ is sufficient for $s = 1$. Later on, we need more.

Virtual Work of Internal and External Forces

- ▶ *Volume force density:* $f \in L^2(\Omega; \mathbb{R}^d)$.

Virtual Work of Internal and External Forces

- ▶ *Volume force density:* $f \in L^2(\Omega; \mathbb{R}^d)$.
- ▶ *Surface load on Neumann part:* $F \in L^2(\Gamma_N; \mathbb{R}^d)$.

Virtual Work of Internal and External Forces

- ▶ *Volume force density:* $f \in L^2(\Omega; \mathbb{R}^d)$.
- ▶ *Surface load on Neumann part:* $F \in L^2(\Gamma_N; \mathbb{R}^d)$.
- ▶ *Work = force · displacement:* The linear functional

$$\ell(v) := \int_{\Omega} f \cdot v \, dx + \int_{\Gamma_N} F \cdot (\gamma v) \, ds, \quad v \in H^1(\Omega; \mathbb{R}^d)$$

describes the *virtual work of external forces*.

Virtual Work of Internal and External Forces

- ▶ *Volume force density:* $f \in L^2(\Omega; \mathbb{R}^d)$.
- ▶ *Surface load on Neumann part:* $F \in L^2(\Gamma_N; \mathbb{R}^d)$.
- ▶ *Work = force · displacement:* The linear functional

$$\ell(v) := \int_{\Omega} f \cdot v \, dx + \int_{\Gamma_N} F \cdot (\gamma v) \, ds, \quad v \in H^1(\Omega; \mathbb{R}^d)$$

describes the *virtual work of external forces*.

- ▶ *Internal forces* are described by the *stress* $\sigma(u) = \mathcal{A} : \varepsilon(u)$.

Virtual Work of Internal and External Forces

- ▶ *Volume force density*: $f \in L^2(\Omega; \mathbb{R}^d)$.
- ▶ *Surface load on Neumann part*: $F \in L^2(\Gamma_N; \mathbb{R}^d)$.
- ▶ *Work = force · displacement*: The linear functional

$$\ell(v) := \int_{\Omega} f \cdot v \, dx + \int_{\Gamma_N} F \cdot (\gamma v) \, ds, \quad v \in H^1(\Omega; \mathbb{R}^d)$$

describes the *virtual work of external forces*.

- ▶ *Internal forces* are described by the *stress* $\sigma(u) = \mathcal{A} : \varepsilon(u)$.
- ▶ *Work = force · displacement*: The symmetric bilinear form

$$a(u, v) := \int_{\Omega} \sigma(u) : \varepsilon(v) \, dx, \quad u, v \in H^1(\Omega; \mathbb{R}^d).$$

describes the *virtual work of internal forces*.

Virtual Work of Internal and External Forces

- ▶ *Volume force density*: $f \in L^2(\Omega; \mathbb{R}^d)$.
- ▶ *Surface load on Neumann part*: $F \in L^2(\Gamma_N; \mathbb{R}^d)$.
- ▶ *Work = force · displacement*: The linear functional

$$\ell(v) := \int_{\Omega} f \cdot v \, dx + \int_{\Gamma_N} F \cdot (\gamma v) \, ds, \quad v \in H^1(\Omega; \mathbb{R}^d)$$

describes the *virtual work of external forces*.

- ▶ *Internal forces* are described by the *stress* $\sigma(u) = \mathcal{A} : \varepsilon(u)$.
- ▶ *Work = force · displacement*: The symmetric bilinear form

$$a(u, v) := \int_{\Omega} \sigma(u) : \varepsilon(v) \, dx, \quad u, v \in H^1(\Omega; \mathbb{R}^d).$$

describes the *virtual work of internal forces*.

- ▶ *Equilibrium* between both: $a(u, v) = \ell(v) \quad \forall v \in H^1(\Omega; \mathbb{R}^d)$.

Standard Properties of a and ℓ

Proposition

The symmetric bilinear form $a : V \times V \rightarrow \mathbb{R}$ defined above is

- i) *bounded, i.e. for all $u, v \in V$ there exists $C_1 < \infty$ such that*

$$|a(u, v)| \leq C_1 \|u\|_{1,\Omega} \|v\|_{1,\Omega}.$$

- ii) *V -elliptic, i.e. for all $v \in V \setminus \{0\}$ there exist $C_2 > 0$ such that*

$$a(v, v) \geq C_2 \|v\|_{1,\Omega}^2.$$

Moreover, the linear functional ℓ defined above is bounded, i.e. for all $v \in V$ there exists $C_3 > 0$ such that

$$|\ell(v)| \leq C_3 \|v\|_{1,\Omega}.$$

Content

1 Introduction – What's the Problem?

2 Geometrical and Physical Setup

- Boundary Decomposition
- Elasticity and Forces
- Admissible Displacements

3 Energy Minimization


4 Weak Formulations

5 Strong Formulations

Admissible Displacements and the Nonpenetration Constraint

- *Homogeneous Dirichlet* boundary conditions on Γ_D are incorporated in the underlying Hilbert space

$$V := \{v \in H^1(\Omega; \mathbb{R}^d) \mid (\gamma v)|_{\Gamma_D} = 0\}$$




Admissible Displacements and the Nonpenetration Constraint

- ▶ *Homogeneous Dirichlet* boundary conditions on Γ_D are incorporated in the underlying Hilbert space

$$V := \{v \in H^1(\Omega; \mathbb{R}^d) \mid (\gamma v)|_{\Gamma_D} = 0\}$$

- ▶ *Normal Component*: For $v \in V$, we define its *normal component* by $v_n := (\gamma v) \cdot \mathbf{n}$ with the *outward normal* $\mathbf{n} \in L^\infty(\Gamma; \mathbb{R}^d)$.




Admissible Displacements and the Nonpenetration Constraint

- ▶ *Homogeneous Dirichlet* boundary conditions on Γ_D are incorporated in the underlying Hilbert space

$$V := \{v \in H^1(\Omega; \mathbb{R}^d) \mid (\gamma v)|_{\Gamma_D} = 0\}$$

- ▶ *Normal Component*: For $v \in V$, we define its *normal component* by $v_n := (\gamma v) \cdot \mathbf{n}$ with the *outward normal* $\mathbf{n} \in L^\infty(\Gamma; \mathbb{R}^d)$.
- ▶ *Nonpenetration constraint*: $u_n \leq g = 0$ almost everywhere on Γ_C .




Admissible Displacements and the Nonpenetration Constraint

- ▶ *Homogeneous Dirichlet* boundary conditions on Γ_D are incorporated in the underlying Hilbert space

$$V := \{v \in H^1(\Omega; \mathbb{R}^d) \mid (\gamma v)|_{\Gamma_D} = 0\}$$

- ▶ *Normal Component*: For $v \in V$, we define its *normal component* by $v_n := (\gamma v) \cdot \mathbf{n}$ with the *outward normal* $\mathbf{n} \in L^\infty(\Gamma; \mathbb{R}^d)$.
- ▶ *Nonpenetration constraint*: $u_n \leq g = 0$ almost everywhere on Γ_C .
- ▶ *Set of admissible displacements*: $K := \{v \in V \mid v_n \leq 0 \text{ a.e. on } \Gamma_C\}$.




Admissible Displacements and the Nonpenetration Constraint

- ▶ *Homogeneous Dirichlet* boundary conditions on Γ_D are incorporated in the underlying Hilbert space

$$V := \{v \in H^1(\Omega; \mathbb{R}^d) \mid (\gamma v)|_{\Gamma_D} = 0\}$$

- ▶ *Normal Component*: For $v \in V$, we define its *normal component* by $v_n := (\gamma v) \cdot \mathbf{n}$ with the *outward normal* $\mathbf{n} \in L^\infty(\Gamma; \mathbb{R}^d)$.
- ▶ *Nonpenetration constraint*: $u_n \leq g = 0$ almost everywhere on Γ_C .
- ▶ *Set of admissible displacements*: $K := \{v \in V \mid v_n \leq 0 \text{ a.e. on } \Gamma_C\}$.
- ▶ K is a closed and convex cone in V , i.e. if $v \in K$ then $\lambda v \in K$ for all $\lambda \in [0, \infty)$.



► *Equilibrium:*

$$a(u, v) = \int_{\Omega} \sigma(u) : \varepsilon(v) \, dx = \int_{\Omega} f \cdot v \, dx + \int_{\Gamma_N} F \cdot (\gamma v) \, ds = \ell(v) \quad \forall v \in V$$

- *Equilibrium:*

$$a(u, v) = \int_{\Omega} \sigma(u) : \varepsilon(v) \, dx = \int_{\Omega} f \cdot v \, dx + \int_{\Gamma_N} F \cdot (\gamma v) \, ds = \ell(v) \quad \forall v \in V$$

- The *potential energy* of the elastic body is given by the quadratic functional

$$\mathcal{J}(v) := \frac{1}{2}a(v, v) - \ell(v), \quad v \in V.$$

Energy Minimization

- *Equilibrium:*

$$a(u, v) = \int_{\Omega} \sigma(u) : \varepsilon(v) \, dx = \int_{\Omega} f \cdot v \, dx + \int_{\Gamma_N} F \cdot (\gamma v) \, ds = \ell(v) \quad \forall v \in V$$

- The *potential energy* of the elastic body is given by the quadratic functional

$$\mathcal{J}(v) := \frac{1}{2}a(v, v) - \ell(v), \quad v \in V.$$

- *Nonpenetration:* Minimize $\mathcal{J}(v)$ over all *admissible* displacements $v \in K$.

Energy Minimization

- *Equilibrium:*

$$a(u, v) = \int_{\Omega} \sigma(u) : \varepsilon(v) \, dx = \int_{\Omega} f \cdot v \, dx + \int_{\Gamma_N} F \cdot (\gamma v) \, ds = \ell(v) \quad \forall v \in V$$

- The *potential energy* of the elastic body is given by the quadratic functional

$$\mathcal{J}(v) := \frac{1}{2}a(v, v) - \ell(v), \quad v \in V.$$

- *Nonpenetration:* Minimize $\mathcal{J}(v)$ over all *admissible* displacements $v \in K$.

Problem (Signorini's Problem – Energy Minimization)

$$\text{Find } u \in K \text{ such that: } \mathcal{J}(u) = \min_{v \in K} \mathcal{J}(v).$$

Proposition (Well-Posedness of Signorini's Problem)

There exists exactly one $u \in K$ that minimizes $\mathcal{J}$ over K . Moreover, the unique minimizer satisfies the a-priori bound

$$\|u\|_{1,\Omega} \leq \frac{2C_3}{C_2} \|\ell\|_{V'}.$$

Proof sketch:

Proposition (Well-Posedness of Signorini's Problem)

There exists exactly one $u \in K$ that minimizes $\mathcal{J}$ over K . Moreover, the unique minimizer satisfies the a-priori bound

$$\|u\|_{1,\Omega} \leq \frac{2C_3}{C_2} \|\ell\|_{V'}.$$

Proof sketch:

- ▶ Prove that $\mathcal{J}$ is coercive and weakly lower semicontinuous.

Proposition (Well-Posedness of Signorini's Problem)

There exists exactly one $u \in K$ that minimizes $\mathcal{J}$ over K . Moreover, the unique minimizer satisfies the a-priori bound

$$\|u\|_{1,\Omega} \leq \frac{2C_3}{C_2} \|\ell\|_{V'}.$$

Proof sketch:

- ▶ Prove that $\mathcal{J}$ is *coercive* and *weakly lower semicontinuous*.
- ▶ This implies existence.

Proposition (Well-Posedness of Signorini's Problem)

There exists exactly one $u \in K$ that minimizes $\mathcal{J}$ over K . Moreover, the unique minimizer satisfies the a-priori bound

$$\|u\|_{1,\Omega} \leq \frac{2C_3}{C_2} \|\ell\|_{V'}.$$

Proof sketch:

- ▶ Prove that $\mathcal{J}$ is *coercive* and *weakly lower semicontinuous*.
- ▶ This implies existence.
- ▶ Prove that $\mathcal{J}$ is *strictly convex*.

Proposition (Well-Posedness of Signorini's Problem)

There exists exactly one $u \in K$ that minimizes $\mathcal{J}$ over K . Moreover, the unique minimizer satisfies the a-priori bound

$$\|u\|_{1,\Omega} \leq \frac{2C_3}{C_2} \|\ell\|_{V'}.$$

Proof sketch:

- ▶ Prove that $\mathcal{J}$ is *coercive* and *weakly lower semicontinuous*.
- ▶ This implies existence.
- ▶ Prove that $\mathcal{J}$ is *strictly convex*.
- ▶ With convexity of K , this implies uniqueness.

Proposition (Well-Posedness of Signorini's Problem)

There exists exactly one $u \in K$ that minimizes $\mathcal{J}$ over K . Moreover, the unique minimizer satisfies the a-priori bound

$$\|u\|_{1,\Omega} \leq \frac{2C_3}{C_2} \|\ell\|_{V'}.$$

Proof sketch:

- ▶ Prove that $\mathcal{J}$ is *coercive* and *weakly lower semicontinuous*.
- ▶ This implies existence.
- ▶ Prove that $\mathcal{J}$ is *strictly convex*.
- ▶ With convexity of K , this implies uniqueness.
- ▶ Stability bound follows from V -ellipticity of a w.r.t. C_2 and boundedness of ℓ w.r.t. C_3 .

Proposition (First-order optimality condition on convex sets)

A function $u \in K$ minimizes $\mathcal{J}$ over K if and only if it satisfies the variational inequality

$$\mathcal{J}'(u)[v - u] = a(u, v - u) - \ell(v - u) \geq 0 \quad \forall v \in K.$$

Weak Formulations / Variational Inequality

Proposition (First-order optimality condition on convex sets)

A function $u \in K$ minimizes $\mathcal{J}$ over K if and only if it satisfies the variational inequality

$$\mathcal{J}'(u)[v - u] = a(u, v - u) - \ell(v - u) \geq 0 \quad \forall v \in K.$$

Problem (Signorini's Problem – Weak Formulation 1)

Find $u \in K$ such that: $a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$

Weak Formulations / Variational Inequality

Proposition (First-order optimality condition on convex sets)

A function $u \in K$ minimizes $\mathcal{J}$ over K if and only if it satisfies the variational inequality

$$\mathcal{J}'(u)[v - u] = a(u, v - u) - \ell(v - u) \geq 0 \quad \forall v \in K.$$

Problem (Signorini's Problem – Weak Formulation 1)

Find $u \in K$ such that: $a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$

Problem (Signorini's Problem – Weak Formulation 2)

Find $u \in K$ such that:
$$\begin{cases} a(u, u) = \ell(u), \\ a(u, v) \geq \ell(v) \quad \forall v \in K. \end{cases}$$

Content

1 Introduction – What's the Problem?

2 Geometrical and Physical Setup

3 Energy Minimization

4 Weak Formulations

5 Strong Formulations

- A Hybrid Formulation
- Classical Strong Formulation

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)_n$ for the stress $\sigma(u)$.

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)_n$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)\mathbf{n}$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.
- ▶ Therefore $v_{\mathbf{n}} = (\gamma v) \cdot \mathbf{n} \in L^2(\Gamma)$.

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)\mathbf{n}$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.
- ▶ Therefore $v_{\mathbf{n}} = (\gamma v) \cdot \mathbf{n} \in L^2(\Gamma)$.
- ▶ For our $u \in V$, we have $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$, i.e. the earlier trace theorem does not apply to $\sigma(u)$.

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)_n$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.
- ▶ Therefore $v_n = (\gamma v) \cdot n \in L^2(\Gamma)$.
- ▶ For our $u \in V$, we have $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$, i.e. the earlier trace theorem does not apply to $\sigma(u)$.
- ▶ We need an additional assumption on the Regularity of $\sigma(u)$.

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)\mathbf{n}$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.
- ▶ Therefore $v_{\mathbf{n}} = (\gamma v) \cdot \mathbf{n} \in L^2(\Gamma)$.
- ▶ For our $u \in V$, we have $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$, i.e. the earlier trace theorem does not apply to $\sigma(u)$.
- ▶ We need an additional assumption on the Regularity of $\sigma(u)$.
- ▶ If we assume $\sigma(u) \in H^1(\Omega; \mathbb{S}^d)$, then $\gamma \sigma(u) \in L^2(\Gamma; \mathbb{S}^d)$ and $(\gamma \sigma(u))\mathbf{n} \in L^2(\Gamma; \mathbb{R}^d)$.

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)\mathbf{n}$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.
- ▶ Therefore $v_{\mathbf{n}} = (\gamma v) \cdot \mathbf{n} \in L^2(\Gamma)$.
- ▶ For our $u \in V$, we have $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$, i.e. the earlier trace theorem does not apply to $\sigma(u)$.
- ▶ We need an additional assumption on the Regularity of $\sigma(u)$.
- ▶ If we assume $\sigma(u) \in H^1(\Omega; \mathbb{S}^d)$, then $\gamma \sigma(u) \in L^2(\Gamma; \mathbb{S}^d)$ and $(\gamma \sigma(u))\mathbf{n} \in L^2(\Gamma; \mathbb{R}^d)$.
- ▶ This assumption is much stronger than needed!

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)\mathbf{n}$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.
- ▶ Therefore $v_{\mathbf{n}} = (\gamma v) \cdot \mathbf{n} \in L^2(\Gamma)$.
- ▶ For our $u \in V$, we have $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$, i.e. the earlier trace theorem does not apply to $\sigma(u)$.
- ▶ We need an additional assumption on the Regularity of $\sigma(u)$.
- ▶ If we assume $\sigma(u) \in H^1(\Omega; \mathbb{S}^d)$, then $\gamma \sigma(u) \in L^2(\Gamma; \mathbb{S}^d)$ and $(\gamma \sigma(u))\mathbf{n} \in L^2(\Gamma; \mathbb{R}^d)$.
- ▶ This assumption is much stronger than needed!
- ▶ Instead, we assume $\sigma(u) \in H(\operatorname{div}; \Omega)$, where

$$H(\operatorname{div}; \Omega) := \{ \tau \in L^2(\Omega; \mathbb{S}^d) \mid \operatorname{div} \tau \in L^2(\Omega; \mathbb{R}^d) \},$$

where $\operatorname{div} \tau$ is understood row-wise in the weak sense.

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)\mathbf{n}$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.
- ▶ Therefore $v_{\mathbf{n}} = (\gamma v) \cdot \mathbf{n} \in L^2(\Gamma)$.
- ▶ For our $u \in V$, we have $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$, i.e. the earlier trace theorem does not apply to $\sigma(u)$.
- ▶ We need an additional assumption on the Regularity of $\sigma(u)$.
- ▶ If we assume $\sigma(u) \in H^1(\Omega; \mathbb{S}^d)$, then $\gamma \sigma(u) \in L^2(\Gamma; \mathbb{S}^d)$ and $(\gamma \sigma(u))\mathbf{n} \in L^2(\Gamma; \mathbb{R}^d)$.
- ▶ This assumption is much stronger than needed!
- ▶ Instead, we assume $\sigma(u) \in H(\operatorname{div}; \Omega)$, where

$$H(\operatorname{div}; \Omega) := \{ \tau \in L^2(\Omega; \mathbb{S}^d) \mid \operatorname{div} \tau \in L^2(\Omega; \mathbb{R}^d) \},$$

where $\operatorname{div} \tau$ is understood row-wise in the weak sense.

- ▶ Then $\operatorname{div} \sigma(u) \in L^2(\Omega; \mathbb{R}^d)$, which is needed for strong equilibrium equations – $\operatorname{div} \sigma(u) = f$ with L^2 -source terms f .

Additional Stress Regularity

- ▶ **Goal:** We need a concept of normal traces $\sigma(u)\mathbf{n}$ for the stress $\sigma(u)$.
- ▶ For displacements $v \in H^1(\Omega; \mathbb{R}^d)$, we have $\gamma v \in H^{1/2}(\Gamma; \mathbb{R}^d) \subset L^2(\Gamma; \mathbb{R}^d)$.
- ▶ Therefore $v_{\mathbf{n}} = (\gamma v) \cdot \mathbf{n} \in L^2(\Gamma)$.
- ▶ For our $u \in V$, we have $\sigma(u) \in L^2(\Omega; \mathbb{S}^d)$, i.e. the earlier trace theorem does not apply to $\sigma(u)$.
- ▶ We need an additional assumption on the Regularity of $\sigma(u)$.
- ▶ If we assume $\sigma(u) \in H^1(\Omega; \mathbb{S}^d)$, then $\gamma \sigma(u) \in L^2(\Gamma; \mathbb{S}^d)$ and $(\gamma \sigma(u))\mathbf{n} \in L^2(\Gamma; \mathbb{R}^d)$.
- ▶ This assumption is much stronger than needed!
- ▶ Instead, we assume $\sigma(u) \in H(\operatorname{div}; \Omega)$, where

$$H(\operatorname{div}; \Omega) := \{ \tau \in L^2(\Omega; \mathbb{S}^d) \mid \operatorname{div} \tau \in L^2(\Omega; \mathbb{R}^d) \},$$

where $\operatorname{div} \tau$ is understood row-wise in the weak sense.

- ▶ Then $\operatorname{div} \sigma(u) \in L^2(\Omega; \mathbb{R}^d)$, which is needed for strong equilibrium equations $-\operatorname{div} \sigma(u) = f$ with L^2 -source terms f .
- ▶ Normal traces on $H(\operatorname{div}; \Omega)$ are understood as follows.

Normal Traces on $H(\operatorname{div}; \Omega)$

- ▶ We denote by $H^{-1/2}(\Gamma; \mathbb{R}^d)$ the dual space of $H^{1/2}(\Gamma; \mathbb{R}^d)$ with duality pairing $\langle \cdot, \cdot \rangle_\Gamma$.

Normal Traces on $H(\operatorname{div}; \Omega)$

- ▶ We denote by $H^{-1/2}(\Gamma; \mathbb{R}^d)$ the dual space of $H^{1/2}(\Gamma; \mathbb{R}^d)$ with duality pairing $\langle \cdot, \cdot \rangle_\Gamma$.
- ▶ For smooth $\tau : \overline{\Omega} \rightarrow \mathbb{S}^d$, *Green's identity* is

$$\int_{\Gamma} (\tau \mathbf{n}) \cdot (\gamma v) \, ds = \underbrace{\int_{\Omega} \tau : \nabla v \, dx + \int_{\Omega} (\operatorname{div} \tau) \cdot v \, dx}_{:= F_{\tau}(v)}$$

for all smooth $v : \overline{\Omega} \rightarrow \mathbb{S}^d$.

Normal Traces on $H(\operatorname{div}; \Omega)$

- ▶ We denote by $H^{-1/2}(\Gamma; \mathbb{R}^d)$ the dual space of $H^{1/2}(\Gamma; \mathbb{R}^d)$ with duality pairing $\langle \cdot, \cdot \rangle_\Gamma$.
- ▶ For smooth $\tau : \overline{\Omega} \rightarrow \mathbb{S}^d$, *Green's identity* is

$$\int_{\Gamma} (\tau \mathbf{n}) \cdot (\gamma v) \, ds = \underbrace{\int_{\Omega} \tau : \nabla v \, dx + \int_{\Omega} (\operatorname{div} \tau) \cdot v \, dx}_{:= F_{\tau}(v)}$$

for all smooth $v : \overline{\Omega} \rightarrow \mathbb{S}^d$.

- ▶ This means $\tau \mathbf{n}$ is defined in a weak sense via Green's identity.

Normal Traces on $H(\operatorname{div}; \Omega)$

- ▶ We denote by $H^{-1/2}(\Gamma; \mathbb{R}^d)$ the dual space of $H^{1/2}(\Gamma; \mathbb{R}^d)$ with duality pairing $\langle \cdot, \cdot \rangle_\Gamma$.
- ▶ For smooth $\tau : \overline{\Omega} \rightarrow \mathbb{S}^d$, *Green's identity* is

$$\int_{\Gamma} (\tau \mathbf{n}) \cdot (\gamma v) \, ds = \underbrace{\int_{\Omega} \tau : \nabla v \, dx + \int_{\Omega} (\operatorname{div} \tau) \cdot v \, dx}_{:= F_{\tau}(v)}$$

for all smooth $v : \overline{\Omega} \rightarrow \mathbb{S}^d$.

- ▶ This means $\tau \mathbf{n}$ is defined in a weak sense via Green's identity.
- ▶ $F_{\tau}(v)$ is still well-defined for $\tau \in H(\operatorname{div}; \Omega)$ and $v \in H^1(\Omega; \mathbb{R}^d)$.

Normal Traces on $H(\operatorname{div}; \Omega)$

- ▶ We denote by $H^{-1/2}(\Gamma; \mathbb{R}^d)$ the dual space of $H^{1/2}(\Gamma; \mathbb{R}^d)$ with duality pairing $\langle \cdot, \cdot \rangle_\Gamma$.
- ▶ For smooth $\tau : \overline{\Omega} \rightarrow \mathbb{S}^d$, *Green's identity* is

$$\int_{\Gamma} (\tau \mathbf{n}) \cdot (\gamma v) \, ds = \underbrace{\int_{\Omega} \tau : \nabla v \, dx + \int_{\Omega} (\operatorname{div} \tau) \cdot v \, dx}_{:= F_{\tau}(v)}$$

for all smooth $v : \overline{\Omega} \rightarrow \mathbb{S}^d$.

- ▶ This means $\tau \mathbf{n}$ is defined in a weak sense via Green's identity.
- ▶ $F_{\tau}(v)$ is still well-defined for $\tau \in H(\operatorname{div}; \Omega)$ and $v \in H^1(\Omega; \mathbb{R}^d)$.
- ▶ F_{τ} depends only on the trace γv and not on v itself, since it vanishes on $H_0^1(\Omega; \mathbb{R}^d)$.

Normal Traces on $H(\operatorname{div}; \Omega)$

- ▶ We denote by $H^{-1/2}(\Gamma; \mathbb{R}^d)$ the dual space of $H^{1/2}(\Gamma; \mathbb{R}^d)$ with duality pairing $\langle \cdot, \cdot \rangle_\Gamma$.
- ▶ For smooth $\tau : \overline{\Omega} \rightarrow \mathbb{S}^d$, *Green's identity* is

$$\int_{\Gamma} (\tau \mathbf{n}) \cdot (\gamma v) \, ds = \underbrace{\int_{\Omega} \tau : \nabla v \, dx + \int_{\Omega} (\operatorname{div} \tau) \cdot v \, dx}_{:= F_{\tau}(v)}$$

for all smooth $v : \overline{\Omega} \rightarrow \mathbb{S}^d$.

- ▶ This means $\tau \mathbf{n}$ is defined in a weak sense via Green's identity.
- ▶ $F_{\tau}(v)$ is still well-defined for $\tau \in H(\operatorname{div}; \Omega)$ and $v \in H^1(\Omega; \mathbb{R}^d)$.
- ▶ F_{τ} depends only on the trace γv and not on v itself, since it vanishes on $H_0^1(\Omega; \mathbb{R}^d)$.
- ▶ Therefore, F_{τ} induces a continuous linear functional on $H^{1/2}(\Gamma; \mathbb{R}^d)$:

$$\langle \gamma_{\mathbf{n}}(\tau), \gamma v \rangle_{\Gamma} := F_{\tau}(v), \quad \gamma_{\mathbf{n}}(\tau) \in H^{-1/2}(\Gamma; \mathbb{R}^d).$$

Normal Traces on $H(\operatorname{div}; \Omega)$

- ▶ We denote by $H^{-1/2}(\Gamma; \mathbb{R}^d)$ the dual space of $H^{1/2}(\Gamma; \mathbb{R}^d)$ with duality pairing $\langle \cdot, \cdot \rangle_\Gamma$.
- ▶ For smooth $\tau : \overline{\Omega} \rightarrow \mathbb{S}^d$, *Green's identity* is

$$\int_{\Gamma} (\tau \mathbf{n}) \cdot (\gamma v) \, ds = \underbrace{\int_{\Omega} \tau : \nabla v \, dx + \int_{\Omega} (\operatorname{div} \tau) \cdot v \, dx}_{:= F_{\tau}(v)}$$

for all smooth $v : \overline{\Omega} \rightarrow \mathbb{S}^d$.

- ▶ This means $\tau \mathbf{n}$ is defined in a weak sense via Green's identity.
- ▶ $F_{\tau}(v)$ is still well-defined for $\tau \in H(\operatorname{div}; \Omega)$ and $v \in H^1(\Omega; \mathbb{R}^d)$.
- ▶ F_{τ} depends only on the trace γv and not on v itself, since it vanishes on $H_0^1(\Omega; \mathbb{R}^d)$.
- ▶ Therefore, F_{τ} induces a continuous linear functional on $H^{1/2}(\Gamma; \mathbb{R}^d)$:

$$\langle \gamma_{\mathbf{n}}(\tau), \gamma v \rangle_{\Gamma} := F_{\tau}(v), \quad \gamma_{\mathbf{n}}(\tau) \in H^{-1/2}(\Gamma; \mathbb{R}^d).$$

- ▶ In the smooth case, $\gamma_{\mathbf{n}}(\tau)$ coincides with the classical traction $\tau \mathbf{n}$. We thus write $\tau \mathbf{n} = \gamma_{\mathbf{n}}(\tau)$.

Stress Regularity Assumption

- ▶ Thus, we further assume $\sigma(u) \in H(\operatorname{div}; \Omega)$ which yields $\sigma(u)_n := \gamma_n(\sigma(u)) \in H^{-1/2}(\Gamma; \mathbb{R}^d)$.

Stress Regularity Assumption

- ▶ Thus, we further assume $\sigma(u) \in H(\operatorname{div}; \Omega)$ which yields $\sigma(u)n := \gamma_n(\sigma(u)) \in H^{-1/2}(\Gamma; \mathbb{R}^d)$.
- ▶ Since $\sigma(u)$ is symmetric, we have

$$\sigma(u) : \nabla v = \sigma(u) : \varepsilon(v) \quad \forall v \in V.$$

Stress Regularity Assumption

- ▶ Thus, we further assume $\sigma(u) \in H(\operatorname{div}; \Omega)$ which yields $\sigma(u)n := \gamma_n(\sigma(u)) \in H^{-1/2}(\Gamma; \mathbb{R}^d)$.
- ▶ Since $\sigma(u)$ is symmetric, we have

$$\sigma(u) : \nabla v = \sigma(u) : \varepsilon(v) \quad \forall v \in V.$$

- ▶ Therefore

$$\langle \sigma(u)n, \gamma v \rangle_\Gamma = \int_\Omega \sigma(u) : \varepsilon(v) \, dx + \int_\Omega \operatorname{div} \sigma(u) \cdot v \, dx \quad \forall v \in V.$$

Stress Regularity Assumption

- ▶ Thus, we further assume $\sigma(u) \in H(\operatorname{div}; \Omega)$ which yields $\sigma(u)n := \gamma_n(\sigma(u)) \in H^{-1/2}(\Gamma; \mathbb{R}^d)$.
- ▶ Since $\sigma(u)$ is symmetric, we have

$$\sigma(u) : \nabla v = \sigma(u) : \varepsilon(v) \quad \forall v \in V.$$

- ▶ Therefore

$$\langle \sigma(u)n, \gamma v \rangle_\Gamma = \int_\Omega \sigma(u) : \varepsilon(v) \, dx + \int_\Omega \operatorname{div} \sigma(u) \cdot v \, dx \quad \forall v \in V.$$

- ▶ *Useful identity:* If $-\operatorname{div} \sigma(u) = f$ almost everywhere in Ω , then

$$a(u, v) - \ell(v) = \langle \sigma(u)n, \gamma v \rangle_\Gamma - \int_{\Gamma_N} F \cdot (\gamma v) \, ds, \quad \forall v \in V.$$

A Hybrid Formulation

Problem (Signorini's Problem – Hybrid Formulation)

For given $f \in L^2(\Omega; \mathbb{R}^d)$ and $F \in L^2(\Gamma_N; \mathbb{R}^d)$, find $u \in K$ such that $\sigma(u) \in H(\text{div}; \Omega)$ and

$$\begin{cases} -\text{div } \sigma(u) = f & \text{a.e. in } \Omega, \end{cases} \quad (1a)$$

$$\begin{cases} \langle \sigma(u)n, \gamma u \rangle_\Gamma = \int_{\Gamma_N} F \cdot (\gamma u) \, ds, \end{cases} \quad (1b)$$

$$\begin{cases} \langle \sigma(u)n, \gamma v \rangle_\Gamma \geq \int_{\Gamma_N} F \cdot (\gamma v) \, ds \quad \forall v \in K. \end{cases} \quad (1c)$$

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Hybrid $\Rightarrow$ Weak:

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\text{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that} \quad a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Hybrid $\Rightarrow$ Weak:

- ▶ Subtract (1b) from (1c):

$$\langle \sigma(u)\mathbf{n}, \gamma(v - u) \rangle_{\Gamma} - \int_{\Gamma_N} F \cdot \gamma(v - u) \, ds \geq 0 \quad \forall v \in K.$$

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that} \quad a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Hybrid $\Rightarrow$ Weak:

- ▶ Subtract (1b) from (1c):

$$\langle \sigma(u)n, \gamma(v - u) \rangle_{\Gamma} - \int_{\Gamma_N} F \cdot \gamma(v - u) \, ds \geq 0 \quad \forall v \in K.$$

- ▶ Insert the identity from the above:

$$a(u, v - u) - \ell(v - u) \geq 0 \quad \forall v \in K,$$

i.e. the weak formulation.

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that} \quad a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: *Weak $\Rightarrow$ Hybrid (Bulk Equation):*

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Bulk Equation):

► Let $\varphi \in \mathcal{D}(\Omega; \mathbb{R}^d)$. Then $u \pm t\varphi \in K$ for $|t| \ll 1$, hence the weak inequality implies

$$a(u, \varphi) = \ell(\varphi) = \int_{\Omega} f \cdot \varphi \, dx = \langle f, \varphi \rangle \quad \forall \varphi \in \mathcal{D}(\Omega; \mathbb{R}^d).$$

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Bulk Equation):

- ▶ Let $\varphi \in \mathcal{D}(\Omega; \mathbb{R}^d)$. Then $u \pm t\varphi \in K$ for $|t| \ll 1$, hence the weak inequality implies

$$a(u, \varphi) = \ell(\varphi) = \int_{\Omega} f \cdot \varphi \, dx = \langle f, \varphi \rangle \quad \forall \varphi \in \mathcal{D}(\Omega; \mathbb{R}^d).$$

- ▶ The left hand side is

$$a(u, \varphi) = \int_{\Omega} \sigma(u) : \varepsilon(\varphi) = \int_{\Omega} \sigma(u) : \nabla \varphi \, dx = \langle -\operatorname{div} \sigma(u), \varphi \rangle.$$

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Bulk Equation):

- ▶ Let $\varphi \in \mathcal{D}(\Omega; \mathbb{R}^d)$. Then $u \pm t\varphi \in K$ for $|t| \ll 1$, hence the weak inequality implies

$$a(u, \varphi) = \ell(\varphi) = \int_{\Omega} f \cdot \varphi \, dx = \langle f, \varphi \rangle \quad \forall \varphi \in \mathcal{D}(\Omega; \mathbb{R}^d).$$

- ▶ The left hand side is

$$a(u, \varphi) = \int_{\Omega} \sigma(u) : \varepsilon(\varphi) = \int_{\Omega} \sigma(u) : \nabla \varphi \, dx = \langle -\operatorname{div} \sigma(u), \varphi \rangle.$$

- ▶ Thus, $-\operatorname{div} \sigma(u) = f$ in $\mathcal{D}'(\Omega; \mathbb{R}^d)$ and by $\sigma(u) \in H(\operatorname{div}; \Omega)$, it holds in $L^2(\Omega; \mathbb{R}^d)$ as well.

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: *Weak $\Rightarrow$ Hybrid (Hybrid Equality):*

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Hybrid Equality):

- Since K is a cone, we have $0, 2u \in K$.

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Hybrid Equality):

- ▶ Since K is a cone, we have $0, 2u \in K$.
- ▶ Testing both in the variational inequality yields $a(u, u) \leq \ell(u)$ and $a(u, u) \geq \ell(u)$ and thus $a(u, u) = \ell(u)$.

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Hybrid Equality):

- ▶ Since K is a cone, we have $0, 2u \in K$.
- ▶ Testing both in the variational inequality yields $a(u, u) \leq \ell(u)$ and $a(u, u) \geq \ell(u)$ and thus $a(u, u) = \ell(u)$.
- ▶ The identity earlier yields

$$0 = a(u, u) - \ell(u) = \langle \sigma(u)n, \gamma u \rangle_{\Gamma} - \int_{\Gamma_N} F \cdot \gamma u \, ds.$$

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Hybrid Equality):

- ▶ Since K is a cone, we have $0, 2u \in K$.
- ▶ Testing both in the variational inequality yields $a(u, u) \leq \ell(u)$ and $a(u, u) \geq \ell(u)$ and thus $a(u, u) = \ell(u)$.
- ▶ The identity earlier yields

$$0 = a(u, u) - \ell(u) = \langle \sigma(u)n, \gamma u \rangle_{\Gamma} - \int_{\Gamma_N} F \cdot \gamma u \, ds.$$

- ▶ This is exactly (1b).

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: *Weak $\Rightarrow$ Hybrid (Hybrid Inequality):*

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Hybrid Inequality):

► With the identity earlier, we have for all $v \in V$ that

$$a(u, v - u) = \langle \sigma(u)n, \gamma(v - u) \rangle_{\Gamma} + \int_{\Omega} f \cdot (v - u) \, dx \geq \ell(v - u) = \int_{\Omega} f \cdot (v - u) \, dx + \int_{\Gamma_N} F \cdot \gamma(v - u) \, ds$$

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Hybrid Inequality):

- ▶ With the identity earlier, we have for all $v \in V$ that

$$a(u, v - u) = \langle \sigma(u)n, \gamma(v - u) \rangle_{\Gamma} + \int_{\Omega} f \cdot (v - u) \, dx \geq \ell(v - u) = \int_{\Omega} f \cdot (v - u) \, dx + \int_{\Gamma_N} F \cdot \gamma(v - u) \, ds$$

- ▶ Subtracting volume term yields: $\langle \sigma(u)n, \gamma(v - u) \rangle_{\Gamma} \geq \int_{\Gamma_N} F \cdot \gamma(v - u) \, ds \quad \forall v \in V.$

Equivalence of Hybrid and Weak Formulation

Proposition (Equivalence of Hybrid and Weak Formulation)

Assume $\sigma(u) \in H(\operatorname{div}; \Omega)$. Then $u \in K$ solves the hybrid problem above if and only if it solves the weak problem

$$\text{Find } u \in K \text{ such that } a(u, v - u) \geq \ell(v - u) \quad \forall v \in K.$$

Proof: Weak $\Rightarrow$ Hybrid (Hybrid Inequality):

- ▶ With the identity earlier, we have for all $v \in V$ that

$$a(u, v - u) = \langle \sigma(u)n, \gamma(v - u) \rangle_{\Gamma} + \int_{\Omega} f \cdot (v - u) \, dx \geq \ell(v - u) = \int_{\Omega} f \cdot (v - u) \, dx + \int_{\Gamma_N} F \cdot \gamma(v - u) \, ds$$

- ▶ Subtracting volume term yields: $\langle \sigma(u)n, \gamma(v - u) \rangle_{\Gamma} \geq \int_{\Gamma_N} F \cdot \gamma(v - u) \, ds \quad \forall v \in V.$
- ▶ This means

$$\langle \sigma(u)n, \gamma v \rangle_{\Gamma} \geq \int_{\Gamma_N} F \cdot \gamma v \, ds + \underbrace{\left[\langle \sigma(u)n, \gamma u \rangle_{\Gamma} - \int_{\Gamma_N} F \cdot \gamma u \, ds \right]}_{=0} \quad \forall v \in V.$$

Content

- 1 Introduction – What's the Problem?
- 2 Geometrical and Physical Setup
- 3 Energy Minimization
- 4 Weak Formulations
- 5 Strong Formulations**
 - A Hybrid Formulation
 - **Classical Strong Formulation**

Additional Regularity Assumptions and their Consequences

- ▶ **Goal:** Formulate classical traces for the stress $\sigma(u)$.

Additional Regularity Assumptions and their Consequences

- ▶ **Goal:** Formulate classical traces for the stress $\sigma(u)$.
- ▶ Previously, $u \in V \subset H^1(\Omega; \mathbb{R}^d)$

Additional Regularity Assumptions and their Consequences

- ▶ **Goal:** Formulate classical traces for the stress $\sigma(u)$.
- ▶ Previously, $u \in V \subset H^1(\Omega; \mathbb{R}^d)$
- ▶ In the classical strong formulation, we assume $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$.

Additional Regularity Assumptions and their Consequences

- ▶ **Goal:** Formulate classical traces for the stress $\sigma(u)$.
- ▶ Previously, $u \in V \subset H^1(\Omega; \mathbb{R}^d)$
- ▶ In the classical strong formulation, we assume $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$.
- ▶ With $\varepsilon(u) = \frac{1}{2} (\nabla u + \nabla u^\top)$, the small strain tensor is an operator $\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d)$

Additional Regularity Assumptions and their Consequences

- ▶ **Goal:** Formulate classical traces for the stress $\sigma(u)$.
- ▶ Previously, $u \in V \subset H^1(\Omega; \mathbb{R}^d)$
- ▶ In the classical strong formulation, we assume $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$.
- ▶ With $\varepsilon(u) = \frac{1}{2} (\nabla u + \nabla u^\top)$, the small strain tensor is an operator $\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d)$
- ▶ We further need the elasticity tensor field $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$ to be *sufficiently* regular (e.g. constant or $\mathcal{A}_{ijkl} \in W^{1,\infty}(\Omega)$ if $3/2 < s \leq 2$).

Additional Regularity Assumptions and their Consequences

- ▶ **Goal:** Formulate classical traces for the stress $\sigma(u)$.
- ▶ Previously, $u \in V \subset H^1(\Omega; \mathbb{R}^d)$
- ▶ In the classical strong formulation, we assume $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$.
- ▶ With $\varepsilon(u) = \frac{1}{2} (\nabla u + \nabla u^\top)$, the small strain tensor is an operator $\varepsilon : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d)$
- ▶ We further need the elasticity tensor field $\mathcal{A} : \Omega \rightarrow \mathcal{L}(\mathbb{S}^d; \mathbb{S}^d)$ to be *sufficiently* regular (e.g. constant or $\mathcal{A}_{ijkl} \in W^{1,\infty}(\Omega)$ if $3/2 < s \leq 2$).
- ▶ With $\sigma(u) = \mathcal{A} : \varepsilon(u)$, the stress tensor is now an operator $\sigma : H^s(\Omega; \mathbb{R}^d) \rightarrow H^{s-1}(\Omega; \mathbb{S}^d)$.

Additional Regularity Assumptions and their Consequences

- ▶ To apply the trace theorem on $\sigma(u)$, we further need the assumption $\Gamma \in \mathcal{C}^{m,1}$ with $m \geq [s - 1]$.

Additional Regularity Assumptions and their Consequences

- ▶ To apply the trace theorem on $\sigma(u)$, we further need the assumption $\Gamma \in \mathcal{C}^{m,1}$ with $m \geq \lceil s - 1 \rceil$.
- ▶ Then, by the trace theorem, $\gamma(\sigma(u)) \in H^{s-3/2}(\Gamma; \mathbb{S}^d)$.

Additional Regularity Assumptions and their Consequences

- ▶ To apply the trace theorem on $\sigma(u)$, we further need the assumption $\Gamma \in \mathcal{C}^{m,1}$ with $m \geq [s - 1]$.
- ▶ Then, by the trace theorem, $\gamma(\sigma(u)) \in H^{s-3/2}(\Gamma; \mathbb{S}^d)$.
- ▶ Since $\Gamma \in \mathcal{C}^{m,1}$, the unit outward normal satisfies $n \in \mathcal{C}^{m-1,1}(\Gamma; \mathbb{R}^d)$.

Additional Regularity Assumptions and their Consequences

- ▶ To apply the trace theorem on $\sigma(u)$, we further need the assumption $\Gamma \in \mathcal{C}^{m,1}$ with $m \geq [s - 1]$.
- ▶ Then, by the trace theorem, $\gamma(\sigma(u)) \in H^{s-3/2}(\Gamma; \mathbb{S}^d)$.
- ▶ Since $\Gamma \in \mathcal{C}^{m,1}$, the unit outward normal satisfies $n \in \mathcal{C}^{m-1,1}(\Gamma; \mathbb{R}^d)$.
- ▶ Hence the traction $t(u) := (\gamma\sigma(u)) n \in H^{s-3/2}(\Gamma; \mathbb{R}^d)$ is well-defined.

Additional Regularity Assumptions and their Consequences

- ▶ To apply the trace theorem on $\sigma(u)$, we further need the assumption $\Gamma \in \mathcal{C}^{m,1}$ with $m \geq [s - 1]$.
- ▶ Then, by the trace theorem, $\gamma(\sigma(u)) \in H^{s-3/2}(\Gamma; \mathbb{S}^d)$.
- ▶ Since $\Gamma \in \mathcal{C}^{m,1}$, the unit outward normal satisfies $n \in \mathcal{C}^{m-1,1}(\Gamma; \mathbb{R}^d)$.
- ▶ Hence the traction $t(u) := (\gamma\sigma(u)) n \in H^{s-3/2}(\Gamma; \mathbb{R}^d)$ is well-defined.
- ▶ If $\sigma(u) \in H(\operatorname{div}; \Omega)$, the two concepts of normal traces coincide:

$$\langle \sigma(u)n, \gamma v \rangle = \int_{\Gamma_N} t(u) \cdot (\gamma v) \, ds \quad \forall v \in V.$$

Additional Regularity Assumptions and their Consequences

- ▶ To apply the trace theorem on $\sigma(u)$, we further need the assumption $\Gamma \in \mathcal{C}^{m,1}$ with $m \geq [s - 1]$.
- ▶ Then, by the trace theorem, $\gamma(\sigma(u)) \in H^{s-3/2}(\Gamma; \mathbb{S}^d)$.
- ▶ Since $\Gamma \in \mathcal{C}^{m,1}$, the unit outward normal satisfies $n \in \mathcal{C}^{m-1,1}(\Gamma; \mathbb{R}^d)$.
- ▶ Hence the traction $t(u) := (\gamma\sigma(u)) n \in H^{s-3/2}(\Gamma; \mathbb{R}^d)$ is well-defined.
- ▶ If $\sigma(u) \in H(\operatorname{div}; \Omega)$, the two concepts of normal traces coincide:

$$\langle \sigma(u)n, \gamma v \rangle = \int_{\Gamma_N} t(u) \cdot (\gamma v) \, ds \quad \forall v \in V.$$

- ▶ Its normal and tangential components are given by

$$\sigma_n(u) := t(u) \cdot n \in H^{s-3/2}(\Gamma), \quad \sigma_t(u) := t(u) - \sigma_n(u) n \in H^{s-3/2}(\Gamma; \mathbb{R}^d).$$

Additional Regularity Assumptions and their Consequences

- ▶ To apply the trace theorem on $\sigma(u)$, we further need the assumption $\Gamma \in \mathcal{C}^{m,1}$ with $m \geq [s - 1]$.
- ▶ Then, by the trace theorem, $\gamma(\sigma(u)) \in H^{s-3/2}(\Gamma; \mathbb{S}^d)$.
- ▶ Since $\Gamma \in \mathcal{C}^{m,1}$, the unit outward normal satisfies $n \in \mathcal{C}^{m-1,1}(\Gamma; \mathbb{R}^d)$.
- ▶ Hence the traction $t(u) := (\gamma\sigma(u)) n \in H^{s-3/2}(\Gamma; \mathbb{R}^d)$ is well-defined.
- ▶ If $\sigma(u) \in H(\text{div}; \Omega)$, the two concepts of normal traces coincide:

$$\langle \sigma(u)n, \gamma v \rangle = \int_{\Gamma_N} t(u) \cdot (\gamma v) \, ds \quad \forall v \in V.$$

- ▶ Its normal and tangential components are given by

$$\sigma_n(u) := t(u) \cdot n \in H^{s-3/2}(\Gamma), \quad \sigma_t(u) := t(u) - \sigma_n(u) n \in H^{s-3/2}(\Gamma; \mathbb{R}^d).$$

- ▶ Furthermore, for $u \in H^s(\Omega; \mathbb{R}^d)$ we have $\gamma u \in (H^{s-1/2}(\Gamma; \mathbb{R}^d))$ with normal/tangential splitting

$$u_n := (\gamma u) \cdot n \in H^{s-1/2}(\Gamma) \quad \text{and} \quad u_t := \gamma u - u_n n \in H^{s-1/2}(\Gamma; \mathbb{R}^d).$$

Classical Strong Formulation under Additional Regularity

Problem (Signorini's Problem – Classical Strong Formulation)

For given $f \in L^2(\Omega; \mathbb{R}^d)$ and $F \in L^2(\Gamma_N; \mathbb{R}^d)$ find $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$ such that the small strain elasticity equations

$$\begin{cases} -\operatorname{div} \sigma(u) = f & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma_D, \\ t(u) = F & \text{on } \Gamma_N, \end{cases} \quad \begin{array}{l} (2a) \\ (2b) \\ (2c) \end{array}$$

and the contact conditions

$$\begin{cases} u_n \leq 0 & \text{on } \Gamma_C, \\ \sigma_n(u) \leq 0 & \text{on } \Gamma_C, \\ \sigma_n(u) u_n = 0 & \text{on } \Gamma_C, \\ \sigma_t(u) = 0 & \text{on } \Gamma_C \end{cases} \quad \begin{array}{l} (3a) \\ (3b) \\ (3c) \\ (3d) \end{array}$$

hold.

Interpretation of the Classical Strong Formulation

- ▶ *Equilibrium equation* – $\operatorname{div} \sigma(u) = f$ in Ω : balance of internal stresses and body forces in the bulk.

Interpretation of the Classical Strong Formulation

- ▶ *Equilibrium equation* $-\operatorname{div} \sigma(u) = f$ in Ω : balance of internal stresses and body forces in the bulk.
- ▶ *Dirichlet condition* $u = 0$ on Γ_D : the body is clamped on Γ_D .

Interpretation of the Classical Strong Formulation

- ▶ *Equilibrium equation* $-\operatorname{div} \sigma(u) = f$ in Ω : balance of internal stresses and body forces in the bulk.
- ▶ *Dirichlet condition* $u = 0$ on Γ_D : the body is clamped on Γ_D .
- ▶ *Neumann condition* $t(u) = F$ on Γ_N : boundary forces acting on the free part of the boundary.

Interpretation of the Classical Strong Formulation

- ▶ *Equilibrium equation* $-\operatorname{div} \sigma(u) = f$ in Ω : balance of internal stresses and body forces in the bulk.
- ▶ *Dirichlet condition* $u = 0$ on Γ_D : the body is clamped on Γ_D .
- ▶ *Neumann condition* $t(u) = F$ on Γ_N : boundary forces acting on the free part of the boundary.
- ▶ *Nonpenetration* $u_n \leq 0$ on Γ_C : the body cannot penetrate the rigid foundation.

Interpretation of the Classical Strong Formulation

- ▶ *Equilibrium equation* $-\operatorname{div} \sigma(u) = f$ in Ω : balance of internal stresses and body forces in the bulk.
- ▶ *Dirichlet condition* $u = 0$ on Γ_D : the body is clamped on Γ_D .
- ▶ *Neumann condition* $t(u) = F$ on Γ_N : boundary forces acting on the free part of the boundary.
- ▶ *Nonpenetration* $u_n \leq 0$ on Γ_C : the body cannot penetrate the rigid foundation.
- ▶ *Compressive contact stress* $\sigma_n(u) \leq 0$ on Γ_C : the foundation can exert only compressive normal forces.

Interpretation of the Classical Strong Formulation

- ▶ *Equilibrium equation* $-\operatorname{div} \sigma(u) = f$ in Ω : balance of internal stresses and body forces in the bulk.
- ▶ *Dirichlet condition* $u = 0$ on Γ_D : the body is clamped on Γ_D .
- ▶ *Neumann condition* $t(u) = F$ on Γ_N : boundary forces acting on the free part of the boundary.
- ▶ *Nonpenetration* $u_n \leq 0$ on Γ_C : the body cannot penetrate the rigid foundation.
- ▶ *Compressive contact stress* $\sigma_n(u) \leq 0$ on Γ_C : the foundation can exert only compressive normal forces.
- ▶ *Complementarity* $\sigma_n(u) u_n = 0$ on Γ_C : either there is no contact pressure, or the body is in contact, but not both effects independently.

Interpretation of the Classical Strong Formulation

- ▶ *Equilibrium equation* $-\operatorname{div} \sigma(u) = f$ in Ω : balance of internal stresses and body forces in the bulk.
- ▶ *Dirichlet condition* $u = 0$ on Γ_D : the body is clamped on Γ_D .
- ▶ *Neumann condition* $t(u) = F$ on Γ_N : boundary forces acting on the free part of the boundary.
- ▶ *Nonpenetration* $u_n \leq 0$ on Γ_C : the body cannot penetrate the rigid foundation.
- ▶ *Compressive contact stress* $\sigma_n(u) \leq 0$ on Γ_C : the foundation can exert only compressive normal forces.
- ▶ *Complementarity* $\sigma_n(u) u_n = 0$ on Γ_C : either there is no contact pressure, or the body is in contact, but not both effects independently.
- ▶ *Frictionless contact* $\sigma_t(u) = 0$ on Γ_C : Foundation does not exert any tangential force on the body.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid:*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Strong $\implies$ Hybrid:

Problem (Signorini's Problem – Hybrid Formulation)

For given $f \in L^2(\Omega; \mathbb{R}^d)$ and $F \in L^2(\Gamma_N; \mathbb{R}^d)$, find $u \in K$ such that $\sigma(u) \in H(\text{div}; \Omega)$ and

$$\begin{cases} -\text{div } \sigma(u) = f & \text{a.e. in } \Omega, \end{cases} \quad (4a)$$

$$\begin{cases} \langle \sigma(u)n, \gamma u \rangle_\Gamma = \int_{\Gamma_N} F \cdot (\gamma u) \, ds, \end{cases} \quad (4b)$$

$$\begin{cases} \langle \sigma(u)n, \gamma v \rangle_\Gamma \geq \int_{\Gamma_N} F \cdot (\gamma v) \, ds \quad \forall v \in K. \end{cases} \quad (4c)$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Bulk equation and $u \in K$ and $\sigma(u) \in H(\operatorname{div}; \Omega)$):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Bulk equation and $u \in K$ and $\sigma(u) \in H(\operatorname{div}; \Omega)$):*

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$ solve the strong problem.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Strong $\implies$ Hybrid (Bulk equation and $u \in K$ and $\sigma(u) \in H(\operatorname{div}; \Omega)$):

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$ solve the strong problem.
- ▶ Bulk Equation $-\operatorname{div} \sigma(u) = f$ appears in both.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Bulk equation and $u \in K$ and $\sigma(u) \in H(\operatorname{div}; \Omega)$):*

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$ solve the strong problem.
- ▶ Bulk Equation $-\operatorname{div} \sigma(u) = f$ appears in both.
- ▶ $u = 0$ on Γ_D (2b) and $u_n \leq 0$ on Γ_C (3a) are incorporated in the requirement $u \in K$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Bulk equation and $u \in K$ and $\sigma(u) \in H(\operatorname{div}; \Omega)$):*

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$ solve the strong problem.
- ▶ Bulk Equation $-\operatorname{div} \sigma(u) = f$ appears in both.
- ▶ $u = 0$ on Γ_D (2b) and $u_n \leq 0$ on Γ_C (3a) are incorporated in the requirement $u \in K$.
- ▶ Since $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$, we have

$$\sigma(u) \in H^{s-1}(\Omega; \mathbb{S}^d) \subset L^2(\Omega; \mathbb{S}^d).$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Bulk equation and $u \in K$ and $\sigma(u) \in H(\operatorname{div}; \Omega)$):*

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$ solve the strong problem.
- ▶ Bulk Equation $-\operatorname{div} \sigma(u) = f$ appears in both.
- ▶ $u = 0$ on Γ_D (2b) and $u_n \leq 0$ on Γ_C (3a) are incorporated in the requirement $u \in K$.
- ▶ Since $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$, we have

$$\sigma(u) \in H^{s-1}(\Omega; \mathbb{S}^d) \subset L^2(\Omega; \mathbb{S}^d).$$

- ▶ Since $f \in L^2(\Omega; \mathbb{R}^d)$ and $-\operatorname{div} \sigma(u) = f$, we have

$$\operatorname{div} \sigma(u) \in L^2(\Omega; \mathbb{R}^d).$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Bulk equation and $u \in K$ and $\sigma(u) \in H(\operatorname{div}; \Omega)$):*

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$ solve the strong problem.
- ▶ Bulk Equation $-\operatorname{div} \sigma(u) = f$ appears in both.
- ▶ $u = 0$ on Γ_D (2b) and $u_n \leq 0$ on Γ_C (3a) are incorporated in the requirement $u \in K$.
- ▶ Since $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$, we have

$$\sigma(u) \in H^{s-1}(\Omega; \mathbb{S}^d) \subset L^2(\Omega; \mathbb{S}^d).$$

- ▶ Since $f \in L^2(\Omega; \mathbb{R}^d)$ and $-\operatorname{div} \sigma(u) = f$, we have

$$\operatorname{div} \sigma(u) \in L^2(\Omega; \mathbb{R}^d).$$

- ▶ This means, $\sigma(u) \in H(\operatorname{div}; \Omega)$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Hybrid Inequality):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Strong $\implies$ Hybrid (Hybrid Inequality):

► We want to show

$$\langle \sigma(u)n, \gamma v \rangle_\Gamma = \int_\Gamma t(u) \cdot (\gamma v) \, ds \geq \int_{\Gamma_N} F \cdot (\gamma v) \, ds \quad \forall v \in K.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Hybrid Inequality):*

► We want to show

$$\langle \sigma(u)n, \gamma v \rangle_\Gamma = \int_\Gamma t(u) \cdot (\gamma v) \, ds \geq \int_{\Gamma_N} F \cdot (\gamma v) \, ds \quad \forall v \in K.$$

► We decompose

$$\int_\Gamma t(u) \cdot (\gamma v) \, ds = \underbrace{\int_{\Gamma_D} t(u) \cdot (\gamma v) \, ds}_{=0 \text{ since } \gamma v=0 \text{ on } \Gamma_D} + \int_{\Gamma_N} \underbrace{t(u)}_{=F \text{ on } \Gamma_C} \cdot (\gamma v) \, ds + \int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Strong $\implies$ Hybrid (Hybrid Inequality):

► We want to show

$$\langle \sigma(u)n, \gamma v \rangle_\Gamma = \int_\Gamma t(u) \cdot (\gamma v) \, ds \geq \int_{\Gamma_N} F \cdot (\gamma v) \, ds \quad \forall v \in K.$$

► We decompose

$$\int_\Gamma t(u) \cdot (\gamma v) \, ds = \underbrace{\int_{\Gamma_D} t(u) \cdot (\gamma v) \, ds}_{=0 \text{ since } \gamma v=0 \text{ on } \Gamma_D} + \int_{\Gamma_N} \underbrace{t(u)}_{=F \text{ on } \Gamma_C} \cdot (\gamma v) \, ds + \int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds.$$

► With $t(u) = \sigma_n(u)n + \sigma_t(u)$ and $\gamma v = v_n n + v_t$, we have

$$t(u) \cdot (\gamma v) = \sigma_n(u) v_n + \underbrace{\sigma_t(u) \cdot v_t}_{=0 \text{ on } \Gamma_C \text{ since } \sigma_t(u)=0 \text{ on } \Gamma_C}.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Hybrid Inequality):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Strong $\implies$ Hybrid (Hybrid Inequality):

► In total:

$$\langle \sigma(u)n, \gamma v \rangle_{\Gamma} = \int_{\Gamma_N} F \cdot (\gamma v) \, ds + \int_{\Gamma_C} \sigma_n(u) v_n \, ds \quad \forall v \in K.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Strong $\implies$ Hybrid (Hybrid Inequality):

- In total:

$$\langle \sigma(u)n, \gamma v \rangle_\Gamma = \int_{\Gamma_N} F \cdot (\gamma v) \, ds + \int_{\Gamma_C} \sigma_n(u) v_n \, ds \quad \forall v \in K.$$

- Since $v \in K$, we have $v_n \leq 0$ on Γ_C and by (3b), we have $\sigma_n(u) \leq 0$ on Γ_C .

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Hybrid Inequality):*

- In total:

$$\langle \sigma(u)n, \gamma v \rangle_\Gamma = \int_{\Gamma_N} F \cdot (\gamma v) \, ds + \int_{\Gamma_C} \sigma_n(u) v_n \, ds \quad \forall v \in K.$$

- Since $v \in K$, we have $v_n \leq 0$ on Γ_C and by (3b), we have $\sigma_n(u) \leq 0$ on Γ_C .
- Thus $\sigma_n(u) v_n \geq 0$ on Γ_C and thus

$$\langle \sigma(u)n, \gamma v \rangle_\Gamma \geq \int_{\Gamma_N} F \cdot (\gamma v) \, ds \quad \forall v \in V.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Hybrid Equality):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Strong $\implies$ Hybrid (Hybrid Equality):*

- ▶ Same as before but with $v = u$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Strong $\implies$ Hybrid (Hybrid Equality):

- ▶ Same as before but with $v = u$.
- ▶ This is

$$\begin{aligned} \langle \sigma(u)n, \gamma u \rangle_{\Gamma} &= \int_{\Gamma_N} F \cdot (\gamma u) \, ds + \underbrace{\int_{\Gamma_C} \sigma_n(u) u_n \, ds}_{=0 \text{ since } \sigma_n(u) u_n = 0 \text{ on } \Gamma_C} . \end{aligned}$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong (Bulk Equation, $u = 0$ on Γ_D and $u_n \leq 0$ on Γ_C):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong (Bulk Equation, $u = 0$ on Γ_D and $u_n \leq 0$ on Γ_C):

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$, $s > 3/2$ solve the hybrid problem.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong (Bulk Equation, $u = 0$ on Γ_D and $u_n \leq 0$ on Γ_C):

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$, $s > 3/2$ solve the hybrid problem.
- ▶ Bulk equation $-\operatorname{div} \sigma(u) = f$ is contained in both problems.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong (Bulk Equation, $u = 0$ on Γ_D and $u_n \leq 0$ on Γ_C):

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$, $s > 3/2$ solve the hybrid problem.
- ▶ Bulk equation $-\operatorname{div} \sigma(u) = f$ is contained in both problems.
- ▶ Since $u \in K$, we have $u = 0$ on Γ_D and $u_n \leq 0$ on Γ_C .

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong (Bulk Equation, $u = 0$ on Γ_D and $u_n \leq 0$ on Γ_C):*

- ▶ Let $u \in H^s(\Omega; \mathbb{R}^d)$, $s > 3/2$ solve the hybrid problem.
- ▶ Bulk equation $-\operatorname{div} \sigma(u) = f$ is contained in both problems.
- ▶ Since $u \in K$, we have $u = 0$ on Γ_D and $u_n \leq 0$ on Γ_C .
- ▶ It remains to prove $t(u) = F$ on Γ_N and the remaining contact conditions.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong ($t(u) = F$ on Γ_N):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong ($t(u) = F$ on Γ_N):

- ▶ Let $g \in H^{1/2}(\Gamma_N; \mathbb{R}^d)$ such that $\text{Ext}_0(g) \in H^{1/2}(\Gamma; \mathbb{R}^d)$, i.e. $g \in H_{00}^{1/2}(\Gamma_N; \mathbb{R}^d)$. Then there exists $v \in H^1(\Omega; \mathbb{R}^d)$ such that

$$(\gamma v)|_{\Gamma_N} = g, \quad (\gamma v)|_{\Gamma_C} = 0, \quad (\gamma v)|_{\Gamma_D} = 0.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong ($t(u) = F$ on Γ_N):

- ▶ Let $g \in H^{1/2}(\Gamma_N; \mathbb{R}^d)$ such that $\text{Ext}_0(g) \in H^{1/2}(\Gamma; \mathbb{R}^d)$, i.e. $g \in H_{00}^{1/2}(\Gamma_N; \mathbb{R}^d)$. Then there exists $v \in H^1(\Omega; \mathbb{R}^d)$ such that

$$(\gamma v)|_{\Gamma_N} = g, \quad (\gamma v)|_{\Gamma_C} = 0, \quad (\gamma v)|_{\Gamma_D} = 0.$$

- ▶ Then $\pm v \in K$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong* ($t(u) = F$ on Γ_N):

- ▶ Let $g \in H^{1/2}(\Gamma_N; \mathbb{R}^d)$ such that $\text{Ext}_0(g) \in H^{1/2}(\Gamma; \mathbb{R}^d)$, i.e. $g \in H_{00}^{1/2}(\Gamma_N; \mathbb{R}^d)$. Then there exists $v \in H^1(\Omega; \mathbb{R}^d)$ such that

$$(\gamma v)|_{\Gamma_N} = g, \quad (\gamma v)|_{\Gamma_C} = 0, \quad (\gamma v)|_{\Gamma_D} = 0.$$

- ▶ Then $\pm v \in K$.
- ▶ Plug $\pm v$ into hybrid inequality:

$$\int_{\Gamma_N} t(u) \cdot g \, ds \geq \int_{\Gamma_N} F \cdot g \, ds, \quad - \int_{\Gamma_N} t(u) \cdot g \, ds \geq - \int_{\Gamma_N} F \cdot f \, ds.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong ($t(u) = F$ on Γ_N):

- ▶ Let $g \in H^{1/2}(\Gamma_N; \mathbb{R}^d)$ such that $\text{Ext}_0(g) \in H^{1/2}(\Gamma; \mathbb{R}^d)$, i.e. $g \in H_{00}^{1/2}(\Gamma_N; \mathbb{R}^d)$. Then there exists $v \in H^1(\Omega; \mathbb{R}^d)$ such that

$$(\gamma v)|_{\Gamma_N} = g, \quad (\gamma v)|_{\Gamma_C} = 0, \quad (\gamma v)|_{\Gamma_D} = 0.$$

- ▶ Then $\pm v \in K$.
- ▶ Plug $\pm v$ into hybrid inequality:

$$\int_{\Gamma_N} t(u) \cdot g \, ds \geq \int_{\Gamma_N} F \cdot g \, ds, \quad - \int_{\Gamma_N} t(u) \cdot g \, ds \geq - \int_{\Gamma_N} F \cdot g \, ds.$$

- ▶ Therefore,

$$\int_{\Gamma_N} (t(u) - F) \cdot g \, ds = 0 \quad \forall g \in H_{00}^{1/2}(\Gamma_N; \mathbb{R}^d) \quad \implies \quad t(u) = F \text{ on } \Gamma_N.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong ($\sigma_t(u) = 0$ on Γ_C):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong ($\sigma_t(u) = 0$ on Γ_C):

- ▶ With $t(u) = F$ on Γ_N , the hybrid inequality yields $\int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds \geq 0$ for all $v \in K$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong ($\sigma_t(u) = 0$ on Γ_C):

- ▶ With $t(u) = F$ on Γ_N , the hybrid inequality yields $\int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds \geq 0$ for all $v \in K$.
- ▶ Let $g_t \in H_{00}^{1/2}(\Gamma_C; \mathbb{R}^d)$ with $g_t \cdot \mathbf{n} = 0$, and choose $v \in H^1(\Omega; \mathbb{R}^d)$ with $(\gamma v)|_{\Gamma_C} = g_t$ and $(\gamma v)|_{\Gamma_N} = 0$ and $(\gamma v)|_{\Gamma_D} = 0$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong ($\sigma_t(u) = 0$ on Γ_C):

- ▶ With $t(u) = F$ on Γ_N , the hybrid inequality yields $\int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds \geq 0$ for all $v \in K$.
- ▶ Let $g_t \in H_{00}^{1/2}(\Gamma_C; \mathbb{R}^d)$ with $g_t \cdot \mathbf{n} = 0$, and choose $v \in H^1(\Omega; \mathbb{R}^d)$ with $(\gamma v)|_{\Gamma_C} = g_t$ and $(\gamma v)|_{\Gamma_N} = 0$ and $(\gamma v)|_{\Gamma_D} = 0$.
- ▶ Then $\pm v \in K$, hence applying hybrid inequality to $\pm v$ yields

$$\int_{\Gamma_C} t(u) \cdot g_t \, ds \geq 0, \quad - \int_{\Gamma_C} t(u) \cdot g_t \, ds \geq 0,$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong ($\sigma_t(u) = 0$ on Γ_C):

- ▶ With $t(u) = F$ on Γ_N , the hybrid inequality yields $\int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds \geq 0$ for all $v \in K$.
- ▶ Let $g_t \in H_{00}^{1/2}(\Gamma_C; \mathbb{R}^d)$ with $g_t \cdot \mathbf{n} = 0$, and choose $v \in H^1(\Omega; \mathbb{R}^d)$ with $(\gamma v)|_{\Gamma_C} = g_t$ and $(\gamma v)|_{\Gamma_N} = 0$ and $(\gamma v)|_{\Gamma_D} = 0$.
- ▶ Then $\pm v \in K$, hence applying hybrid inequality to $\pm v$ yields

$$\int_{\Gamma_C} t(u) \cdot g_t \, ds \geq 0, \quad - \int_{\Gamma_C} t(u) \cdot g_t \, ds \geq 0,$$

- ▶ With $t(u) = \sigma_n(u)\mathbf{n} + \sigma_t(u)$ this means

$$\int_{\Gamma_C} (\sigma_n(u)\mathbf{n} + \sigma_t(u)) \cdot g_t \, ds = \int_{\Gamma_C} \sigma_n(u) \underbrace{(\mathbf{n} \cdot g_t)}_{=0} \, ds + \int_{\Gamma_C} \sigma_t(u) \cdot g_t \, ds = 0.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong ($\sigma_n(u) \leq 0 = 0$ on Γ_C):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong ($\sigma_n(u) \leq 0 = 0$ on Γ_C):*

- ▶ Again, $\int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds \geq 0$ for all $v \in K$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: Hybrid $\implies$ Strong ($\sigma_n(u) \leq 0 = 0$ on Γ_C):

- ▶ Again, $\int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds \geq 0$ for all $v \in K$.
- ▶ Let $\eta \in H_{00}^{1/2}(\Gamma_C)$ with $\eta \leq 0$ on Γ_C and choose $v \in H^1(\Omega; \mathbb{R}^d)$ with $(\gamma v)|_{\Gamma_C} = \eta n$ and $(\gamma v)|_{\Gamma_N} = 0$ and $(\gamma v)|_{\Gamma_D} = 0$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong* ($\sigma_n(u) \leq 0 = 0$ on Γ_C):

- ▶ Again, $\int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds \geq 0$ for all $v \in K$.
- ▶ Let $\eta \in H_{00}^{1/2}(\Gamma_C)$ with $\eta \leq 0$ on Γ_C and choose $v \in H^1(\Omega; \mathbb{R}^d)$ with $(\gamma v)|_{\Gamma_C} = \eta \mathbf{n}$ and $(\gamma v)|_{\Gamma_N} = 0$ and $(\gamma v)|_{\Gamma_D} = 0$.
- ▶ Then, $v_n = (\gamma v) \cdot \mathbf{n} = \eta \leq 0$, i.e. $v \in K$ and $\int_{\Gamma_C} t(u) \cdot (\eta \mathbf{n}) \, ds \geq 0$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong* ($\sigma_n(u) \leq 0 = 0$ on Γ_C):

- ▶ Again, $\int_{\Gamma_C} t(u) \cdot (\gamma v) \, ds \geq 0$ for all $v \in K$.
- ▶ Let $\eta \in H_{00}^{1/2}(\Gamma_C)$ with $\eta \leq 0$ on Γ_C and choose $v \in H^1(\Omega; \mathbb{R}^d)$ with $(\gamma v)|_{\Gamma_C} = \eta \mathbf{n}$ and $(\gamma v)|_{\Gamma_N} = 0$ and $(\gamma v)|_{\Gamma_D} = 0$.
- ▶ Then, $v_n = (\gamma v) \cdot \mathbf{n} = \eta \leq 0$, i.e. $v \in K$ and $\int_{\Gamma_C} t(u) \cdot (\eta \mathbf{n}) \, ds \geq 0$.
- ▶ With $t(u) = \sigma_n(u) \mathbf{n} + \sigma_t(u)$ and $\sigma_t(u) = 0$ on Γ_C , we have

$$\int_{\Gamma_C} \sigma_n(u) \eta \, ds \geq 0 \quad \forall \eta \in H_{00}^{1/2}(\Gamma_C), \eta \leq 0.$$

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong ($\sigma_n(u) u_n = 0$ on Γ_C):*

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong ($\sigma_n(u) u_n = 0$ on Γ_C):*

- ▶ Hybrid equality yields $\int_{\Gamma_C} t(u) \cdot \gamma u \, ds = 0$.

Proposition (Equivalence of Strong and Hybrid Formulation)

Let $u \in H^s(\Omega; \mathbb{R}^d)$ with $s > 3/2$. Then u solves the strong problem if and only if it solves the hybrid problem.

Proof: *Hybrid $\implies$ Strong* ($\sigma_n(u) u_n = 0$ on Γ_C):

- ▶ Hybrid equality yields $\int_{\Gamma_C} t(u) \cdot \gamma u \, ds = 0$.
- ▶ With $t(u) = \sigma_n(u) n + \sigma_t(u)$ and $\sigma_t(u) = 0$ on Γ_C , we have

$$\int_{\Gamma_C} t(u) \cdot \gamma u \, ds = \int_{\Gamma_C} \underbrace{\sigma_n(u)}_{\leq 0} \underbrace{u_n}_{\leq 0} \, ds = 0.$$

□