

Formula Page

Algebra of set theory

Union: $C = A \cup B$; **Intersection:** $C = A \cap B$; **Complement:** $C = A^c$

Commutative Laws: $A \cup B = B \cup A$, $A \cap B = B \cap A$

Associative Laws: $(A \cup B) \cup C = A \cup (B \cup C)$, $(A \cap B) \cap C = A \cap (B \cap C)$

Distributive Laws: $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$, $(A \cap B) \cup C = (A \cup C) \cap (B \cup C)$

De Morgan's Law: $(A \cap B)^c = A^c \cup B^c$, $(A \cup B)^c = A^c \cap B^c$

Computing probabilities

k samples from n objects

	ordered	unordered
replacement	n^k	$\frac{(n+k-1)!}{(n-1)! k!}$
no replacement	$\frac{n!}{(n-k)!}$	$\frac{n!}{k! (n-k)!}$

Conditional probabilities, Bayes' Theorem, Law of total probability

Addition Law: $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

Multiplication Law: $P(A \cap B) = P(A|B)P(B)$

Bayes' theorem: $P(B|A) = \frac{P(A|B)P(B)}{P(A)}$

Law of total probability: $P(A) = P(A|B_1)P(B_1) + P(A|B_2)P(B_2) + \dots + P(A|B_n)P(B_n)$

Type of probabilities:

Events	B_1	...	B_n
Prior probabilities	$P(B_1)$...	$P(B_n)$
Conditional probabilities	$P(A B_1)$...	$P(A B_n)$
Joint probabilities	$P(A B_1)P(B_1)$, $P(A \cap B_1)$...	$P(A B_n)P(B_n)$, $P(A \cap B_n)$
Posterior probabilities	$P(B_1 A)$...	$P(B_n A)$

Independence

Pairwise independence \nRightarrow Mutual independence

Discrete rv

$$E(X) = \sum_i x_i \cdot p_X(x_i) = \int_{-\infty}^{+\infty} x f_X(x) dx$$

$$Y = f(X), E(Y) = \sum_i f(x_i) \cdot p_X(x_i)$$

$$E(cX + d) = cE(X) + d$$

$$\text{Var}(X) = E[(X - \mu)^2] = \sum_i (x_i - \mu)^2 p_X(x_i) = \int_{-\infty}^{+\infty} (x - \mu)^2 f_X(x) dx = E(X^2) - [E(X)]^2$$

$$\text{Var}(cX + d) = c^2 \text{Var}(X)$$

Bernoulli distribution: X takes on only two values e.g. 0 and 1, with probabilities $1 - p$ and p .

$$p(x) = \begin{cases} p^x(1-p)^{1-x}, & \text{if } x = 0 \text{ or } x = 1 \\ 0, & \text{otherwise} \end{cases}, \text{Mean: } p, \text{Variance: } p(1-p)$$

Binomial distribution: X is the sum of n variables that follow Bernoulli distribution.

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}, \text{Mean: } np, \text{Variance: } np(1-p)$$

Limit: Poisson Distribution, $n \rightarrow \infty, p \rightarrow 0, np \rightarrow \infty, X \sim \text{Poisson}(np)$

Geometric distribution (total number of trials): X is the total trials until the first success.

$$P(X = k) = (1-p)^{k-1} p, \text{Mean: } 1/p, \text{Variance: } (1-p)/p^2$$

Geometric distribution (number of failures): X is the total fail trials until the first success.

$P(X = k) = (1 - p)^k p$, Mean: $(1 - p)/p$, Variance: $(1 - p)/p^2$

Poisson Distribution: X is the number of times an event occurs in a given interval of time.

$P(X = k) = \frac{\lambda^k}{k!} e^{-\lambda}$, Mean: λ , Variance: λ

Limit: Normal Distribution, $\lambda \rightarrow \infty$, $X \sim N(\lambda, \lambda)$

Continuous rv

pth quantile x_p : $F(x_p) = P(X \leq x_p) = p$

Uniform Distribution:

$$\text{pdf: } f_X(x) = \begin{cases} \frac{1}{b-a}, & a \leq x \leq b \\ 0, & x < a \text{ or } x > b \end{cases}, \text{ cdf: } F_X(x) = \begin{cases} 0, & x \leq a \\ \frac{x-a}{b-a}, & a < x < b \\ 1, & x \geq b \end{cases}, \text{ Mean: } \frac{a+b}{2}, \text{ Variance: } \frac{(b-a)^2}{12}$$

Normal distribution: $X \sim N(\mu, \sigma^2)$, pdf: $f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\}$, Mean: μ , Variance: σ^2

$$f(x, y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left[-\frac{1}{2(1-\rho^2)}\left\{\frac{(x-\mu_X)^2}{\sigma_X^2} + \frac{(y-\mu_Y)^2}{\sigma_Y^2} - \frac{2\rho(x-\mu_X)(y-\mu_Y)}{\sigma_X\sigma_Y}\right\}\right]$$

$$(X, Y) \sim N_2(\boldsymbol{\mu}, \Sigma) \Rightarrow \nRightarrow X \sim N(\mu_X, \sigma_X^2) \text{ and } Y \sim N(\mu_Y, \sigma_Y^2)$$

Exponential distribution: the time until next event. λ : count per time unit

pdf: $f_X(x) = \lambda \exp(-\lambda x)$, $x \geq 0, \lambda > 0$, cdf: $F_X(x) = 1 - \exp(-\lambda x)$, $x \geq 0$

Mean: $1/\lambda$, Variance: $1/\lambda^2$

χ^2 distribution: pdf: $f_X(x) = \frac{1}{\Gamma(p/2)2^{p/2}} x^{p/2-1} e^{-x/2}$, $x > 0$; $Z \sim N(0, 1)$, $Z^2 \sim \chi_1^2$

Joint Distribution

$$f_X(x) = \int_{-\infty}^{+\infty} f(x, y) dy$$

$$P(x_1 < X \leq x_2, y_1 < Y \leq y_2) = F(x_2, y_2) - F(x_2, y_1) - F(x_1, y_2) + F(x_1, y_1)$$

Covariance

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y)$$

$$\text{Cov}(aW + bX, cY + dZ) = ac\text{Cov}(W, Y) + ad\text{Cov}(W, Z) + bc\text{Cov}(X, Y) + bd\text{Cov}(X, Z)$$

$$\text{Var}(aX + bY) = a^2\text{Var}(X) + b^2\text{Var}(Y) + 2ab\text{Cov}(X, Y)$$

$$\text{independent} \Rightarrow \nRightarrow \rho(X, Y) = 0$$

Limiting Theorems

Chebyshev's Inequality:

Let X be a random variable with known $E(X)$ and $\text{Var}(X) = \sigma^2$ but unknown distribution

$$P(|X - E(X)| \geq k\sigma) \leq \frac{1}{k^2}$$

Law of Large numbers:

The sample mean will be close to μ if the sample size is sufficiently large.

Central Limit Theorem:

Let X_1, X_2, \dots, X_n be a sequence of independent random variables having a common distribution with mean μ and variance σ^2 . We can compute probabilities for the sample mean using the cdf of the Normal distribution:

$$\bar{X} \sim N(\mu, \sigma^2/n)$$