

Groups, Analysis, and Geometry Seminars:

Harmonic Analysis of $SU(2)$

Tim Gou

20th August, 2020

Suppose f is 2π -periodic, complex valued, integrable over $[0, 2\pi)$, then

$$\widehat{f}(n) = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-inx} dx \quad (1)$$

with $n \in \mathbb{Z}$, is the Fourier transform of f . Question: What is e^{inx} ? Why is $n \in \mathbb{Z}$? Answer: e^{inx} is the character of circle group, denoted by \mathbb{T} (i.e. $e^{ix} \in \mathbb{T}$).

A character is a continuous homomorphism from a locally compact Abelian group G to \mathbb{T} : $\chi : G \rightarrow \mathbb{T}$ where

$$\chi(gh) = \chi(g)\chi(h) \quad (2)$$

for $g, h \in G$. Let's work out $\chi : \mathbb{R} \rightarrow \mathbb{T}$ first, where \mathbb{R} is the group $(\mathbb{R}, +)$. Since $\chi(0) = 1$ (identity to identity) and χ is continuous, then $\exists a > 0$ such that

$$\int_0^a \chi(y) dy. \quad (3)$$

Let $\xi = \int_0^a \chi(y) dy$, then

$$\chi(x)\xi = \int_0^a \chi(x+y) dy = \int_x^{a+x} \chi(t) dt \quad (4)$$

so

$$\chi(x) = \xi^{-1} \int_x^{a+x} \chi(t) dt \quad (5)$$

and

$$\begin{aligned} \chi'(x) &= \xi^{-1} (\chi(a+x) - \chi(x)) \\ &= \xi^{-1} \chi(x) (\chi(a) - 1) \\ &= c\chi(x). \end{aligned} \quad (6)$$

We have an ODE

$$\chi'(x) = c\chi(x) \quad (7)$$

where solving the equation gives us

$$\chi(x) = e^{cx}. \quad (8)$$

Solving it gives us $\chi(x) = e^{cx}$. Since $|\chi| = 1$, then $c = i\lambda$ with $\lambda \in \mathbb{R}$. Thus $\chi(x) = e^{i\lambda x}$ and we have characters of \mathbb{R} , all the χ_λ form a dual group of \mathbb{R} , denoted by $\widehat{\mathbb{R}}$.

Since we identify each χ_λ with $\lambda \in \mathbb{R}$, then

$$\widehat{\mathbb{R}} \cong \mathbb{R}. \quad (9)$$

To work out $\widehat{\mathbb{T}}$, notice that

$$\mathbb{T} \cong \mathbb{R}/2\pi\mathbb{Z} \quad (10)$$

i.e. each element in $[0, 2\pi)$ is a representative of the cosets of $\mathbb{R}/2\pi\mathbb{Z}$. Suppose $x, y \in \mathbb{R}/2\pi\mathbb{Z}$ and $x + y = 2\pi$, then

$$\chi(x + y) = \chi(0) = 1 = e^{i\lambda(x+y)} \quad (11)$$

we know $\lambda \in \mathbb{R}$, but the only way $e^{i\lambda(x+y)} = 1$ is that $\lambda \in \mathbb{Z}$. So all the $\chi_n(x) = e^{inx}$ for the dual group $\widehat{\mathbb{T}}$, and

$$\widehat{\mathbb{T}} \cong \mathbb{Z}. \quad (12)$$

Similarly, we have $\mathbb{R}^n \cong \mathbb{R}^n$ and $\widehat{\mathbb{T}} \cong \mathbb{Z}^n$.

Theorem 0.1: If G is compact, \widehat{G} is discrete.

In addition, $\{e^{inx} : n \in \mathbb{Z}\}$ form an orthonormal basis for the Hilbert space $L^2(\mathbb{T})$, with respect to its inner product, i.e.

$$\begin{aligned} \langle e^{imx}, e^{inx} \rangle &= \frac{1}{2\pi} \int_0^{2\pi} e^{imx} e^{-inx} dx \\ &= \delta_{mn} = \begin{cases} 1, & m = n \\ 0, & m \neq n \end{cases} \end{aligned} \quad (13)$$