

Deriving syntax and axioms for quantitative regular behaviours

Filippo Bonchi¹ Marcello Bonsangue^{1,2} Jan Rutten^{1,3}
Alexandra Silva¹

¹Centrum voor Wiskunde en Informatica

²LIACS - Leiden University

³Vrije Universiteit Amsterdam


CONCUR, September 2009

Specify and reason about systems.

Motivation

Specify and reason about systems.

state-machines
e.g. DFA, LTS, PA,



Motivation

Specify and reason about systems.


Syntax
RE, CCS, ...

$b^*a(b^*a)^*$

$a.b.0 + a.c.0$

$a(\frac{1}{2}.0 \oplus \frac{1}{2}.0) + \dots$

state-machines
e.g. DFA, LTS, PA,



Motivation

Specify

and

reason

about

systems.

Syntax

RE, CCS, ...

$$b^*a(b^*a)^*$$

$$a.b.0 + a.c.0$$

$$a.(\frac{1}{2}.0 \oplus \frac{1}{2}.0) + \dots$$

Axiomatization

KA, ...


$$1 + a.a^* = a^*$$

$$P + 0 = P$$

$$p.P \oplus p'.P = (p+p').P$$

state-machines

e.g. DFA, LTS, PA,



Motivation

Specify and reason about systems.

Syntax
RE, CCS, ...

$$b^*a(b^*a)^*$$

$$a.b.0 + a.c.0$$

$$a.(\frac{1}{2}.0 \oplus \frac{1}{2}.0) + \dots$$


Axiomatization
KA, ...

$$1 + a.a^* = a^*$$

$$P + 0 = P$$


$$p.P \oplus p'.P = (p+p').P$$

state-machines
e.g. DFA, LTS, PA,



Can we do all of this **uniformly** in a single framework?

What do these things have in common?



$$(S, t: S \rightarrow 2 \times S^A)$$

$$(S, t: S \rightarrow \mathcal{P}S^A)$$


$$(S, t: S \rightarrow \mathcal{P}\mathcal{D}_\omega(S)^A)$$

$$(S, t: S \rightarrow \mathcal{D}_\omega(S) + (A \times S) + 1)$$


$$(S, t: S \rightarrow \mathcal{P}(\mathcal{D}_\omega(\mathcal{P}S^A)))$$

$$(S, t: S \rightarrow GS)$$


G-coalgebras




What do these things have in common?




$$(S, t: S \rightarrow 2 \times S^A)$$




$$(S, t: S \rightarrow \mathcal{P}S^A)$$



$$(S, t: S \rightarrow \mathcal{PD}_\omega(S)^A)$$



$$(S, t: S \rightarrow \mathcal{D}_\omega(S) + (A \times S) + 1)$$




$$(S, t: S \rightarrow \mathcal{P}(\mathcal{D}_\omega(\mathcal{P}S^A)))$$


$$(S, t: S \rightarrow GS)$$

G-coalgebras


What do these things have in common?




$$(S, t: S \rightarrow 2 \times S^A)$$




$$(S, t: S \rightarrow \mathcal{P}S^A)$$



$$(S, t: S \rightarrow \mathcal{PD}_\omega(S)^A)$$



$$(S, t: S \rightarrow \mathcal{D}_\omega(S) + (A \times S) + 1)$$




$$(S, t: S \rightarrow \mathcal{P}(\mathcal{D}_\omega(\mathcal{P}S^A)))$$


$$(S, t: S \rightarrow GS)$$

G-coalgebras

What do these things have in common?




$$(S, t: S \rightarrow 2 \times S^A)$$




$$(S, t: S \rightarrow \mathcal{P}S^A)$$



$$(S, t: S \rightarrow \mathcal{PD}_\omega(S)^A)$$



$$(S, t: S \rightarrow \mathcal{D}_\omega(S) + (A \times S) + 1)$$




$$(S, t: S \rightarrow \mathcal{P}(\mathcal{D}_\omega(\mathcal{P}S^A)))$$


$$(S, t: S \rightarrow GS)$$

G-coalgebras


What do these things have in common?




$$(S, t: S \rightarrow 2 \times S^A)$$




$$(S, t: S \rightarrow \mathcal{P}S^A)$$



$$(S, t: S \rightarrow \mathcal{PD}_\omega(S)^A)$$



$$(S, t: S \rightarrow \mathcal{D}_\omega(S) + (A \times S) + 1)$$




$$(S, t: S \rightarrow \mathcal{P}(\mathcal{D}_\omega(\mathcal{P}S)^A))$$


$$(S, t: S \rightarrow GS)$$

G-coalgebras


What do these things have in common?




$$(S, t: S \rightarrow 2 \times S^A)$$




$$(S, t: S \rightarrow \mathcal{P}S^A)$$



$$(S, t: S \rightarrow \mathcal{PD}_\omega(S)^A)$$



$$(S, t: S \rightarrow \mathcal{D}_\omega(S) + (A \times S) + 1)$$




$$(S, t: S \rightarrow \mathcal{P}(\mathcal{D}_\omega(\mathcal{P}S)^A))$$


$$(S, t: S \rightarrow GS)$$

G-coalgebras


What do these things have in common?




$$(S, t: S \rightarrow 2 \times S^A)$$




$$(S, t: S \rightarrow \mathcal{P}S^A)$$



$$(S, t: S \rightarrow \mathcal{PD}_\omega(S)^A)$$



$$(S, t: S \rightarrow \mathcal{D}_\omega(S) + (A \times S) + 1)$$



$$(S, t: S \rightarrow \mathcal{P}(\mathcal{D}_\omega(\mathcal{P}S^A)))$$

$$(S, t: S \rightarrow GS)$$

G-coalgebras

The power of G

$$(S, t:S \rightarrow GS)$$

The functor G determines:

- 1 notion of observational equivalence (coalg. bisimulation)
- 2 behaviour (final coalgebra)
- 3 set of expressions describing finite systems
- 4 axioms to prove bisimulation equivalence of expressions

The power of G

$$(S, t:S \rightarrow GS)$$

The functor G determines:

- 1 notion of observational equivalence (coalg. bisimulation)
- 2 behaviour (final coalgebra)
- 3 set of expressions describing finite systems
- 4 axioms to prove bisimulation equivalence of expressions

The power of G

$$(S, t: S \rightarrow GS)$$

The functor G determines:

- ❶ notion of observational equivalence (coalg. bisimulation)
- ❷ behaviour (final coalgebra)
- ❸ set of expressions describing finite systems
- ❹ axioms to prove bisimulation equivalence of expressions

❶ + ❷ are classic coalgebra; ❸ + ❹ are LICS'09 and CONCUR'09

Coalgebras

Quantitative coalgebras

- Generalizations of deterministic automata
- Quantitative coalgebras: set of states S and $t : S \rightarrow GS$

$$G ::= Id \mid B \mid G \times G \mid G + G \mid G^A \mid \mathbb{M}^G$$

\mathbb{M} is a monoid. $\mathcal{P} = 2^{Id}$ and $\mathcal{D}_\omega = \mathbb{R}^{Id}$

Examples

- $G = 2 \times Id^A$ Deterministic automata
- $G = (B \times Id)^A$ Mealy machines
- $G = (\mathcal{P}Id)^A$ LTS
- $G = \mathcal{PD}_\omega(S)^A$ Simple Segala systems
- ...

Quantitative coalgebras

- Generalizations of deterministic automata
- Quantitative coalgebras: set of states S and $t : S \rightarrow GS$

$$G ::= Id \mid B \mid G \times G \mid G + G \mid G^A \mid \mathbb{M}^G$$

\mathbb{M} is a monoid. $\mathcal{P} = 2^{Id}$ and $\mathcal{D}_\omega = \mathbb{R}^{Id}$

Examples

- $G = 2 \times Id^A$ Deterministic automata
- $G = (B \times Id)^A$ Mealy machines
- $G = (\mathcal{P}Id)^A$ LTS
- $G = \mathcal{PD}_\omega(S)^A$ Simple Segala systems
- ...

Quantitative coalgebras

- Generalizations of deterministic automata
- Quantitative coalgebras: set of states S and $t : S \rightarrow GS$

$$G ::= Id \mid B \mid G \times G \mid G + G \mid G^A \mid \mathbb{M}^G$$

\mathbb{M} is a monoid. $\mathcal{P} = 2^{Id}$ and $\mathcal{D}_\omega = \mathbb{R}^{Id}$

Examples

- | | |
|----------------------------------|------------------------|
| • $G = 2 \times Id^A$ | Deterministic automata |
| • $G = (B \times Id)^A$ | Mealy machines |
| • $G = (\mathcal{P}Id)^A$ | LTS |
| • $G = \mathcal{PD}_\omega(S)^A$ | Simple Segala systems |
| • ... | |

In this talk...

- ... we present a **systematic** way to derive from the functor **G**: languages of (generalized) regular expressions and
- ... sound and complete axiomatizations thereof for **quantitative systems**;
- ... we show the correspondence between language and systems (generalizing **Kleene's theorem**);
- ... we apply the framework to several types of probabilistic automata **recovering old results and deriving new ones**.

G-expressions

$$E \quad ::= \quad \emptyset \mid \epsilon \mid E \cdot E \mid E + E \mid E^*$$


$$E_G \quad ::= \quad ?$$

G-expressions

$$E \quad ::= \quad \emptyset \mid \epsilon \mid E \cdot E \mid E + E \mid E^*$$

$$E_G \quad ::= \quad ?$$

How do we define E_G ?



G-expressions

$$\begin{array}{lcl} \text{Exp} \ni \varepsilon & ::= & \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \\ & & \mid b \quad B \\ & & \mid l\langle \varepsilon \rangle \mid r\langle \varepsilon \rangle \quad G_1 \times G_2 \\ & & \mid l[\varepsilon] \mid r[\varepsilon] \quad G_1 + G_2 \\ & & \mid a(\varepsilon) \quad G^A \\ & & \mid m \cdot \varepsilon \quad M^G \end{array}$$

G-expressions

$$\begin{array}{lcl} \text{Exp} \ni \varepsilon & ::= & \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \\ & & \mid b \quad B \\ & & \mid l\langle \varepsilon \rangle \mid r\langle \varepsilon \rangle \quad G_1 \times G_2 \\ & & \mid l[\varepsilon] \mid r[\varepsilon] \quad G_1 + G_2 \\ & & \mid a(\varepsilon) \quad G^A \\ & & \mid m \cdot \varepsilon \quad M^G \end{array}$$

G-expressions

$$\begin{array}{lcl} \text{Exp} \ni \varepsilon & ::= & \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \\ & & \mid b \quad B \\ & & \mid l\langle \varepsilon \rangle \mid r\langle \varepsilon \rangle \quad G_1 \times G_2 \\ & & \mid l[\varepsilon] \mid r[\varepsilon] \quad G_1 + G_2 \\ & & \mid a(\varepsilon) \quad G^A \\ & & \mid m \cdot \varepsilon \quad M^G \end{array}$$

G-expressions

$$\begin{array}{lcl} \text{Exp} \ni \varepsilon & ::= & \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \\ & & \mid b \quad B \\ & & \mid l\langle \varepsilon \rangle \mid r\langle \varepsilon \rangle \quad G_1 \times G_2 \\ & & \mid l[\varepsilon] \mid r[\varepsilon] \quad G_1 + G_2 \\ & & \mid a(\varepsilon) \quad G^A \\ & & \mid m \cdot \varepsilon \quad M^G \end{array}$$

G-expressions

$$\begin{array}{lcl} \text{Exp} \ni \varepsilon & ::= & \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \\ & & \mid b \quad B \\ & & \mid l\langle \varepsilon \rangle \mid r\langle \varepsilon \rangle \quad G_1 \times G_2 \\ & & \mid l[\varepsilon] \mid r[\varepsilon] \quad G_1 + G_2 \\ & & \mid a(\varepsilon) \quad G^A \\ & & \mid m \cdot \varepsilon \quad M^G \end{array}$$

G-expressions

$$\begin{array}{lcl} \text{Exp} \ni \varepsilon & ::= & \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \\ & & \mid b \quad B \\ & & \mid l\langle \varepsilon \rangle \mid r\langle \varepsilon \rangle \quad G_1 \times G_2 \\ & & \mid l[\varepsilon] \mid r[\varepsilon] \quad G_1 + G_2 \\ & & \mid a(\varepsilon) \quad G^A \\ & & \mid m \cdot \varepsilon \quad M^G \end{array}$$

Examples

LTS expressions – $G = (\mathcal{P}Id)^A = (2^{Id})^A$

$$\varepsilon ::= \emptyset \mid \underbrace{\varepsilon \oplus \varepsilon \mid \mu X. \varepsilon}_G \mid$$

Examples

LTS expressions – $G = (\mathcal{P}Id)^A = (2^{Id})^A$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{a(\quad)}_{_A}$$

Examples

LTS expressions – $G = (\mathcal{P}Id)^A = (2^{Id})^A$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{a(\underbrace{1 \cdot \varepsilon}_{2^{Id}})}_{\substack{2^{Id} \\ _A}} \mid a(\underbrace{0 \cdot \varepsilon}_{2^{Id}})$$

Examples

LTS expressions – $G = (\mathcal{P}Id)^A = (2^{Id})^A$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{a(1 \cdot \varepsilon)}_{\substack{2^{Id} \\ _A}} \mid a(\underbrace{0 \cdot \varepsilon}_{2^{Id}})$$

$$\varepsilon ::= \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \mid a. \varepsilon$$

Examples

LTS expressions – $G = (\mathcal{P}Id)^A = (2^{Id})^A$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{a(\underbrace{1 \cdot \varepsilon}_{2^{Id}})}_{\substack{2^{Id} \\ _A}} \mid a(\underbrace{0 \cdot \varepsilon}_{2^{Id}})$$

$$\varepsilon ::= \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \mid a. \varepsilon$$

Markov Chain expressions – $G = \mathcal{D}_\omega(Id) = \mathbb{R}^{Id}$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid$$

Examples

LTS expressions – $G = (\mathcal{P}Id)^A = (2^{Id})^A$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{a(\underbrace{1 \cdot \varepsilon}_{2^{Id}})}_{\substack{2^{Id} \\ _A}} \mid a(\underbrace{0 \cdot \varepsilon}_{2^{Id}})$$

$$\varepsilon ::= \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \mid a. \varepsilon$$

Markov Chain expressions – $G = \mathcal{D}_\omega(Id) = \mathbb{R}^{Id}$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{p \cdot \varepsilon}_{\mathbb{R}^{Id}}$$

Examples

LTS expressions – $G = (\mathcal{P}Id)^A = (2^{Id})^A$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{a(\underbrace{1 \cdot \varepsilon}_{2^{Id}})}_{\substack{2^{Id} \\ _A}} \mid a(\underbrace{0 \cdot \varepsilon}_{2^{Id}})$$

$$\varepsilon ::= \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \mid a. \varepsilon$$

Markov Chain expressions – $G = \mathcal{D}_w(Id) = \mathbb{R}^{Id} \quad \sum_i p_i = 1$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{p \cdot \varepsilon}_{\mathbb{R}^{Id}}$$

Examples

LTS expressions – $G = (\mathcal{P}Id)^A = (2^{Id})^A$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{a(\underbrace{1 \cdot \varepsilon}_{2^{Id}})}_{\neg A} \mid a(\underbrace{0 \cdot \varepsilon}_{2^{Id}})$$

$$\varepsilon ::= \emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \gamma \mid a. \varepsilon$$

Markov Chain expressions – $G = \mathcal{D}_w(Id) = \mathbb{R}^{Id} \quad \sum_i p_i = 1$

$$\varepsilon ::= \underbrace{\emptyset \mid \varepsilon \oplus \varepsilon \mid \mu x. \varepsilon}_G \mid \underbrace{p \cdot \varepsilon}_{\mathbb{R}^{Id}}$$

$$\varepsilon ::= \mu x. \varepsilon \mid \bigoplus_{i \in 1 \dots n} p_i \cdot \varepsilon \quad \text{for } p_i \in (0, 1] \text{ such that } \sum_{i \in 1 \dots n} p_i = 1$$

Kleene's Theorem

The goal is:

G – expressions **correspond to** Finite G – coalgebras and vice-versa.
What does it mean **correspond**?

Final coalgebras exist for quantitative coalgebras.

Kleene's Theorem

The goal is:

G – expressions **correspond to** Finite G – coalgebras and vice-versa.

What does it mean **correspond**?

Final coalgebras exist for quantitative coalgebras.

$$\begin{array}{ccc} S & \xrightarrow{h} & \Omega_G \xleftarrow{[\![\cdot]\!]} Exp_G \\ \alpha \downarrow & & \downarrow \omega_G \\ GS & \xrightarrow{Gh} & G\Omega_G \end{array}$$

Kleene's Theorem

The goal is:

G – expressions **correspond to** Finite G – coalgebras and vice-versa.
What does it mean **correspond**?

Final coalgebras exist for quantitative coalgebras.

$$\begin{array}{ccccc}
 S & \xrightarrow{\quad h \quad} & \Omega_G & \xleftarrow{\quad \llbracket \cdot \rrbracket \quad} & Exp_G \\
 \alpha \downarrow & & \downarrow \omega_G & & \downarrow \lambda_G \\
 GS & \xrightarrow{\quad Gh \quad} & G\Omega_G & \xleftarrow{\quad G\llbracket \cdot \rrbracket \quad} & GExp_G
 \end{array}$$

Kleene's Theorem

The goal is:

G – expressions **correspond to** Finite G – coalgebras and vice-versa.
What does it mean **correspond**?

Final coalgebras exist for quantitative coalgebras.

$$\begin{array}{ccccc}
 S & \xrightarrow{h} & \Omega_G & \xleftarrow{[\![\cdot]\!]} & Exp_G \\
 \alpha \downarrow & & \downarrow \omega_G & & \downarrow \lambda_G \\
 GS & \xrightarrow{Gh} & G\Omega_G & \xleftarrow{G[\![\cdot]\!] } & GExp_G
 \end{array}$$

correspond \equiv mapped to the same element of the final coalgebra
 \equiv **bisimilar**

A generalized Kleene Theorem

Theorem

- 1 *Let (S, g) be a G -coalgebra. If S is finite then there exists for any $s \in S$ a G -expression ε_s such that $\varepsilon_s \sim s$.*
- 2 *For all G -expressions ε , there exists a finite G -coalgebra (S, g) such that $\exists_{s \in S} s \sim \varepsilon$.*

The proof provides algorithms to construct an expression from a system and vice-versa.

A generalized Kleene Theorem

Theorem

- 1 *Let (S, g) be a G -coalgebra. If S is finite then there exists for any $s \in S$ a G -expression ε_s such that $\varepsilon_s \sim s$.*
- 2 *For all G -expressions ε , there exists a finite G -coalgebra (S, g) such that $\exists_{s \in S} s \sim \varepsilon$.*

The proof provides algorithms to construct an expression from a system and vice-versa.

Axiomatization

$$\left. \begin{array}{lcl} \varepsilon_1 \oplus \varepsilon_2 & \equiv & \varepsilon_2 \oplus \varepsilon_1 \\ \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3) & \equiv & (\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \\ \varepsilon_1 \oplus \varepsilon_1 & \equiv & \varepsilon_1, \text{ } \textcolor{red}{G \text{ polynomial}} \\ \varepsilon \oplus \emptyset & \equiv & \varepsilon \end{array} \right\} \textcolor{blue}{G}$$

$$\left. \begin{array}{lcl} \mu X. \gamma & \equiv & \gamma[\mu X. \gamma / X] \\ \gamma[\varepsilon / X] \equiv \varepsilon & \Rightarrow & \mu X. \gamma \equiv \varepsilon \end{array} \right\} \textcolor{blue}{FP}$$

$$\left. \begin{array}{lcl} \emptyset & \equiv & 0 \\ m_1 \cdot \varepsilon \oplus m_2 \cdot \varepsilon & \equiv & (m_1 + m_2) \cdot \varepsilon \end{array} \right\} \textcolor{blue}{M^G}$$

$$\left. \begin{array}{lcl} \emptyset & \equiv & \perp_B \\ b_1 \oplus b_2 & \equiv & b_1 \vee b_2 \end{array} \right\} \textcolor{blue}{B}$$

$$\left. \begin{array}{lcl} I(\emptyset) & \equiv & \emptyset \\ I(\varepsilon_1) \oplus I(\varepsilon_2) & \equiv & I(\varepsilon_1 \oplus \varepsilon_2) \\ r(\emptyset) & \equiv & \emptyset \\ r(\varepsilon_1) \oplus r(\varepsilon_2) & \equiv & r(\varepsilon_1 \oplus \varepsilon_2) \end{array} \right\} \textcolor{blue}{G_1 \times G_2}$$

Sound and complete w.r.t \sim

Similar for $G_1 + G_2$ and G^A

Axiomatization

$$\left. \begin{array}{ll} \varepsilon_1 \oplus \varepsilon_2 & \equiv \varepsilon_2 \oplus \varepsilon_1 \\ \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3) & \equiv (\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \\ \varepsilon_1 \oplus \varepsilon_1 & \equiv \varepsilon_1, \text{ } \textcolor{red}{G} \text{ polynomial} \\ \varepsilon \oplus \emptyset & \equiv \varepsilon \end{array} \right\} \textcolor{blue}{G}$$

$$\left. \begin{array}{ll} \mu x. \gamma & \equiv \gamma[\mu x. \gamma / x] \\ \gamma[\varepsilon / x] \equiv \varepsilon & \Rightarrow \mu x. \gamma \equiv \varepsilon \end{array} \right\} \textcolor{blue}{FP}$$

$$\left. \begin{array}{ll} \emptyset & \equiv 0 \\ m_1 \cdot \varepsilon \oplus m_2 \cdot \varepsilon & \equiv (m_1 + m_2) \cdot \varepsilon \end{array} \right\} \textcolor{blue}{M}^G$$

Sound and complete w.r.t \sim

$$\left. \begin{array}{ll} \emptyset & \equiv \perp_B \\ b_1 \oplus b_2 & \equiv b_1 \vee b_2 \end{array} \right\} \textcolor{blue}{B}$$

$$\left. \begin{array}{ll} l(\emptyset) & \equiv \emptyset \\ l(\varepsilon_1) \oplus l(\varepsilon_2) & \equiv l(\varepsilon_1 \oplus \varepsilon_2) \\ r(\emptyset) & \equiv \emptyset \\ r(\varepsilon_1) \oplus r(\varepsilon_2) & \equiv r(\varepsilon_1 \oplus \varepsilon_2) \end{array} \right\} G_1 \times G_2$$

Similar for $G_1 + G_2$ and G^A

Axiomatization

$$\left. \begin{array}{ll} \varepsilon_1 \oplus \varepsilon_2 & \equiv \varepsilon_2 \oplus \varepsilon_1 \\ \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3) & \equiv (\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \\ \varepsilon_1 \oplus \varepsilon_1 & \equiv \varepsilon_1, \text{ } \textcolor{red}{G} \text{ polynomial} \\ \varepsilon \oplus \emptyset & \equiv \varepsilon \end{array} \right\} \textcolor{blue}{G}$$

$$\left. \begin{array}{ll} \mu X. \gamma & \equiv \gamma[\mu X. \gamma / X] \\ \gamma[\varepsilon / X] \equiv \varepsilon & \Rightarrow \mu X. \gamma \equiv \varepsilon \end{array} \right\} \textcolor{blue}{FP}$$

$$\left. \begin{array}{ll} \emptyset & \equiv 0 \\ m_1 \cdot \varepsilon \oplus m_2 \cdot \varepsilon & \equiv (m_1 + m_2) \cdot \varepsilon \end{array} \right\} \textcolor{blue}{M}^G$$

Sound and complete w.r.t \sim

$$\left. \begin{array}{ll} \emptyset & \equiv \perp_B \\ b_1 \oplus b_2 & \equiv b_1 \vee b_2 \end{array} \right\} \textcolor{blue}{B}$$

$$\left. \begin{array}{ll} I(\emptyset) & \equiv \emptyset \\ I(\varepsilon_1) \oplus I(\varepsilon_2) & \equiv I(\varepsilon_1 \oplus \varepsilon_2) \\ r(\emptyset) & \equiv \emptyset \\ r(\varepsilon_1) \oplus r(\varepsilon_2) & \equiv r(\varepsilon_1 \oplus \varepsilon_2) \end{array} \right\} G_1 \times G_2$$

Similar for $G_1 + G_2$ and G^A

Axiomatization

$$\left. \begin{array}{ll} \varepsilon_1 \oplus \varepsilon_2 & \equiv \varepsilon_2 \oplus \varepsilon_1 \\ \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3) & \equiv (\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \\ \varepsilon_1 \oplus \varepsilon_1 & \equiv \varepsilon_1, \text{ } \textcolor{red}{G} \text{ polynomial} \\ \varepsilon \oplus \emptyset & \equiv \varepsilon \end{array} \right\} \textcolor{blue}{G}$$

$$\left. \begin{array}{ll} \mu X. \gamma & \equiv \gamma[\mu X. \gamma / X] \\ \gamma[\varepsilon / X] \equiv \varepsilon & \Rightarrow \mu X. \gamma \equiv \varepsilon \end{array} \right\} \textcolor{blue}{FP}$$

$$\left. \begin{array}{ll} \emptyset & \equiv 0 \\ m_1 \cdot \varepsilon \oplus m_2 \cdot \varepsilon & \equiv (m_1 + m_2) \cdot \varepsilon \end{array} \right\} \textcolor{blue}{M}^G$$

$$\left. \begin{array}{ll} \emptyset & \equiv \perp_B \\ b_1 \oplus b_2 & \equiv b_1 \vee b_2 \end{array} \right\} \textcolor{blue}{B}$$

$$\left. \begin{array}{ll} I(\emptyset) & \equiv \emptyset \\ I(\varepsilon_1) \oplus I(\varepsilon_2) & \equiv I(\varepsilon_1 \oplus \varepsilon_2) \\ r(\emptyset) & \equiv \emptyset \\ r(\varepsilon_1) \oplus r(\varepsilon_2) & \equiv r(\varepsilon_1 \oplus \varepsilon_2) \end{array} \right\} G_1 \times G_2$$

Sound and complete w.r.t \sim

Similar for $G_1 + G_2$ and G^A

Axiomatization

$$\left. \begin{array}{ll} \varepsilon_1 \oplus \varepsilon_2 & \equiv \varepsilon_2 \oplus \varepsilon_1 \\ \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3) & \equiv (\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \\ \varepsilon_1 \oplus \varepsilon_1 & \equiv \varepsilon_1, \text{ } \textcolor{red}{G} \text{ polynomial} \\ \varepsilon \oplus \emptyset & \equiv \varepsilon \end{array} \right\} \textcolor{blue}{G}$$

$$\left. \begin{array}{ll} \mu X. \gamma & \equiv \gamma[\mu X. \gamma / X] \\ \gamma[\varepsilon / X] \equiv \varepsilon & \Rightarrow \mu X. \gamma \equiv \varepsilon \end{array} \right\} \textcolor{blue}{FP}$$

$$\left. \begin{array}{ll} \emptyset & \equiv 0 \\ m_1 \cdot \varepsilon \oplus m_2 \cdot \varepsilon & \equiv (m_1 + m_2) \cdot \varepsilon \end{array} \right\} \textcolor{blue}{M}^G$$

$$\left. \begin{array}{ll} \emptyset & \equiv \perp_B \\ b_1 \oplus b_2 & \equiv b_1 \vee b_2 \end{array} \right\} \textcolor{blue}{B}$$

$$\left. \begin{array}{ll} l(\emptyset) & \equiv \emptyset \\ l(\varepsilon_1) \oplus l(\varepsilon_2) & \equiv l(\varepsilon_1 \oplus \varepsilon_2) \\ r(\emptyset) & \equiv \emptyset \\ r(\varepsilon_1) \oplus r(\varepsilon_2) & \equiv r(\varepsilon_1 \oplus \varepsilon_2) \end{array} \right\} G_1 \times G_2$$

Sound and complete w.r.t \sim

Similar for $G_1 + G_2$ and G^A

Axiomatization

$$\left. \begin{array}{ll} \varepsilon_1 \oplus \varepsilon_2 & \equiv \varepsilon_2 \oplus \varepsilon_1 \\ \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3) & \equiv (\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \\ \varepsilon_1 \oplus \varepsilon_1 & \equiv \varepsilon_1, \text{ } \textcolor{red}{G \text{ polynomial}} \\ \varepsilon \oplus \emptyset & \equiv \varepsilon \end{array} \right\} \textcolor{blue}{G}$$

$$\left. \begin{array}{ll} \mu X. \gamma & \equiv \gamma[\mu X. \gamma / X] \\ \gamma[\varepsilon / X] \equiv \varepsilon & \Rightarrow \mu X. \gamma \equiv \varepsilon \end{array} \right\} \textcolor{blue}{FP}$$

$$\left. \begin{array}{ll} \emptyset & \equiv 0 \\ m_1 \cdot \varepsilon \oplus m_2 \cdot \varepsilon & \equiv (m_1 + m_2) \cdot \varepsilon \end{array} \right\} \textcolor{blue}{M^G}$$

Sound and complete w.r.t \sim

$$\left. \begin{array}{ll} \emptyset & \equiv \perp_B \\ b_1 \oplus b_2 & \equiv b_1 \vee b_2 \end{array} \right\} \textcolor{blue}{B}$$

$$\left. \begin{array}{ll} l(\emptyset) & \equiv \emptyset \\ l(\varepsilon_1) \oplus l(\varepsilon_2) & \equiv l(\varepsilon_1 \oplus \varepsilon_2) \\ r(\emptyset) & \equiv \emptyset \\ r(\varepsilon_1) \oplus r(\varepsilon_2) & \equiv r(\varepsilon_1 \oplus \varepsilon_2) \end{array} \right\} G_1 \times G_2$$

Similar for $G_1 + G_2$ and G^A

Axiomatization

$$\left. \begin{array}{ll} \varepsilon_1 \oplus \varepsilon_2 & \equiv \varepsilon_2 \oplus \varepsilon_1 \\ \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3) & \equiv (\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \\ \varepsilon_1 \oplus \varepsilon_1 & \equiv \varepsilon_1, \text{ } G \text{ polynomial} \\ \varepsilon \oplus \emptyset & \equiv \varepsilon \end{array} \right\} G$$

$$\left. \begin{array}{ll} \mu X. \gamma & \equiv \gamma[\mu X. \gamma / X] \\ \gamma[\varepsilon / X] \equiv \varepsilon & \Rightarrow \mu X. \gamma \equiv \varepsilon \end{array} \right\} FP$$

$$\left. \begin{array}{ll} \emptyset & \equiv 0 \\ m_1 \cdot \varepsilon \oplus m_2 \cdot \varepsilon & \equiv (m_1 + m_2) \cdot \varepsilon \end{array} \right\} M^G$$

$$\left. \begin{array}{ll} \emptyset & \equiv \perp_B \\ b_1 \oplus b_2 & \equiv b_1 \vee b_2 \end{array} \right\} B$$

$$\left. \begin{array}{ll} l(\emptyset) & \equiv \emptyset \\ l(\varepsilon_1) \oplus l(\varepsilon_2) & \equiv l(\varepsilon_1 \oplus \varepsilon_2) \\ r(\emptyset) & \equiv \emptyset \\ r(\varepsilon_1) \oplus r(\varepsilon_2) & \equiv r(\varepsilon_1 \oplus \varepsilon_2) \end{array} \right\} G_1 \times G_2$$

Sound and complete w.r.t \sim

Similar for $G_1 + G_2$ and G^A

Results I : Segala systems – $\mathcal{P}(D_\omega(Id))^A$

$$\varepsilon:: = \emptyset \mid \varepsilon \boxplus \varepsilon \mid \mu X. \varepsilon \mid X \mid a(\{\varepsilon'\})$$

$$\varepsilon':: = \bigoplus_{i \in 1 \dots n} p_i \cdot \varepsilon_i$$

where $a \in A$, $p_i \in (0, 1]$ and $\sum_{i \in 1 \dots n} p_i = 1$

$$(\varepsilon_1 \boxplus \varepsilon_2) \boxplus \varepsilon_3 \equiv \varepsilon_1 \boxplus (\varepsilon_2 \boxplus \varepsilon_3)$$

$$\varepsilon_1 \boxplus \varepsilon_2 \equiv \varepsilon_2 \boxplus \varepsilon_1$$

$$\varepsilon \boxplus \emptyset \equiv \varepsilon$$

$$\varepsilon \boxplus \varepsilon \equiv \varepsilon$$

$$(\varepsilon'_1 \oplus \varepsilon'_2) \oplus \varepsilon'_3 \equiv \varepsilon'_1 \oplus (\varepsilon'_2 \oplus \varepsilon'_3)$$

$$\varepsilon'_1 \oplus \varepsilon'_2 \equiv \varepsilon'_2 \oplus \varepsilon'_1$$

$$(p_1 \cdot \varepsilon) \oplus (p_2 \cdot \varepsilon) \equiv (p_1 + p_2) \cdot \varepsilon$$

$$\varepsilon[\mu X. \varepsilon / X] \equiv \mu X. \varepsilon$$

$$\gamma[\varepsilon / X] \equiv \varepsilon \Rightarrow \mu X. \gamma \equiv \varepsilon$$

Same syntax and axioms as in [Deng and Palamidessi'05]

Results I : Segala systems – $\mathcal{P}(D_\omega(Id))^A$

$$\varepsilon:: = \emptyset \mid \varepsilon \boxplus \varepsilon \mid \mu x. \varepsilon \mid x \mid a(\{\varepsilon'\})$$

$$\varepsilon':: = \bigoplus_{i \in 1 \dots n} p_i \cdot \varepsilon_i$$

where $a \in A$, $p_i \in (0, 1]$ and $\sum_{i \in 1 \dots n} p_i = 1$

$$(\varepsilon_1 \boxplus \varepsilon_2) \boxplus \varepsilon_3 \equiv \varepsilon_1 \boxplus (\varepsilon_2 \boxplus \varepsilon_3)$$

$$\varepsilon_1 \boxplus \varepsilon_2 \equiv \varepsilon_2 \boxplus \varepsilon_1$$

$$\varepsilon \boxplus \emptyset \equiv \varepsilon$$

$$\varepsilon \boxplus \varepsilon \equiv \varepsilon$$

$$(\varepsilon'_1 \oplus \varepsilon'_2) \oplus \varepsilon'_3 \equiv \varepsilon'_1 \oplus (\varepsilon'_2 \oplus \varepsilon'_3)$$

$$\varepsilon'_1 \oplus \varepsilon'_2 \equiv \varepsilon'_2 \oplus \varepsilon'_1$$


$$(p_1 \cdot \varepsilon) \oplus (p_2 \cdot \varepsilon) \equiv (p_1 + p_2) \cdot \varepsilon$$

$$\varepsilon[\mu x. \varepsilon / x] \equiv \mu x. \varepsilon$$

$$\gamma[\varepsilon / x] \equiv \varepsilon \Rightarrow \mu x. \gamma \equiv \varepsilon$$

Same syntax and axioms as in [Deng and Palamidessi'05]

Results I : Segala systems – $\mathcal{P}(D_\omega(Id))^A$



\Uparrow
Kleene's Theorem

$$\Downarrow$$

$$a(\{1/2 \cdot \emptyset \oplus 1/2 \cdot \emptyset\}) \boxplus a(\{1/3 \cdot \emptyset \oplus 2/3 \cdot \emptyset\}) \boxplus b(\{1 \cdot \emptyset\})$$

Results I : Segala systems – $\mathcal{P}(D_\omega(Id))^A$



↑↑
Kleene's Theorem

$$\begin{aligned} & \Downarrow \\ & a(\{1/2 \cdot \emptyset \oplus 1/2 \cdot \emptyset\}) \boxplus a(\{1/3 \cdot \emptyset \oplus 2/3 \cdot \emptyset\}) \boxplus b(\{1 \cdot \emptyset\}) \\ \equiv & a(\{1 \cdot \emptyset\}) \boxplus a(\{1 \cdot \emptyset\}) \boxplus b(\{1 \cdot \emptyset\}) \end{aligned}$$

Results I : Segala systems – $\mathcal{P}(D_\omega(Id))^A$



↑↑
Kleene's Theorem

$$\begin{aligned} & \Downarrow \\ & a(\{1/2 \cdot \emptyset \oplus 1/2 \cdot \emptyset\}) \boxplus a(\{1/3 \cdot \emptyset \oplus 2/3 \cdot \emptyset\}) \boxplus b(\{1 \cdot \emptyset\}) \\ \equiv & a(\{1 \cdot \emptyset\}) \boxplus a(\{1 \cdot \emptyset\}) \boxplus b(\{1 \cdot \emptyset\}) \\ \equiv & a(\{1 \cdot \emptyset\}) \boxplus b(\{1 \cdot \emptyset\}) \end{aligned}$$

Results II : Stratified systems – $D_\omega(Id) + (B \times Id) + 1$

$$\varepsilon ::= \mu x. \varepsilon \mid x \mid \langle b, \varepsilon \rangle \mid \bigoplus_{i \in 1 \dots n} p_i \cdot \varepsilon_i \mid \downarrow$$

where $b \in B$, $p_i \in (0, 1]$ and $\sum_{i \in 1 \dots n} p_i = 1$

$$(\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \equiv \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3)$$

$$\varepsilon_1 \oplus \varepsilon_2 \equiv \varepsilon_2 \oplus \varepsilon_1$$

$$(p_1 \cdot \varepsilon) \oplus (p_2 \cdot \varepsilon) \equiv (p_1 + p_2) \cdot \varepsilon$$

$$\varepsilon[\mu x. \varepsilon / x] \equiv \mu x. \varepsilon$$

$$\gamma[\varepsilon / x] \equiv \varepsilon \Rightarrow \mu x. \gamma \equiv \varepsilon$$

Same syntax as in [van Glabbeek, Smolka and Steffen'95] and new axiomatization (inexistent).

Results II : Stratified systems – $D_\omega(Id) + (B \times Id) + 1$

$$\varepsilon ::= \mu x. \varepsilon \mid x \mid \langle b, \varepsilon \rangle \mid \bigoplus_{i \in 1 \dots n} p_i \cdot \varepsilon_i \mid \downarrow$$

where $b \in B$, $p_i \in (0, 1]$ and $\sum_{i \in 1 \dots n} p_i = 1$

$$(\varepsilon_1 \oplus \varepsilon_2) \oplus \varepsilon_3 \equiv \varepsilon_1 \oplus (\varepsilon_2 \oplus \varepsilon_3)$$

$$\varepsilon_1 \oplus \varepsilon_2 \equiv \varepsilon_2 \oplus \varepsilon_1$$

$$(p_1 \cdot \varepsilon) \oplus (p_2 \cdot \varepsilon) \equiv (p_1 + p_2) \cdot \varepsilon$$

$$\varepsilon[\mu x. \varepsilon / x] \equiv \mu x. \varepsilon$$

$$\gamma[\varepsilon / x] \equiv \varepsilon \Rightarrow \mu x. \gamma \equiv \varepsilon$$

Same syntax as in [van Glabbeek, Smolka and Steffen'95] and new axiomatization (inexistent).

Results III : Pnueli-Zuck systems – $\mathcal{PD}_\omega \mathcal{P}(Id)^A$

$$\varepsilon:: = \emptyset \mid \varepsilon \boxplus \varepsilon \mid \mu x. \varepsilon \mid x \mid \{\varepsilon'\}$$

$$\varepsilon':: = \bigoplus_{i \in 1 \dots n} p_i \cdot \varepsilon''_i$$

$$\varepsilon'':: = \emptyset \mid \varepsilon'' \boxplus \varepsilon'' \mid a(\{\varepsilon\})$$

where $a \in A$, $p_i \in (0, 1]$ and $\sum_{i \in 1 \dots n} p_i = 1$

$$(\varepsilon_1 \boxplus \varepsilon_2) \boxplus \varepsilon_3 \equiv \varepsilon_1 \boxplus (\varepsilon_2 \boxplus \varepsilon_3)$$

$$\varepsilon_1 \boxplus \varepsilon_2 \equiv \varepsilon_2 \boxplus \varepsilon_1$$

$$\varepsilon \boxplus \emptyset \equiv \varepsilon$$

$$\varepsilon \boxplus \varepsilon \equiv \varepsilon$$

$$(\varepsilon'_1 \oplus \varepsilon'_2) \oplus \varepsilon'_3 \equiv \varepsilon'_1 \oplus (\varepsilon'_2 \oplus \varepsilon'_3) \quad \varepsilon'_1 \oplus \varepsilon'_2 \equiv \varepsilon'_2 \oplus \varepsilon'_1$$

$$(p_1 \cdot \varepsilon'') \oplus (p_2 \cdot \varepsilon'') \equiv (p_1 + p_2) \cdot \varepsilon''$$

$$\varepsilon[\mu x. \varepsilon / x] \equiv \mu x. \varepsilon$$

$$\gamma[\varepsilon / x] \equiv \varepsilon \Rightarrow \mu x. \gamma \equiv \varepsilon$$

New syntax and axiomatization.

Results III : Pnueli-Zuck systems – $\mathcal{PD}_\omega \mathcal{P}(Id)^A$

$$\varepsilon:: = \emptyset \mid \varepsilon \boxplus \varepsilon \mid \mu x. \varepsilon \mid x \mid \{\varepsilon'\}$$

$$\varepsilon':: = \bigoplus_{i \in 1 \dots n} p_i \cdot \varepsilon''_i$$

$$\varepsilon'':: = \emptyset \mid \varepsilon'' \boxplus \varepsilon'' \mid a(\{\varepsilon\})$$

where $a \in A$, $p_i \in (0, 1]$ and $\sum_{i \in 1 \dots n} p_i = 1$

$$(\varepsilon_1 \boxplus \varepsilon_2) \boxplus \varepsilon_3 \equiv \varepsilon_1 \boxplus (\varepsilon_2 \boxplus \varepsilon_3)$$

$$\varepsilon_1 \boxplus \varepsilon_2 \equiv \varepsilon_2 \boxplus \varepsilon_1$$

$$\varepsilon \boxplus \emptyset \equiv \varepsilon$$

$$\varepsilon \boxplus \varepsilon \equiv \varepsilon$$

$$(\varepsilon'_1 \oplus \varepsilon'_2) \oplus \varepsilon'_3 \equiv \varepsilon'_1 \oplus (\varepsilon'_2 \oplus \varepsilon'_3) \quad \varepsilon'_1 \oplus \varepsilon'_2 \equiv \varepsilon'_2 \oplus \varepsilon'_1$$

$$(p_1 \cdot \varepsilon'') \oplus (p_2 \cdot \varepsilon'') \equiv (p_1 + p_2) \cdot \varepsilon''$$

$$\varepsilon[\mu x. \varepsilon / x] \equiv \mu x. \varepsilon$$

$$\gamma[\varepsilon / x] \equiv \varepsilon \Rightarrow \mu x. \gamma \equiv \varepsilon$$

New syntax and axiomatization.

Conclusions and future work

Conclusions

- Framework to **uniformly** derive language and axioms for quantitative coalgebras (weighted automata, probabilistic automata, etc)
- Examples show the effectiveness of the framework: known syntaxes recovered, new ones derived.

Future work

- Apply the framework to other systems, *e.g.* alternating systems.
- Automation: `Circ` — Coinductive prover