

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

BY

XXXXXXX

17/56EB0XX

A PROJECT SUBMITTED TO THE DEPARTMENT OF
MATHEMATICS, FACULTY OF PHYSICAL SCIENCES, UNIVERSITY
OF ILORIN, ILORIN, KWARA STATE, NIGERIA.

IN PARTIAL FULFILMENT OF REQUIREMENTS FOR THE AWARD
OF BACHELOR OF SCIENCE (B. Sc.) DEGREE IN MATHEMATICS.

February, 2022

CERTIFICATION

This is to certify that this project was carried out by **XXXXXXXXXXXXXXXXXXXXXXX**
with Matriculation Number 17/56EB0XX in the Department of Mathemat-
ics, Faculty of Physical Sciences, University of Ilorin, Ilorin, Nigeria, for the
award of Bachelor of Science (B.Sc.) degree in Mathematics.

.....

Dr. E.O. Titiloye
Supervisor

.....

Date

.....

Prof. K. Rauf
Head of Department

.....

Date

.....

Prof.o
External Examiner

.....

Date

ACKNOWLEDGMENTS

All praises

i

DEDICATION

I would like to dedicate the project to God

ABSTRACT

In this project

Table of Contents

TITLE PAGE	1
CERTIFICATION	i
ACKNOWLEDGMENTS	ii
DEDICATION	iii
ABSTRACT	iv
TABLE OF CONTENTS	v
1	1
2	2
3	3
4 WORKED EXAMPLES	4
4.1 Non-linear Volterra Integral Equation using the Adomian De- composition Method	4
4.2 Non-linear Fredholm Integral Equation using the Adomian Decomposition Method	8

5	SUMMARY AND CONCLUSION	11
5.1	Summary	11
5.2	Conclusion	11
	REFERENCES	12

Chapter 1

Chapter 2

Chapter 3

Chapter 4

WORKED EXAMPLES

4.1 Non-linear Volterra Integral Equation using the Adomian Decomposition Method

Example 1

$$y(x) = x + \int_0^x y^4(t)dt$$

Solution

$$\sum_{n=0}^{\infty} y_n(x) = x + \int_0^x \sum_{n=0}^{\infty} A_n(t)dt$$

The Adomian polynomial for y^4

$$A_0 = y_0^4, \quad A_1 = 4y_1y_0^3, \quad A_2 = 4y_2y_0^3 + 6y_1^2y_0^2, \quad A_3 = 4y_3y_0^3 + 12y_2y_1y_0^2 + 4y_1^3y_0$$

where

$$y_0(x) = x, \quad y_{k+1}(x) = \int_0^x A_k(t)dt, k \geq 0$$

$$\begin{aligned}
y_1(x) &= \int_0^x y_0^4(t) dt = \int_0^x t^4 dt = \frac{t^5}{5} \Big|_0^x = \frac{1}{5} x^5 \\
y_2(x) &= \int_0^x (4y_1 y_0^3) dt = \int_0^x \left(\frac{4}{5} t^5 \cdot t^3 \right) dt \\
&= \frac{4}{5} \int_0^x t^8 dt = \frac{4}{5} \frac{t^9}{9} \Big|_0^x = \frac{4}{45} x^9 \\
y_3(x) &= \int_0^x (4y_2 y_0^3 + 6y_1^2 y_0^2) dt \\
&= \int_0^x 4 \left(\frac{4}{45} t^9 \right) (t^3) + 6 \left(\frac{1}{5} t^5 \right)^2 (t^2) dt \\
&= \int_0^x \left(\frac{16}{45} t^{12} + \frac{6}{5} t^{12} \right) dt \\
&= \frac{14}{9} \int_0^x t^{12} dt = \frac{14}{9} \frac{t^{13}}{13} \Big|_0^x = \frac{14}{117} x^{13} \\
\therefore y(x) &= x + \frac{1}{5} x^5 + \frac{4}{45} x^9 + \frac{14}{117} x^{13} + \dots
\end{aligned}$$

Example 2

$$y(x) = x + \int_0^x (t-x) y^2(t) dt$$

Solution

$$\sum_{n=0}^{\infty} y_n(x) = x + \int_0^x (t-x) \sum_{n=0}^{\infty} A_n(t) dt$$

the Adomian polynomial of y^2

$$A_0 = y_0^2, \quad A_1 = 2y_0 y_1, \quad A_2 = 2y_2 y_0 + y_1^2, \quad A_3 = 2y_0 y_3 + 2y_1 y_2,$$

$$y_0(x) = x, \quad y_{k+1}(x) = \int_0^x (t-x) A_k(t) dt$$

$$\begin{aligned}
y_1(x) &= \int_0^x (t-x)t^2 dt = \int_0^x (t^3 - xt^2) dt = \left. \frac{t^4}{4} - \frac{xt^3}{3} \right|_0^x \\
&= \frac{x^4}{4} - \frac{x^4}{3} = -\frac{1}{12}x^4 \\
y_2(x) &= \int_0^\infty (t-x)(2t) \left(-\frac{1}{12}t^4 \right) dt = \int_0^x (t-x) \left(-\frac{1}{6}t^5 \right) dt \\
&= -\frac{1}{6} \int_0^x (t^6 - xt^5) dt = -\frac{1}{6} \left[\frac{t^7}{7} - \frac{xt^6}{6} \right]_0^x \\
&= -\frac{1}{6} \left[\frac{x^7}{7} - \frac{x^7}{6} \right] = -\frac{1}{6} \left[-\frac{1}{42}x^7 \right] = \frac{1}{180}x^7 \\
y_3(x) &= \int_0^x (t-x) \left[\left(\frac{2}{180}t^7 \right) (t) + \left(-\frac{1}{12}t^4 \right)^2 \right] dt \\
&= \int_0^x (t-x) \left(\frac{1}{90}t^8 + \frac{1}{12}t^8 \right) dt = \int_0^x (t-x) \left(\frac{17}{180}t^8 \right) dt \\
&= \frac{17}{180} \int_0^x (t^9 - xt^8) dt = \frac{17}{180} \left[\frac{t^{10}}{10} - \frac{xt^9}{9} \right]_0^x \\
&= \frac{17}{180} \left[\frac{x^{10}}{10} - \frac{x^{10}}{9} \right] = \frac{17}{180} \left[-\frac{1}{90}x^{10} \right] = -\frac{17}{16200}x^{10} \\
\therefore y(x) &= x - \frac{1}{12}x^4 + \frac{1}{180}x^7 - \frac{17}{16200}x^{10}
\end{aligned}$$

Example 3

$$y(x) = x + \int_0^x (x^2t - xt^2)y^2(t)dt$$

Solution

$$\sum_{n=0}^{\infty} y_n(x) = x + \int_0^x (x^2t - xt^2) \sum_{n=0}^{\infty} A_n(t) dt$$

$$y_0(x) = x, \quad y_{k+1}(x) = \int_0^\infty (x^2t - xt^2)A_k(t)dt$$

$$\begin{aligned} y_1(x) &= \int_0^x (x^2t - xt^2)(t^2)dt = \int_0^x (x^2t^3 - xt^4)dt \\ &= \left. \frac{x^2t^4}{4} - \frac{xt^5}{5} \right|_0^x = \frac{x^6}{4} - \frac{x^6}{5} = \frac{1}{20}x^6 \end{aligned}$$

$$\begin{aligned} y_2(x) &= \int_0^x (x^2t - xt^2)(2y_0y_1)dt \\ &= \int_0^x (x^2t - xt^2) \left(\frac{2}{10}t^7 \right) dt = \frac{1}{10} \int_0^x (x^2t^8 - xt^9)dt \\ &= \frac{1}{10} \left[\frac{x^2t^9}{9} - \frac{xt^{10}}{10} \right]_0^x = \frac{1}{10} \left[\frac{x^{11}}{9} - \frac{x^{11}}{10} \right] = \frac{1}{900}x^{11} \end{aligned}$$

$$\begin{aligned} y_3(x) &= \int_0^x (x^2t - xt^2)(2y_2y_0 + y_1^2)dt \\ &= \int_0^x (x^2t - xt^2) \left[\left(\frac{2}{900}t^{11} \right) (t) + \left(\frac{1}{20}t^6 \right)^2 \right] dt \\ &= \int_0^x (x^2t - xt^2) \left(\frac{1}{450}t^{12} + \frac{1}{400}t^{12} \right) dt \\ &= \int_0^x (x^2t - xt^2) \left(\frac{17}{3600}t^{12} \right) dt \\ &= \frac{17}{3600} \int_0^x (x^2t^{13} - xt^{14})dt \\ &= \frac{17}{3600} \left[\frac{x^2t^{14}}{14} - \frac{xt^{15}}{15} \right]_0^x = \frac{17}{3600} \left[\frac{x^{16}}{14} - \frac{x^{16}}{15} \right] \\ &= \frac{17}{3600} \left[\frac{1}{210}x^{16} \right] = \frac{17}{756000}x^{16} \\ y(x) &= x + \frac{1}{20}x^6 + \frac{1}{900}x^{11} + \frac{17}{756000}x^{16} + \dots \end{aligned}$$

4.2 Non-linear Fredholm Integral Equation using the Adomian Decomposition Method

Example 1

$$y(x) = 3 + \lambda \int_0^1 ty^2(t)dt$$

Solution

$$y_0(x) = 3, \quad y_{k+1}(x) = \lambda \int_0^1 y^2(t)dt$$

$$y_1(x) = \lambda \int_0^1 ty_0^2(t)dt$$

$$= \lambda \int_0^1 9t dt = \frac{9\lambda t^2}{2} \Big|_0^1 = \frac{9\lambda}{2}$$

$$y_2(x) = \lambda \int_0^1 (2y_0y_1)t dt$$

$$= \lambda \int_0^1 (6) \left(\frac{9\lambda}{2} \right) t dt = \frac{27\lambda^2 t^2}{2} \Big|_0^1 = \frac{27\lambda^2}{2}$$

$$y_3(x) = \lambda \int_0^1 \left(81\lambda^2 + \frac{81\lambda^2}{2} \right) t dt = \lambda \int_0^1 \left(\frac{243\lambda^2}{2} \right) t dt$$

$$= \frac{243\lambda^3 t^2}{4} \Big|_0^1 = \frac{243\lambda^3}{4}$$

$$y_4(x) = \lambda \int_0^1 (2y_0y_3 + 2y_1y_2)t dt$$

$$= \lambda \int_0^1 \left(\frac{729}{2}\lambda^3 + \frac{243}{2}\lambda^3 \right) t dt$$

$$\begin{aligned}
&= \lambda \int_0^1 (486\lambda^3)t dt = \frac{486\lambda^4}{2} \Big|_0^1 \\
&= 243\lambda^4
\end{aligned}$$

$$y(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + y_4(x) + \dots$$

$$y(x) = 3 + \frac{9\lambda}{2} + \frac{27\lambda^2}{2} + \frac{243\lambda^3}{4} + 243\lambda^4 + \dots$$

Example 2

$$y(x) = 4 + \lambda \int_0^1 t^2 y^2(t) dt$$

Solution

The Adomian polynomial for $y^2(x)$

$$A_0(x) = y_0^2, \quad A_1(x) = 2y_0y_1,$$

$$A_2(x) = 2y_0y_2 + y_1^2, \quad A_3 = 2y_0y_3 + 2y_1y_2$$

$$y_0(x) = 4$$

$$y_1(x) = \lambda \int_0^1 y^2(t)t^2 dt = \lambda \int_0^1 16t^2 dt = \frac{16}{3}t^3 \lambda \Big|_0^1 = \frac{16}{3}\lambda$$

$$y_2(x) = \lambda \int_0^1 (2y_0y_1)t^2 dt = \lambda \int_0^1 \frac{128}{3}\lambda t^2 dt = \frac{\lambda^2 128}{9}t^3 \Big|_0^1 = \frac{128}{9}\lambda^2$$

$$\begin{aligned}
y_3(x) &= \lambda \int_0^1 (2y_0y_2 + y_1^2)t^2 dt = \lambda \int_0^1 \left(\frac{1024\lambda^2}{9} + \frac{256\lambda^2}{9} \right) t^2 dt \\
&= \lambda \int_0^1 \frac{1280\lambda^2}{9} t^2 dt = \frac{1280}{27}\lambda^3 t^3 \Big|_0^1 = \frac{1280}{27}\lambda^3
\end{aligned}$$

$$\begin{aligned}
y_4(x) &= \lambda \int_0^1 (2y_0y_3 + 2y_1y_2)t^2 dt = \lambda \int_0^1 \left(\frac{5120}{27}\lambda^3 + \frac{4096}{27}\lambda^3 \right) t^2 dt \\
&= \lambda \int_0^1 \left(\frac{9216}{27}\lambda^3 \right) t^3 dt = \frac{9216}{81}\lambda^4 t^4 \Big|_0^1 = \frac{9216}{81}\lambda^4 \\
y(x) &= 4 + \frac{16}{3}\lambda + \frac{128}{9}\lambda^2 + \frac{1280}{27}\lambda^3 + \frac{9216}{81}\lambda^4 + \dots
\end{aligned}$$

Example 3

$$y(x) = 1 + \lambda \int_0^1 y^4(t) dt$$

Solution

The Adomian polynomials of $y^4(t)$ are

$$A_0(x) = y_0^4, \quad A_1(x) = 4y_1y_0^3, \quad A_2(x) = 4y_2y_0^3 + 6y_1^2y_0^2$$

$$A_3(x) = 4y_3y_0^3 + 12y_2y_1y_0^2 + 4y_1^3y_0$$

$$y_0(x) = 1$$

$$y_1(x) = \lambda \int_0^1 y_0^4(t) dt = \lambda \int_0^1 (1^4) dt = \lambda t \Big|_0^1 = \lambda$$

$$y_2(x) = \lambda \int_0^1 4y_1y_0^3(t) dt = \lambda \int_0^1 4\lambda dt = 4\lambda^2 t \Big|_0^1 = 4\lambda^2$$

$$\begin{aligned}
y_3(x) &= \lambda \int_0^1 4y_2y_0^3 + 6y_1^2y_0^2(t) dt = \lambda \int_0^1 (16\lambda^2 + 6\lambda^2) dt \\
&= \lambda \int_0^1 22\lambda^2 dt = 22\lambda^3 t \Big|_0^1 = 22\lambda^3
\end{aligned}$$

$$y_4(x) = \lambda \int_0^1 (4y_3y_0^3 + 12y_2y_1y_0^2 + 4y_1^3y_0) dt$$

$$\begin{aligned}
&= \lambda \int_0^1 (88\lambda^3 + 48\lambda^3 + 4\lambda^3) dt = \lambda \int_0^1 140\lambda^3 dt \\
&= 140\lambda^4 t \Big|_0^1 = 140\lambda^4 \\
y(x) &= 1 + \lambda + 4\lambda^2 + 22\lambda^3 + 140\lambda^4 + \cdots
\end{aligned}$$

Chapter 5

SUMMARY AND CONCLUSION

5.1 Summary

5.2 Conclusion

REFERENCES