


journal homepage: www.elsevier.com/locate/csbj

Mini review

Rhizosphere microbes enhance plant salt tolerance: Toward crop production in saline soil

Yunpeng Liu^a, Weibing Xun^b, Lin Chen^c, Zhihui Xu^b, Nan Zhang^b, Haichao Feng^b, Qiang Zhang^d, Ruifu Zhang^{a,b,*}^a Key Laboratory of Microbial Resources Collection and Preservation, Ministry of Agriculture and Rural Affairs, Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing 100081, PR China^b Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, National Engineering Research Center for Organic-based Fertilizers, Jiangsu Collaborative Innovation Center for Solid Organic Waste Resource Utilization, Nanjing Agricultural University, Nanjing 210095, PR China^c Experimental Center of Forestry in North China, Chinese Academy of Forestry, Beijing 102300, PR China^d Heze Kingenta Ecological Engineering Co., Ltd, Heze, Shandong 274000, PR China

ARTICLE INFO

Article history:

Received 31 July 2022

Received in revised form 22 November 2022

Accepted 22 November 2022

Available online 25 November 2022

Keywords:

Plant salt tolerance

Rhizosphere microbe

Saline soil

Microbe-enhanced salt tolerance mechanisms

ABSTRACT

The world's population continues to increase and thus requires more food production to take place in nonarable land, such as saline soil; therefore, it is urgent to find solutions to enhance the salinity tolerance of crops. As the second genome of plants, the rhizosphere microbiome plays critical roles in plant fitness under stress conditions. Many beneficial microbes that help plants cope with salinity stress have been identified, highlighting their roles in mitigating salt stress-induced negative effects on plants. However, a comprehensive review of the microbial species that are able to confer plant salt tolerance and the underlying mechanisms is still lacking. In this review, we compared the representative fungal and bacterial taxa that demonstrate the ability to enhance plant growth in saline soil. We also reviewed the mechanisms by which rhizosphere microbes enhance plant salt stress tolerance, i.e., by re-establishing ion and osmotic homeostasis, preventing damage to plant cells, and resuming plant growth under salt stress. Finally, future research efforts to explore the rhizosphere microbiome for agricultural sustainability are proposed.

© 2022 The Author(s). Published by Elsevier B.V. on behalf of Research Network of Computational and Structural Biotechnology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction	6544
2. Microbial taxa enhancing plant tolerance to salt stress	6544
2.1. Arbuscular mycorrhizal fungi (AMF)	6544
2.2. <i>Trichoderma</i> spp.	6544
2.3. <i>Serendipita indica</i> (formerly <i>Piriformospora indica</i>)	6544
2.4. <i>Enterobacter</i> spp.	6545
2.5. <i>Bacillus</i> spp.	6545
2.6. <i>Pseudomonas</i> spp.	6545
3. Mechanisms by which microbes enhance plant salt tolerance	6545
3.1. Re-establishing ion and osmotic homeostasis	6545
3.1.1. Ion balance	6545
3.1.2. Osmotic balance	6546
3.2. Preventing damage to plant cells	6547
3.3. Resuming plant growth under salt stress	6547

* Corresponding author at: Key Laboratory of Microbial Resources Collection and Preservation, Ministry of Agriculture and Rural Affairs, Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing 100081, PR China.

E-mail address: rfzhang@njau.edu.cn (R. Zhang).

4. Summary and perspective	6548
CRedit authorship contribution statement	6549
Declaration of Competing Interest	6549
Appendix A. Supplementary data	6549
References	6549

1. Introduction

Agricultural production worldwide is facing the challenge of increasing population size. Food production is increasingly restricted by limited cultivated land due to soil degradation issues, such as salinization. Saline soil accounts for 6 % of the world's total land area and is a severe hindrance to agricultural production [1]. Most domesticated crops are salt sensitive and cannot be cultivated in saline soil [2]. Currently, the increasing food demand of global populations makes finding a solution to enhance crop tolerance to salinity more urgent.

As the functions of rhizosphere microbes in plant fitness have been recognized [3], many beneficial microbes that help plants cope with salinity stress have been isolated and reapplied to the soil to enhance crop production in saline soil. These microbes can be used in different ways; the common method is to make microbial fertilizers or microbial agents [4,5], which can be applied to the soil directly or combined with chemical or organic fertilizers. In addition, some strains can be coated on crop seeds before sowing [6]. Their efficiency in increasing plant growth or promoting seed germination under saline conditions has been shown, indicating a role in the mitigation of salt stress-induced negative effects on plants. The plant salt tolerance induced by some microbial groups, such as bacterial endophytes [7], the halophyte microbiome [8], microbial 1-aminocyclopropane-1-carboxylic acid (ACC) deaminase [9] or microbial coinoculation [10], has been reviewed, as well as the microbial metabolites responsible for enhancing plant salt tolerance [11]. However, a comprehensive comparison of the microbial species that are able to confer plant salt tolerance and the underlying mechanisms is still lacking.

Here, we reviewed the fungal and bacterial isolates with clear evidence of promoting plant growth or seed germination under saline stress. Specifically, we discussed the challenges for the agricultural application of each kind of microbial inoculant. We also summarized the mechanisms by which microbes enhance plant tolerance to salt stress. The salt stress signaling and tolerance mechanism of plants have been well reviewed [2,12–18] and will not be included in this review.

2. Microbial taxa enhancing plant tolerance to salt stress

2.1. Arbuscular mycorrhizal fungi (AMF)

AMF are well-known beneficial fungi belonging to the phylum Glomeromycota that have close symbiotic interactions with their host plants. The main function of AM symbiosis seems to be assisting plants with nutrient uptake by extending the active absorption area of roots [10,19]. Several studies have also demonstrated the effect of AMF on plant productivity under salt stress. AMF, including the genera *Glomus*, *Funneliformis*, *Septoglomus*, *Gigaspora*, *Acaulospora*, *Claroideoglomus*, and *Rhizophagus* (*Funneliformis*, *Claroideoglomus* and *Rhizophagus* used to belong to *Glomus*), have shown the capability to enhance plant salt tolerance (Table S1). The plants that showed enhanced production under saline stress upon AMF inoculation include lettuce (up to 80 mM NaCl) [20], maize (up to 100 mM NaCl) [21], wheat (up to 200 mM NaCl) [22], rice (up to 120 mM NaCl) [23] and some leguminous (up to 200 mM NaCl) [24] and woody plants (Table S1). The characteris-

tics and advances of AMF in mitigating plant salt stress include integrity with significantly enhanced plant nutrient uptake, such as nitrogen (N) and potassium (K), especially phosphorus (P) [25]. AMF-induced plant salt tolerance is always correlated with phosphorus, which plays an important role in photosynthesis to supply energy for the stress response [26]. However, the symbiosis of AMF with plants is negatively affected by salt stress because AMF are not salt tolerant, which has been confirmed by several studies [25,27–30]. Pandey et al. [31] found that native AMF isolated from saline soil are more effective in mitigating the negative effect of salt stress on *Cajanus cajan*, suggesting that the salt tolerance of AMF themselves is also a limiting factor for performance under salt stress.

2.2. *Trichoderma* spp.

Trichoderma spp. are free-living soil filamentous fungi that are beneficial to plants [32]. *Trichoderma* strains are nonsymbiotic, while some strains can infect root outer cells [32]. *Trichoderma* has diverse beneficial functions for plants, such as enhancing plant stress resistance and promoting plant growth. *T. harzianum*, *T. virens*, *T. atroviride*, *T. asperelloides*, *T. longibrachiatum*, *T. yunnanense*, and *T. afroharzianum* have shown the ability to promote seed germination and plant growth of *Arabidopsis* (up to 100 mM NaCl) [33], cucumber (up to 200 mM NaCl) [34], wheat (up to 200 mM NaCl) [6] and barley (up to 200 mM NaCl) [35] under salt stress (Table S1). These strains promoted root development and affected root system architecture. In addition, most *Trichoderma* strains isolated from soil are efficient decomposers of dead plant residues that drive soil carbon cycling and plant growth [32]. *Trichoderma* strains are not an outstanding microbial group regarding salt tolerance when compared with some bacterial strains, such as *Bacillus* spp. and *Enterobacter* spp. Contreras-Cornejo et al. [36] demonstrated that 200 mM NaCl significantly inhibited the hyphal growth of *T. virens* and *T. atroviride*. Zhang et al. [37] found similar results that 30 mg/mL NaCl inhibited the growth of *Trichoderma longibrachiatum* T6. Therefore, both salt tolerance for itself and plant salt tolerance enhancement are necessary for the successful cultivation of a good functional *Trichoderma* isolate in saline soil.

2.3. *Serendipita indica* (formerly *Piriformospora indica*)

Piriformospora indica is a natural soil-inhabiting filamentous fungus that promotes plant growth upon symbiosis. *P. indica* is an endophyte for a broad range of host plants (Franken 2012). It was first isolated from an AMF spore by Verma et al. in 1998. However, unlike AMF, *P. indica* can be cultivated on defined media. Several reviews have concluded the beneficial effects of *P. indica* on plants [10,38–40]. In recent years, *P. indica* has shown impressive potential in mitigating salt stress in *Arabidopsis* (up to 200 mM NaCl) [41], rice (up to 300 mM NaCl) [42], barley (up to 300 mM NaCl) [43] and tomato (up to 150 mM NaCl) [44] (Table S1). Lanza et al. [45] demonstrated that free-living *P. indica* is sensitive to salt; it can only tolerate up to 100 mM NaCl, and hyphal growth was significantly inhibited due to the toxic effect of Na⁺ rather than osmotic stress. Interestingly, Vahabi et al. [41] demonstrated that the growth promotion of *Arabidopsis* seedlings under salt stress is

strongly dependent on physical contact between the two symbionts.

2.4. *Enterobacter* spp.

The specific role of *Enterobacter* spp. as beneficial endophytes in enhancing plant salt tolerance has been demonstrated in their interactions with *Arabidopsis* (up to 200 mM NaCl) [46], *rice* (up to 150 mM NaCl) [47], *wheat* (up to 200 mM NaCl) [48], *maize* (electrical conductivity (EC) up to 13.6 dS/m) [49], *tomato* (up to 200 mM NaCl) [46], *mung bean* (1 % NaCl) [50], *okra* (75 mM NaCl) [51] and *quinoa* (400 mM NaCl) [52] (Table S1). *Enterobacter* spp. were first recognized as beneficial plant bacteria due to ACC deaminase activity [53]. *Enterobacter* sp. SA187 is a well-studied isolate that promotes plant growth under salt stress. It seems that the modulation of ethylene signaling in plants plays a key role in SA187-induced salt tolerance [54]. Ali et al. [55] demonstrated that *E. cloacae* PM2 can survive under salinity stress of up to 3 M NaCl in Luria-Bertani (LB) medium. However, the careful selection of plant-safe strains is necessary given that members of the *Enterobacter* genus are generally known as opportunistic pathogens of plants and humans [56].

2.5. *Bacillus* spp.

Bacillus spp. are gram-positive spore-forming bacteria from the phylum *Firmicutes* and represent the most widely used and well-studied bacteria in agricultural production. Most of the *Bacillus* strains that are beneficial to plants belong to the *subtilis* subgroup. These bacteria are generally root surface colonizers or root endophytes that exert diverse plant beneficial functions. *Bacillus* spp. have been reported to induce salt tolerance in *Arabidopsis* (up to 250 mM NaCl) [57,58], *maize* (up to 100 mM NaCl) [59], *barley* (up to 200 mM NaCl) [60], *rice* (up to 200 mM NaCl) [61], *wheat* (up to 400 mM NaCl) [62], *tomato* (up to 300 mM NaCl) [63], *chick-pea* (2 % NaCl) [64], *cotton* (up to 200 mM NaCl) [65], *soybean* (up to 240 mM NaCl) [66] and some halophytes (Table S1). *Bacillus* can tolerate high salt concentrations and form stress-tolerant spores when exposed to adverse environments [55]. The feature of spore formation greatly increases bacterial survival in soil.

2.6. *Pseudomonas* spp.

Pseudomonas spp. are gram-negative bacteria from the phylum *Proteobacteria* and represent one of the most abundant rhizobacterial groups. The species enhancing plant salt tolerance include *P. putida*, *P. simiae*, *P. syringae*, *P. fluorescens*, *P. migulae*, *P. thivervalensis*, *P. geniculata*, *P. corrugate*, *P. stutzeri*, *P. chlororaphis*, *P. extremorientalis*, *P. extremorientalis*, *P. aurantiaca* and *P. chlororaphis*. *Arabidopsis* (up to 200 mM NaCl) [67], *maize* (up to 150 mM NaCl) [68], *wheat* (up to 200 mM NaCl) [69], *cotton* (0.35 % NaCl) [70], *cucumber* (100 mM NaCl) [71], *peanut* (100 mM NaCl) [72], *soybean* (100 mM NaCl) [73,74], *tomato* (up to 185 mM NaCl) [75], *sunflower* (EC up to 15.9 dS/m) [76], *oats* and *barley* (EC up to 9.4 dS/m) [77] have shown elevated growth under salt stress upon inoculation with *Pseudomonas* spp. (Table S1). *Pseudomonas* is a naturally highly abundant bacterial group in the rhizosphere, suggesting that *Pseudomonas* is adapted to the rhizosphere and may perform well in colonizing roots and thus enhancing plant salt tolerance. However, as nonspore-forming bacteria, the relatively low survival rate of microbial fertilizers or microbial agents during the process of production and storage may be disadvantageous for their application in agriculture.

3. Mechanisms by which microbes enhance plant salt tolerance

High salinity is a complex type of stress for plants. On the one hand, high amounts of Na^+ and Cl^- in the cytoplasm (mainly Na^+) are directly toxic to cells, as they interfere with many protein functions, thus influencing cell physiology [2,16]. On the other hand, high salt concentrations in soil lead to hyperosmotic stress in plant cells and subsequent difficulty in acquiring water and soil nutrients [2,16]. Both direct ion stress and osmotic stress can be sensed immediately after plants are exposed to high-salinity conditions. After perceiving the primary signal, secondary signals, such as Ca^{2+} , ROS and ABA, are generated to induce stress-responsive genes [18]. However, if ROS accumulate, the cells are damaged, and programmed cell death is initiated. ABA accumulation induces stomatal closure to reduce transpiration, which also reduces water use efficiency (WUE) and CO_2 use efficiency (CUE). To counter the negative effects of Na^+ and ROS causing cell damage and reducing WUE and CUE, plants must produce more ROS-scavenging enzymes, transport Na^+ into vacuoles, and produce osmolytes, which are costly with regard to energy, thereby leading to growth inhibition [1,13,17,18].

The NaCl tolerance of microbes far exceeds the salt tolerance of plants. Microbes can concentrate compatible solutes to protect against the osmotic imbalance between the cytoplasm and the environment and express diverse Na^+ transporters to reduce cytoplasmic Na^+ concentrations. In addition to tolerating high salinity, many fungal and bacterial strains can enhance plant tolerance to salinity and mitigate the salt stress-induced inhibition of plant growth. Resuming growth under stress conditions is risky because growth inhibition by stress physiology is actually an energy trade-off with tolerance [16]. Nevertheless, microbes have multiple effects on plant ion and osmotic balance, ROS regulation and detoxification, and photosynthesis. These processes are interrelated and ultimately promote plant growth under salt stress via three mechanisms: re-establishing ion and osmotic homeostasis, preventing damage to plant cells, and resuming plant growth under salt stress (Fig. 1).

3.1. Re-establishing ion and osmotic homeostasis

3.1.1. Ion balance

The accumulation of Na^+ in the plant cytoplasm is toxic to plant cells. Na^+ competes with other ions to bind proteins, which leads to the inactivation of enzymes. Na^+ also inhibits the uptake of K^+ and other ions due to the impact of ion balance between the two sides of the cytoplasmic membrane. Therefore, the accumulation of Na^+ in the cytoplasm must be reduced, and the ion balance must be re-established for roots to acquire other ions.

To counter the high Na^+ concentration in the environment, plants need to reduce Na^+ influx through the plasma membrane, which is mediated by the high-affinity K^+ channel (HKT) [78,78,79]. Many microbial strains have been shown to increase plant shoot and/or root K^+/Na^+ ratios by increasing K^+ and/or reducing Na^+ in the cytoplasm, but the underlying mechanisms vary. HKT1, which plays critical roles in both Na^+ influx across the plasma membrane and long-distance Na^+ transport, is considered a key determinant of plant salt tolerance [79]. Expression of HKT1 in shoots promotes plant salt tolerance by reducing Na^+ accumulation in shoots [80]. The specific role of HKT1 in the decrease in microbe-mediated plant Na^+ under salt stress has been recognized. For example, Zhang et al. [81] demonstrated that *B. subtilis* GB03 stimulated HKT1 expression in shoots while reducing it in roots under 100 mM NaCl by producing VOCs. *Athkt1;1*, but not *sos3*, mutant plants failed to show GB03-induced plant growth and a decrease in Na^+ under salt stress [81]. Niu et al. [82] demon-




Fig. 1. Mechanisms by which microbes enhance plant salt tolerance. 1. To re-establish the ion balance of plants under salt stress, microbes reduce the Na^+ content in the cytoplasm by decreasing Na^+ uptake, increasing Na^+ efflux and compartmentalizing Na^+ into vacuoles. To regulate the osmotic balance of plants under salt stress, microbes produce osmolytes and induce plants to produce osmolytes, and they also reduce ABA to increase water uptake. 2. To prevent ROS-induced damage to plants under salt stress, microbes can reduce plant ROS generation and stimulate plants to scavenge ROS. 3. To resume plant growth under salt stress, microbes promote photosynthesis and produce ACC deaminase or other compounds to regulate plant ethylene signaling.

strated that enhanced *Puccinellia tenuiflora* salt tolerance is dependent on modulating the expression of *HKT1*. In contrast, *Pseudomonas simiae* AU decreased the expression of *HKT1* in soybean shoots under salt stress [73]. *P. indica* colonized *Arabidopsis* roots under salt stress and showed enhanced expression of *HKT1* [83]. Inoculation with AMF significantly induced *HKT1*;1 and *HKT1*;6 in both the roots and shoots of lettuce under 60 mM NaCl, but in the same study, inoculation with another AMF strain, *Claroideoglomus lamellosum*, did not significantly change the expression profiles for *HKT1*;1 and *HKT1*;6 in lettuce [84].

The excess Na^+ in the cytoplasm under salt stress can be either exported out of the protoplast by the plasma membrane Na^+/H^+ antiporter [17] or compartmentalized into the vacuole by the vacuolar Na^+/H^+ exchanger (NHX) [85]. Increasing the Na^+ efflux by the SOS pathway and vacuolar NHX-driven vacuolar Na^+ compartmentalization is another strategy for reducing cytoplasmic Na^+ accumulation [86]. SOS1, a plasma membrane Na^+/H^+ antiporter controlled by the regulation cascade of SOS2 and SOS3, is believed to be a key determinant of Na^+ exclusion in roots [17]. Chen et al.

[87,88] showed that *B. amyloliquefaciens* SQR9 strongly stimulated the expression of *NHX1*, *NHX2* and *NHX3* in maize under salt stress by producing spermidine. However, either mutation of *NHX1* or blocking the SOS pathway, but not mutation with *HKT1*, leads to deficiency in SQR9- and spermidine-induced salt tolerance, indicating the determinant role of *NHX1* and *SOS1* in SQR9-induced salt tolerance [88]. Baek et al. [89] demonstrated a similar conclusion that the SOS pathway is necessary for *B. orydicola* YC7007-induced *Arabidopsis* tolerance to salt stress. There are also some cases in which microbial inoculation increases K^+ uptake but has no effect on Na^+ accumulation [54].

3.1.2. Osmotic balance

Soil salinity is always accompanied by hyperosmotic stress in plants. A high salt concentration is hyperosmotic to plant cells and will affect root water absorption or even water loss. This situation also limits the uptake of nutrients by roots and results in ABA accumulation. ABA signaling will lead to stomatal closure, which restricts transpiration and thus makes it harder to absorb water

and reduce photosynthesis. Accumulating soluble osmoprotectants in the cytoplasm is a general strategy to counter hyperosmotic stress. These osmolytes, including charged metabolites, polyols, soluble sugars and complex sugars, vary based on plant species [2].

Most of the microbes that show mitigation effects on plant salt stress also increase the WUE and stimulate the production of different kinds of osmolytes by plants. The production of osmolytes under saline conditions can be controlled by microbes. For example, *Bacillus subtilis* GB03 produced VOCs that induced glycine betaine production by *Arabidopsis* under saline conditions in an ABA-dependent manner [81]. The mutation of trehalose synthesis in *Pseudomonas putida* UW4 completely blocked the promotion of tomato root development and growth under saline conditions, suggesting a critical role of microbe-produced osmolytes in plant salt tolerance [90,91]. Salt stress-induced ABA accumulation is believed to be a signature of the stress response and contributes to stomatal closure [92]. Although ABA and proline accumulation in salt-stressed plants generally occurs, inoculation with microbes has inconsistent effects on ABA and proline contents under salt stress. While most studies propose that the observed ABA reduction in plants upon microbial inoculation is responsible for restoration of the osmotic balance under salt stress, *Bacillus licheniformis* SA03 [93], *Paenibacillus yonginensis* DCY84T [94] and *Bacillus mesonae* H20-5 [95] increased the ABA contents in *Chrysanthemum*, *Panax ginseng* and *tomato* under saline conditions, respectively. By inhibiting ABA synthesis using fluridone, Zhou et al. [93] demonstrated the key role of SA03-induced ABA in enhancing plant salt tolerance. The effects of microbial inoculations on plant proline accumulation upon saline stress also vary, and the physiological significance of proline accumulation in salt-stressed plants is controversial (Table S1). Proline protects plants from saline stress by increasing the osmosis of the cytoplasm and is induced upon salt stress and strengthened by microbial inoculation. However, a decrease in proline in inoculated plants compared with that of noninoculated plants under saline conditions has also been widely reported [53,96–98]. Proline is an osmolyte and a signature of the stress response. Increased levels of proline upon microbial inoculation undoubtedly benefit salt tolerance, but the decreased levels of proline upon inoculation can also reflect a decrease in salt stress [99]. Therefore, the observed inconsistency in proline accumulation may also be due to the difference in the stress period. As plant endogenous osmolytes, proline and poly- γ -glutamic acid produced by microbes can also protect plants directly or act as a signal to induce plant salt tolerance [100,101].

3.2. Preventing damage to plant cells

Stress signaling following Ca^{2+} influx is always accompanied by ROS production [102]. ROS act as critical signals by activating various secondary messengers in the stress response, but they are also deleterious to plant cells by impairing DNA, proteins and lipids when they accumulate in plant cells [103]. The cell damage caused by ROS, including superoxide anion, hydrogen peroxide, hydroxyl radical, and singlet oxygen, is irreversible, especially for the elicitation of programmed cell death when hydroxyl radicals are produced [103]. Under normal conditions, ROS production and elimination are balanced; however, salt stress causes an imbalance due to the increase in ROS [102]. It is widely reported that plant electrolyte leakage and lipid peroxidation, as signatures of plasma membrane impairment by ROS [102], could be reduced by microbes.

3.2.1. Reducing ROS generation. Inoculation with microbes that reduce ROS generation may exert beneficial effects in preventing or arresting oxidative damage. ROS are produced from two sources: metabolic ROS are mainly produced in the chloroplast, mitochondria and peroxisome via different pathways, and signal-

ing ROS are produced in the apoplast by NADPH oxidases [104]. Under salt stress, there is a decrease in transpiration rates, which reduces CO_2 availability to the plants. Salt stress also exposes chloroplasts to high excitation energy, which causes ROS generation [105]. It is evident that microbial inoculation is capable of stimulating stomatal conductance and increasing the CO_2 availability of plants under saline conditions. In particular, increasing chlorophyll and carotenoid contents in salt-stressed plants is a common feature of microbe-primed plant salt tolerance (Table S1). Native AMF-inoculated maize showed higher efficiencies of photosystem II and stomatal conductance, which are believed to decrease photorespiration and ROS production [106,107]. Sulfur metabolism is connected to photosynthesis due to its role in forming iron-sulfur cluster proteins, which act as electron carriers in photosynthesis. Andres-Barrao et al. [108] found that *Enterobacter* sp. SA187 suppressed *Arabidopsis* salt stress-induced ROS accumulation in plastids by increasing root sulfur assimilation. Inoculation with SA187 also rescued the hypersensitivity of *Arabidopsis* LSU mutants, in which four genes encoding chloroplast-targeted proteins (LSU1–LSU4) that activate Fe-superoxide dismutase were silenced [108].

3.2.2. Scavenging excess ROS. In addition to reducing plant ROS generation, microbes can stimulate the ROS detoxification system in plants. Both the enzymatic antioxidant system, including superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), glutathione peroxidase (GPX), and peroxiredoxin (PRX), and the nonenzymatic system, including ascorbic acid, glutathione (GSH) and proline, are induced when plants are under salt stress; microbial inoculation can strengthen this induction process and stimulate plants to scavenge excess ROS. For example, Waller et al. [43] found that *P. indica*-infested roots under saline conditions increased the dehydroascorbate reductase activity and ascorbate content and decreased the level of dehydroascorbate. The critical role of stimulating plant ROS scavenging is confirmed by the fact that the salt tolerance induced by *B. velezensis* and the produced spermidine is dependent on plant GSH production [88]. It is evident that *Pseudomonas*-produced phenazine also contributes to catalase induction and thus ROS scavenging in plants under saline stress [69]. Moreover, some genes involved in Fe acquisition were activated in *Bacillus licheniformis* SA03-inoculated plants under salt stress [93]. Considering the Fenton reaction between iron and H_2O_2 that generates hydroxyl radicals [104], the modulation of iron homeostasis in plant cells by microbial inoculants may also contribute to the avoidance of hydroxyl radicals causing cell damage and subsequent cell death.

In contrast, several microbes with high antioxidant activity attenuated the salt stress-induced antioxidant system. For example, inoculation with *Brevibacterium linens* RS16 was found to reduce the salt stress-induced plant emissions of all major classes of volatile organic compounds (VOCs), which serve as antioxidants to tolerate stress [109]. An endophytic isolate of the fungus *Yarrowia lipolytica* selected on the basis of high antioxidant activity has reduced the peroxidase and catalase activity levels of maize in saline conditions compared to noninoculated maize [97]. These examples suggest that microbes can also enhance the salt tolerance of plants by directly scavenging ROS, which results in a reduction in stress-induced antioxidant production in plants.

3.3. Resuming plant growth under salt stress

Transporting ions, producing osmolytes and scavenging ROS, as discussed above, are all energy-consuming processes. For plants under salt-stressed conditions, resuming growth is a difficult choice for energy distribution and is precisely controlled by a very complex signaling transduction network.

3.3.1. Promotion of photosynthesis. An important determinant of microbe-induced salt tolerance is the improvement of photosynthesis to compensate for the energy cost of tolerance. It is generally recognized that salt stress greatly reduces the chlorophyll content, stomatal conductance and intercellular CO₂ concentration and strongly inhibits the net photosynthetic and transpiration rates [110]. As discussed above, almost all microbe-induced plant growth under salinity stress occurs in conjunction with the mitigation of stress-reduced photosynthesis. Increased biomass is always correlated with the net photosynthetic rate (Table S1). Increased transpiration by microbe inoculation will then lead to higher water absorption and nutrient uptake by roots, including those of phosphorus and sulfur, and thus promote photosynthesis. There is a possibility that the increased water absorption and nutrient uptake in stressed plants upon microbe inoculation is also a result of the microbial manipulation of the root system architecture, which occurs commonly in plant–microbe interactions [36]. The mechanism by which microbes initiate this positive feedback in plants under saline conditions is not clear. Since microbes can always directly or indirectly influence plant hormone signaling, which controls photosynthesis, it is necessary to understand the roles of phytohormones in microbe-induced plant growth under saline conditions.

3.3.2. Modulating IAA and ethylene signaling. IAA is important for plant growth and root development but is not a generally known central phytohormone in salt-stress responses in plants [111]. Salt stress-induced plant growth inhibition results from the suppression of IAA signaling, as shown by the decreased GUS density in the *DR5:uidA* line and the severely reduced fresh weight of auxin transport or signaling mutants, such as *arf7arf19* and *tir1afb2afb3*, under saline conditions [36]. Several studies have demonstrated the microbial functions in activating IAA signaling and increasing endogenous IAA content in salt-stressed plants [36,37,70,112]. IAA signaling has been proven to play a key role in *Trichoderma*- and *Bacillus*-modulated root systemic architecture (RSA) under nonstress conditions since mutations in genes involved in auxin transport or signaling reduced the growth-promoting and root developmental effects of inoculation [36,113,114]. It is likely that activating IAA signaling is required for microbe-induced plant growth and root development under saline stress, but definitive evidence is still lacking.

Ethylene is undoubtedly an important phytohormone involved in regulating plant growth under stress, but its role in plant salt tolerance is controversial [115]. Producing ACC deaminase, which is encoded by *acdS*, is an important characteristic of plant salt tolerance-inducing microbes [9]. ACC deaminase degrades ACC, the ethylene precursor, thereby reducing ethylene production and suppressing ethylene signaling, resulting in the activation of auxin signaling [9]. The significance of ACC deaminase-producing microbes in elevating salt tolerance has been recognized in *canola*, *wheat*, *tomato*, *barley* and *red pepper* [116–119]. This function is evident based on the deficiency in decreasing plant ethylene under saline stress of the bacterial *acdS* mutant and on the enhanced salt-stress tolerance and reduced ethylene content in the *acdS* transgenic line of *Camelina sativa* [120] and rice [121]. In another example of microbes influencing ethylene signaling, *Enterobacter* sp. SA187 produces 2-keto-4-methylthiobutyric acid (KMBA), which can be converted into ethylene in planta, as well as ACC. de Zélécourt et al. [54] demonstrated that KMBA is responsible for modulating the plant ethylene signaling pathway and the SA187-induced salt tolerance of *Arabidopsis*. They deployed the pEBF2::GUS reporter in *Arabidopsis* and found that SA187 inoculation activated the expression of EBF2 in root tips as well as ACC treatment. By using several *Arabidopsis* mutants, de Zélécourt et al. [54] pro-

posed that both SA187- and KMBA-induced salt tolerance is dependent on ethylene signaling rather than ethylene production or JA and ABA signaling. Conclusively, it seems that ACC deaminase and KMBA have somewhat similar effects on downstream ethylene signaling in different ways. Interestingly, de Zélécourt et al. [54] demonstrated that 100 nM ACC during salt stress could largely mimic the beneficial activity of SA187 on *Arabidopsis*, which should be opposite to the effect of ACC deaminase. Due to the controversial role of ethylene in plant abiotic stress tolerance, further investigation of the signaling perception in the ethylene signaling pathway upon ACC treatment, ACC deaminase and KMBA is needed to clarify the contradictory phenomenon.

Recently, two research groups simultaneously increased the interaction between microbe-induced plant salt tolerance and plant immunity. Loo et al. [122] proposed that microbe-associated molecular patterns (MAMPs) responsible for plant pattern-triggered immunity (PTI) could induce *Arabidopsis* salt tolerance when sensed by pattern-recognizing receptors (PRRs), a mechanism termed pattern-triggered salt tolerance (PTST). Rolli et al. [123] demonstrated that LYK4, which is responsible for the plant perception of chitin that triggers PTI, mediates *Enterobacter* sp.-triggered salt tolerance in *Arabidopsis thaliana*. These studies help to make sense of the crosstalk between microbe-induced biotic stress resistance and microbe-induced abiotic stress tolerance.

4. Summary and perspective

Due to the requirements of sustainable development and the multifunctional property of rhizosphere microbes, developing a microbial inoculant-based strategy is a promising direction for crop cultivation in saline soil. In recent years, scientists have drawn much attention to the isolation of strains and their effects on ion/osmotic balance, cell damage prevention and growth promotion of plants under saline stress. However, several issues remain that restrict the microbial strategy from taking further steps as a solution for enhancing crop tolerance in saline land.

- 1) The microbe-produced signaling molecules responsible for inducing plant salt tolerance are still lacking and remain to be identified. The identification of these molecules may support the development of chemical agents for promoting production in saline soil.
- 2) The detailed mechanisms by which microbes enhance plant salt tolerance, which is important for agricultural production, need to be deeply investigated. For example, many publications have demonstrated the regulation of plant NHXs or HKTs by microbes, but the underlying mechanisms are not clear. The mechanisms pertaining to the regulation of photosynthesis, antioxidants, osmolytes and growth by microbes are also not clear.
- 3) Some microbial species show efficiency in helping with plant tolerance to salt stress but are difficult to industrialize due to their relatively weak adaptability to either the production environment, storage environment, or saline soil environment. Therefore, matched technologies for microbial agent production and application need to be developed.
- 4) Enhancing crop production in saline soil is the final scope of research on microbe-induced plant salt tolerance, but there is still a lack of reports on the effect of microbial application on crop production in large-scale field experiments in saline soil, especially for some typical large saline soil land areas. The effects of microbes on crop production in different kinds of saline land may vary due to other factors, such as climate and soil type.

Funding

This work was funded by the National Key Research and Development Program (2021YFF1000404), the Central Public-interest Scientific Institution Basal Research Fund (No. Y2022QC15), the Agricultural Science and Technology Innovation Program (ASTIP No. CAAS-ZDRW202201) and the TaiShan Industrial Experts Program (2018TSCYCX-37).

CRediT authorship contribution statement

Yunpeng Liu: Funding acquisition, Writing – original draft. **Weibing Xun:** Writing– review and editing. **Lin Chen:** Visualization, Writing – original draft. **Haichao Feng:** Validation. **Zhihui Xu:** Writing– review& editing. **Nan Zhang:** Writing– review & editing. **Qiang Zhang:** Validation. **Ruifu Zhang:** Conceptualization.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.csbj.2022.11.046>.

References

- Munns R, Tester M. Mechanisms of salinity tolerance. *Annu Rev Plant Biol* 2008;59:651–81. <https://doi.org/10.1146/annurev.arplant.59.032607.092911>.
- Yang Y, Guo Y. Elucidating the molecular mechanisms mediating plant salt-stress responses. *New Phytol* 2018;217:523–39. <https://doi.org/10.1111/nph.14920>.
- Lugtenberg B, Kamilova F. Plant-growth-promoting rhizobacteria. *Annu Rev Microbiol* 2009;63:541–56. <https://doi.org/10.1146/annurev.micro.62.081307.162918>.
- Tchakounté GVT, Berger B, Patz S, Becker M, Fankem H, Taffouo VD, et al. Selected rhizosphere bacteria help tomato plants cope with combined phosphorus and salt stresses. *Microorganisms* 2020;8:1–16. <https://doi.org/10.3390/microorganisms8111844>.
- Adesemoye AO, Kloepper JW. Plant-microbes interactions in enhanced fertilizer-use efficiency. *Appl Microbiol Biotechnol* 2009;85:1–12. <https://doi.org/10.1007/s00253-009-2196-0>.
- Oljira AM, Hussain T, Waghmode TR, Zhao H, Sun H, Liu X, et al. *Trichoderma* enhances net photosynthesis, water use efficiency, and growth of wheat (*Triticum aestivum* L.) under salt stress. *Microorganisms* 2020;8:1–19. <https://doi.org/10.3390/microorganisms8101565>.
- Kushwaha P, Kashyap PL, Bhardwaj AK, Kuppusamy P, Srivastava AK, Tiwari RK. Bacterial endophyte mediated plant tolerance to salinity: growth responses and mechanisms of action. *World J Microbiol Biotechnol* 2020;36:1–16. <https://doi.org/10.1007/s11274-020-2804-9>.
- Etesami H, Beattie GA. Mining halophytes for plant growth-promoting halotolerant bacteria to enhance the salinity tolerance of non-halophytic crops. *Front Microbiol* 2018;9. <https://doi.org/10.3389/fmicb.2018.00148>.
- Orozco-Mosqueda M del C, Glick BR, Santoyo G. ACC deaminase in plant growth-promoting bacteria (PGPB): An efficient mechanism to counter salt stress in crops. *Microbiological Research* 2020;235:126439. <https://doi.org/10.1016/j.micres.2020.126439>.
- Khalvandi M, Amerian M, Pirdashti H, Keramati S. Does co-inoculation of mycorrhiza and *Piriformospora indica* fungi enhance the efficiency of chlorophyll fluorescence and essential oil composition in peppermint under irrigation with saline water from the Caspian Sea? *PLoS One* 2021;16:1–21. <https://doi.org/10.1371/journal.pone.0254076>.
- Egamberdieva D, Wirth S, Bellingrath-Kimura SD, Mishra J, Arora NK. Salt-tolerant plant growth promoting rhizobacteria for enhancing crop productivity of saline soils. *Front Microbiol* 2019;10. <https://doi.org/10.3389/fmicb.2019.02791>.
- Deinlein U, Stephan AB, Horie T, Luo W, Xu G, Schroeder JI. Plant salt-tolerance mechanisms. *Trends Plant Sci* 2014;19:371–9. <https://doi.org/10.1016/j.tplants.2014.02.001>.
- Lamers J, Der Meer TV, Testerink C. How plants sense and respond to stressful environments. *Plant Physiol* 2020;182:1624–35. <https://doi.org/10.1104/PP.19.01464>.
- Liang W, Ma X, Wan P, Liu L. Plant salt-tolerance mechanism: A review. *Biochem Biophys Res Commun* 2018;495:286–91. <https://doi.org/10.1016/j.bbrc.2017.11.043>.
- Ondrasek G, Rathod S, Manohara KK, Gireesh C, Anantha MS, Sakhar AS, et al. Salt stress in plants and mitigation approaches. *Plants* 2022;11:1–21. <https://doi.org/10.3390/plants11060717>.
- Zhu J-K. Plant salt tolerance. *Trends Plant Sci* 2001;6:66–71.
- Zhu JK. Salt and drought stress signal transduction in plants. *Annu Rev Plant Biol* 2002;53:247–73. <https://doi.org/10.1146/annurev.arplant.53.091401.143329>.
- Zhu J-K. Abiotic stress signaling and responses in plants. *Cell* 2016;167:313–24. <https://doi.org/10.1016/j.cell.2016.08.029>.
- Evelin H, Devi TS, Gupta S, Kapoor R. Mitigation of salinity stress in plants by arbuscular mycorrhizal symbiosis: Current understanding and new challenges. *Frontiers. Plant Sci* 2019;10. <https://doi.org/10.3389/fpls.2019.00470>.
- Aroca R, Ruiz-Lozano JM, Zamarreño ángel M, Paz JA, García-Mina JM, Pozo MJ, et al. Arbuscular mycorrhizal symbiosis influences strigolactone production under salinity and alleviates salt stress in lettuce plants. *Journal of Plant Physiology* 2013;170:47–55. [10.1016/j.jplph.2012.08.020](https://doi.org/10.1016/j.jplph.2012.08.020).
- Feng G, Zhang FS, Li XL, Tian CY, Tang C, Rengel Z. Improved tolerance of maize plants to salt stress by arbuscular mycorrhiza is related to higher accumulation of soluble sugars in roots. *Mycorrhiza* 2002;12:185–90. <https://doi.org/10.1007/s00572-002-0170-0>.
- Eroglu G, Cabral C, Ravnskov S, Bak Topbjerg H, Wollenweber B. Arbuscular mycorrhiza influences carbon-use efficiency and grain yield of wheat grown under pre- and post-anthesis salinity stress. *Plant Biol* 2020;22:863–71. <https://doi.org/10.1111/plb.13123>.
- Parvin S, Van Geel M, Yeasmin T, Verbruggen E, Honnay O. Effects of single and multiple species inocula of arbuscular mycorrhizal fungi on the salinity tolerance of a Bangladeshi rice (*Oryza sativa* L.) cultivar. *Mycorrhiza* 2020;30:431–44. <https://doi.org/10.1007/s00572-020-00957-9>.
- Yamato M, Ikeda S, Iwase K. Community of arbuscular mycorrhizal fungi in a coastal vegetation on Okinawa island and effect of the isolated fungi on growth of sorghum under salt-treated conditions. *Mycorrhiza* 2008;18:241–9. <https://doi.org/10.1007/s00572-008-0177-2>.
- Al-Karaki GN. Growth of mycorrhizal tomato and mineral acquisition under salt stress. *Mycorrhiza* 2000;10:51–4. <https://doi.org/10.1007/s005720000055>.
- Romero-Munar A, Baraza E, Gullás J, Cabot C. Arbuscular mycorrhizal fungi confer salt tolerance in giant reed (*Arundo donax* L.) plants grown under low phosphorus by reducing leaf N⁺ concentration and improving phosphorus use efficiency. *Frontiers. Plant Sci* 2019;10:1–14. <https://doi.org/10.3389/fpls.2019.00843>.
- Duc NH, Vo AT, Haddidi I, Daoud H, Posta K. Arbuscular mycorrhizal fungi improve tolerance of the medicinal plant *Eclipta prostrata* (L.) and induce major changes in polyphenol profiles under salt stresses. *Frontiers. Plant Sci* 2021;11:1–18. <https://doi.org/10.3389/fpls.2020.612299>.
- Giri B, Kapoor R, Mukerji KG. Improved tolerance of *Acacia nilotica* to salt stress by arbuscular mycorrhiza. *glomus fasciculatum* may be partly related to elevated K/Na ratios in root and shoot tissues. *Microb Ecol* 2007;54:753–60. <https://doi.org/10.1007/s00248-007-9239-9>.
- Wang J, Zhai L, Ma J, Zhang J, Wang GG, Liu X, et al. Comparative physiological mechanisms of arbuscular mycorrhizal fungi in mitigating salt-induced adverse effects on leaves and roots of *Zelkova serrata*. *Mycorrhiza* 2020;30:341–55. <https://doi.org/10.1007/s00572-020-00954-y>.
- Wang Y, Wang M, Li Y, Wu A, Huang J. Enhancements of arbuscular mycorrhizal fungi on growth and nitrogen acquisition of *Chrysanthemum morifolium* under salt stress. *PLoS One* 2017;35:36.
- Pandey R, Garg N. High effectiveness of *Rhizopagus irregularis* is linked to superior modulation of antioxidant defence mechanisms in *Cajanus cajan* (L.) Millsp. genotypes grown under salinity stress. *Mycorrhiza* 2017;27:669–82. <https://doi.org/10.1007/s00572-017-0778-8>.
- Harman GE, Howell CR, Viterbo A, Chet I, Lorito M. *Trichoderma* species—opportunistic, avirulent plant symbionts. *Nat Rev Microbiol* 2004;2:43–56. <https://doi.org/10.1038/nrmicro797>.
- Brotman Y, Landau U, Cuadros-Inostroza Á, Takayuki T, Fernie AR, Chet I, et al. *Trichoderma*-plant root colonization: escaping early plant defense responses and activation of the antioxidant machinery for saline stress tolerance. *PLoS Pathog* 2013;9:e1003221–e. <https://doi.org/10.1371/journal.ppat.1003221>.
- Zhang F, Wang Y, Liu C, Chen F, Ge H, Tian F, et al. *Trichoderma harzianum* mitigates salt stress in cucumber via multiple responses. *Ecotoxicol Environ Saf* 2019;170:436–45. <https://doi.org/10.1016/j.ecoenv.2018.11.084>.
- Gupta S, Smith PMC, Boughton BA, Rupasinghe TWT, Natera SHA, Roessner U. Inoculation of barley with *Trichoderma harzianum* T-22 modifies lipids and metabolites to improve salt tolerance. *J Exp Bot* 2021;72:7229–46. <https://doi.org/10.1093/jxb/erab335>.
- Contreras-Cornejo HA, Macías-Rodríguez L, Alfaro-Cuevas R, López-Bucio J. *Trichoderma* spp. improve growth of *Arabidopsis* seedlings under salt stress through enhanced root development, osmolyte production, and Na⁺ elimination through root exudates. *Molecular Plant-Microbe Interactions* 2014;27:503–14. [10.1094/MPMI-09-13-0265-R](https://doi.org/10.1094/MPMI-09-13-0265-R).
- Zhang S, Gan Y, Xu B. Mechanisms of the IAA and ACC-deaminase producing strain of *Trichoderma longibrachiatum* T6 in enhancing wheat seedling tolerance to NaCl stress. *BMC Plant Biology* 2019;19:22–22. [10.1186/s12870-018-1618-5](https://doi.org/10.1186/s12870-018-1618-5).

- [38] Franken P. The plant strengthening root endophyte *Piriformospora indica*: Potential application and the biology behind. *Appl Microbiol Biotechnol* 2012;96:1455–64. <https://doi.org/10.1007/s00253-012-4506-1>.
- [39] Gill SS, Gill R, Trivedi DK, Anjum NA, Sharma KK, Ansari MW, et al. *Piriformospora indica*: potential and significance in plant stress tolerance. *Frontiers in Microbiology* 2016;7:332–332. 10.3389/fmicb.2016.00332.
- [40] Jisha S, Sabu KK. Multifunctional aspects of *Piriformospora indica* in plant endosymbiosis. *Mycology* 2019;10:182–90. <https://doi.org/10.1080/21501203.2019.1600063>.
- [41] Vahabi K, Dorcheh SK, Monajembashi S, Westermann M, Reichelt M, Falkenberg D, et al. Stress promotes *Arabidopsis*-*Piriformospora indica* interaction. *Plant Signal Behav* 2016;11. <https://doi.org/10.1080/15592324.2015.1136763>.
- [42] Jogawat A, Saha S, Bakshi M, Dayaman V, Kumar M, Dua M, et al. *Piriformospora indica* rescues growth diminution of rice seedlings during high salt stress. *Plant Signal Behav* 2013;8:e26891–e. <https://doi.org/10.4161/psb.26891>.
- [43] Waller F, Achatz B, Baltruschat H, Fodor J, Becker K, Fischer M, et al. The endophytic fungus *Piriformospora indica* reprograms barley to salt stress tolerance, disease resistance, and higher yield. *Proceedings of the National Academy of Sciences of the United States of America* 2005;102:13386–91. 10.1073/pnas.0504423102.
- [44] Ghorbani A, Razavi SM, Ghasemi Omran VO, Pirdashti H. *Piriformospora indica* inoculation alleviates the adverse effect of NaCl stress on growth, gas exchange and chlorophyll fluorescence in tomato (*Solanum lycopersicum* L.). *Plant Biol* 2018;20:729–36. <https://doi.org/10.1111/plb.12717>.
- [45] Lanza M, Haro R, Conchillo LB, Benito B. The endophyte *Serendipita indica* reduces the sodium content of *Arabidopsis* plants exposed to salt stress: fungal ENA ATPases are expressed and regulated at high pH and during plant co-cultivation in salinity. *Environ Microbiol* 2019;21:3364–78. <https://doi.org/10.1111/1462-2920.14619>.
- [46] Kim K, Jang Y-J, Lee S-M, Oh B-T, Chae J-C, Lee K-J. Alleviation of salt stress by enterobacter sp. EJ01 in tomato and Arabidopsis is accompanied by up-regulation of conserved salinity responsive factors in plants. *Molecules and Cells* 2014;37:109–17. 10.14348/molcells.2014.2239.
- [47] Khan MA, Asaf S, Khan AL, Adhikari A, Jan R, Ali S, et al. Plant growth-promoting endophytic bacteria augment growth and salinity tolerance in rice plants. *Plant Biol* 2020;22:850–62. <https://doi.org/10.1111/plb.13124>.
- [48] Singh RP, Pandey DM, Jha PN, Ma Y. ACC deaminase producing rhizobacterium *Enterobacter cloacae* ZNP-4 enhance abiotic stress tolerance in wheat plant. *PLoS One* 2022;17:1–23. <https://doi.org/10.1371/journal.pone.0267127>.
- [49] Nadeem SM, Zahir ZA, Naveed M, Arshad M. Rhizobacteria containing ACC-deaminase confer salt tolerance in maize grown on salt-affected fields. *Can J Microbiol* 2009;55:1302–9. <https://doi.org/10.1139/W09-092>.
- [50] Desai S, Mistry J, Shah F, Chandwani S, Amarean N, Supriya NR. Salt-tolerant bacteria enhance the growth of mung bean (*Vigna radiata* L.) and uptake of nutrients, and mobilize sodium ions under salt stress condition. *Int J Phytorem* 2022;1–8. <https://doi.org/10.1080/15226514.2022.2057419>.
- [51] Habib SH, Kausar H, Saud HM. Plant growth-promoting rhizobacteria enhance salinity stress tolerance in *Okra* through ROS-scavenging enzymes. *Biomed Res Int* 2016;1–10. <https://doi.org/10.1155/2016/6284547>.
- [52] Yang A, Akhtar SS, Iqbal S, Amjad M, Naveed M, Zahir ZA, et al. Enhancing salt tolerance in quinoa by halotolerant bacterial inoculation. *Funct Plant Biol* 2016;43:632–42. <https://doi.org/10.1071/FP15265>.
- [53] Nadeem SM, Zahir ZA, Naveed M, Arshad M. Preliminary investigations on inducing salt tolerance in maize through inoculation with rhizobacteria containing ACC deaminase activity. *Can J Microbiol* 2007;53:1141–9. <https://doi.org/10.1139/W07-081>.
- [54] de Zélicourt A, Synek L, Saad MM, Alzubaidy H, Jalal R, Xie Y, et al. Ethylene induced plant stress tolerance by *Enterobacter* sp. SA187 is mediated by 2-keto-4-methylthiobutyric acid production. *PLoS Genet* 2018;14:1–28. <https://doi.org/10.1371/journal.pgen.1007273>.
- [55] Ali B, Wang X, Saleem MH, Sumaira, Hafeez A, Afridi MS, et al. PGPR-mediated salt tolerance in maize by modulating plant physiology, antioxidant defense, compatible solutes accumulation and bio-surfactant producing genes. *Plants* 2022;11:345–345. 10.3390/plants11030345.
- [56] Nazarowec-White M, Farber JM. *Enterobacter sakazakii*: A review. *Int J Food Microbiol* 1997;34:103–13. [https://doi.org/10.1016/S0168-1605\(96\)01172-5](https://doi.org/10.1016/S0168-1605(96)01172-5).
- [57] Subramanian S, Ricci E, Souleimanov A, Smith DL. A proteomic approach to lipo-chitoooligosaccharide and thuricin 17 effects on soybean germination unstressed and salt stress. *PLoS One* 2016;11:1–19. <https://doi.org/10.1371/journal.pone.0160660>.
- [58] Subramanian S, Souleimanov A, Smith DL. Proteomic studies on the effects of lipo-chitoooligosaccharide and thuricin 17 under unstressed and salt stressed conditions in *Arabidopsis thaliana*. *Front Plant Sci* 2016;7:1–13. <https://doi.org/10.3389/fpls.2016.01314>.
- [59] Ullah S, Bano A. Isolation of plant-growth-promoting rhizobacteria from rhizospheric soil of halophytes and their impact on maize (*Zea mays* L.) under induced soil salinity. *Can J Microbiol* 2015;61:307–13. <https://doi.org/10.1139/cjm-2014-0668>.
- [60] Mahmoud OMB, Slimene IB, Zribi OT, Abdellly C, Djébali N. Response to salt stress is modulated by growth-promoting rhizobacteria inoculation in two contrasting barley cultivars. *Acta Physiol Plant* 2017;39. <https://doi.org/10.1007/s11738-017-2421-x>.
- [61] Sultana S, Paul SC, Parveen S, Alam S, Rahman N, Jannat B, et al. Isolation and identification of salt-tolerant plant-growth-promoting rhizobacteria and their application for rice cultivation under salt stress. *Can J Microbiol* 2020;66:144–60. <https://doi.org/10.1139/cjm-2019-0323>.
- [62] Wang R, Wang C, Feng Q, Liou RM, Lin YF. Biological inoculant of salt-tolerant bacteria for plant growth stimulation under different saline soil conditions. *J Microbiol Biotechnol* 2021;31:398–407. <https://doi.org/10.4014/JMB.2009.09032>.
- [63] Fan P, Chen D, He Y, Zhou Q, Tian Y, Gao L. Alleviating salt stress in tomato seedlings using *Arthrobacter* and *Bacillus megaterium* isolated from the rhizosphere of wild plants grown on saline-alkaline lands. *Int J Phytorem* 2016;18:1113–21. <https://doi.org/10.1080/15226514.2016.1183583>.
- [64] Egamberdieva D, Wirth SJ, Shurigin VV, Hashem A, Abd Allah EF. Endophytic bacteria improve plant growth, symbiotic performance of chickpea (*Cicer arietinum* L.) and induce suppression of root rot caused by *Fusarium solani* under salt stress. *Front Microbiol* 2017;8:1–13. <https://doi.org/10.3389/fmicb.2017.01887>.
- [65] Saleem S, Iqbal A, Ahmed F, Ahmad M. Phytobeneficial and salt stress mitigating efficacy of IAA producing salt tolerant strains in *Gossypium hirsutum*. *Saudi Journal of Biological Sciences* 2021;28:5317–24. <https://doi.org/10.1016/j.sjbs.2021.05.056>.
- [66] Khan MA, Sahile AA, Jan R, Asaf S, Hamayun M, Imran M, et al. Halotolerant bacteria mitigate the effects of salinity stress on soybean growth by regulating secondary metabolites and molecular responses. *BMC Plant Biol* 2021;21:1–15. <https://doi.org/10.1186/s12870-021-02937-3>.
- [67] Srivastava S, Srivastava S. Prescience of endogenous regulation in *Arabidopsis thaliana* by *Pseudomonas putida* MTCC 5279 under phosphate starved salinity stress condition. *Sci Rep* 2020;10:1–15. <https://doi.org/10.1038/s41598-020-62725-1>.
- [68] Singh S, Singh UB, Trivedi M, Sahu PK, Paul S, Paul D, et al. Seed biopriming with salt-tolerant endophytic pseudomonas geniculata-modulated biochemical responses provide ecological fitness in maize (*Zea mays* L.) grown in saline sodic soil. *Int J Environ Res Public Health* 2020;17. <https://doi.org/10.3390/ijerph17010253>.
- [69] Yuan P, Pan H, Boak EN, Pierson LS, Pierson EA. Phenazine-producing rhizobacteria promote plant growth and reduce redox and osmotic stress in wheat seedlings under saline conditions. *Front Plant Sci* 2020;11:1–12. <https://doi.org/10.3389/fpls.2020.575314>.
- [70] Yao L, Wu Z, Zheng Y, Kaleem I, Li C. Growth promotion and protection against salt stress by *Pseudomonas putida* Rs-198 on cotton. *Eur J Soil Biol* 2010;46:49–54. <https://doi.org/10.1016/j.ejsobi.2009.11.002>.
- [71] Nascimento FX, Urón P, Glick BR, Giachini A, Rossi MJ. Genomic analysis of the 1-aminocyclopropane-1-carboxylate deaminase-producing *Pseudomonas thirervallensis* sc5 reveals its multifaceted roles in soil and in beneficial interactions with plants. *Front Microbiol* 2021;12. <https://doi.org/10.3389/fmicb.2021.752288>.
- [72] Sharma S, Kulkarni J, Jha B. Halotolerant rhizobacteria promote growth and enhance salinity tolerance in peanut. *Front Microbiol* 2016;7:1–11. <https://doi.org/10.3389/fmicb.2016.01600>.
- [73] Vaishnav A, Kumari S, Jain S, Varma A, Tuteja N, Choudhary DK. PGPR-mediated expression of salt tolerance gene in soybean through volatiles under sodium nitroprusside. *J Basic Microbiol* 2016;56:1274–88. <https://doi.org/10.1002/jobm.201600188>.
- [74] Vaishnav A, Kumari S, Jain S, Varma A, Choudhary DK. Putative bacterial volatile-mediated growth in soybean (*Glycine max* L. Merrill) and expression of induced proteins under salt stress. *J Appl Microbiol* 2015;119:539–51. <https://doi.org/10.1111/jam.12866>.
- [75] Ali S, Charles TC, Glick BR. Amelioration of high salinity stress damage by plant growth-promoting bacterial endophytes that contain ACC deaminase. *Plant Physiol Biochem* 2014;80:160–7. <https://doi.org/10.1016/j.plaphy.2014.04.003>.
- [76] Fatima T, Arora NK. *Pseudomonas entomophila* PE3 and its exopolysaccharides as biostimulants for enhancing growth, yield and tolerance responses of sunflower under saline conditions. *Microbiological Research* 2021;244:126671–126671. 10.1016/j.micres.2020.126671.
- [77] Chang P, Gerhardt KE, Huang XD, Yu XM, Glick BR, Gerwing PD, et al. Plant growth-promoting bacteria facilitate the growth of barley and oats in salt-impacted soil: implications for phytoremediation of saline soils. *Int J Phytorem* 2014;16:1133–47. <https://doi.org/10.1080/15226514.2013.821447>.
- [78] Rus A, Lee BH, Muñoz-Mayor A, Sharkhuu A, Miura K, Zhu JK, et al. AtHKT1 facilitates Na⁺ homeostasis and K⁺ nutrition in planta. *Plant Physiol* 2004;136:2500–11. <https://doi.org/10.1104/pp.104.042234>.
- [79] Platten JD, Cotsaftis O, Berthomieu P, Bohnert H, Davenport RJ, Fairbairn DJ, et al. Nomenclature for HKT transporters, key determinants of plant salinity tolerance. *Trends Plant Sci* 2006;11:372–4. <https://doi.org/10.1016/j.tplants.2006.06.001>.
- [80] Moller IS, Gilliam M, Jha D, Mayo GM, Roy SJ, Coates JC, et al. Shoot Na⁺ exclusion and increased salinity tolerance engineered by cell type-specific alteration of Na⁺ transport in *Arabidopsis*. *Plant Cell* 2009;21:2163–78. <https://doi.org/10.1105/tpc.108.064568>.
- [81] Zhang H, Kim MS, Sun Y, Dowd SE, Shi H, Paré PW. Soil bacteria confer plant salt tolerance by tissue-specific regulation of the sodium transporter HKT1. *Mol Plant Microbe Interact* 2008;21:737–44. <https://doi.org/10.1094/MPMI-21-6-0737>.

- [82] Niu SQ, Li HR, Paré PW, Aziz M, Wang SM, Shi H, et al. Induced growth promotion and higher salt tolerance in the halophyte grass *Puccinellia tenuiflora* by beneficial rhizobacteria. *Plant and Soil* 2016;407:217–30. <https://doi.org/10.1007/s11104-015-2767-z>.
- [83] Abdelaziz ME, Kim D, Ali S, Fedoroff NV, Al-Babili S. The endophytic fungus *Piriformospora indica* enhances *Arabidopsis thaliana* growth and modulates Na⁺/K⁺ homeostasis under salt stress conditions. *Plant Sci* 2017;263:107–15. <https://doi.org/10.1016/j.plantsci.2017.07.006>.
- [84] Santander C, Aroca R, Cartes P, Vidal G, Cornejo P. Aquaporins and cation transporters are differentially regulated by two arbuscular mycorrhizal fungi strains in lettuce cultivars growing under salinity conditions. *Plant Physiol Biochem* 2021;158:396–409. <https://doi.org/10.1016/j.plaphy.2020.11.025>.
- [85] Apse MP, Aharon GS, Snedden WA, Blumwald E. Salt tolerance conferred by overexpression of a vacuolar Na⁺/H⁺ antiporter in *Arabidopsis*. *Science* 1999;285:1256–8. <https://doi.org/10.1126/science.285.5431.1256>.
- [86] Drodzowicz Y, Rea P. Vacuolar H⁺ pyrophosphatases: from the evolutionary backwaters into the mainstream. *Trends Plant Sci* 2001;6:206–11. [https://doi.org/10.1016/S1360-1385\(01\)01923-9](https://doi.org/10.1016/S1360-1385(01)01923-9).
- [87] Chen L, Liu Y, Wu G, Veronican Njeri K, Shen Q, Zhang N, et al. Induced maize salt tolerance by rhizosphere inoculation of *Bacillus amyloliquefaciens* SQR9. *Physiol Plant* 2016;158:34–44. <https://doi.org/10.1111/ppi.12441>.
- [88] Chen L, Liu Y, Wu G, Zhang N, Shen Q, Zhang R. Beneficial rhizobacterium *Bacillus amyloliquefaciens* SQR9 induces plant salt tolerance through spermidine production. *Mol Plant Microbe Interact* 2017;30:423–32. <https://doi.org/10.1094/MPMI-07-17-0027-R>.
- [89] Baek D, Rokibuzzaman M, Khan A, Kim MC, Park HJ, Yun DJ, et al. Plant-growth promoting *Bacillus oryzae* yc7007 modulates stress-response gene expression and provides protection from salt stress. *Front Plant Sci* 2020;10:1–13. <https://doi.org/10.3389/fpls.2019.01646>.
- [90] Del Carmen O-M, Duan J, DiBernardo M, Zetter E, Campos-García J, Glick BR, et al. The production of ACC deaminase and trehalose by the plant growth promoting bacterium *Pseudomonas* sp. UW4 synergistically protect tomato plants against salt stress. *Front Microbiol* 2019;10:1–10. <https://doi.org/10.3389/fmicb.2019.01392>.
- [91] Yan J, Smith MD, Glick BR, Liang Y. Effects of ACC deaminase containing rhizobacteria on plant growth and expression of Toc GTPases in tomato (*Solanum lycopersicum*) under salt stress. *Botany* 2014;92:775–81. <https://doi.org/10.1139/cjb-2014-0038>.
- [92] Hedrich R, Shabala S. Stomata in a saline world. *Curr Opin Plant Biol* 2018;46:87–95. <https://doi.org/10.1016/j.pbi.2018.07.015>.
- [93] Zhou C, Zhu L, Xie Y, Li F, Xiao X, Ma Z, et al. *Bacillus licheniformis* SA03 confers increased saline-alkaline tolerance in *chrysanthemum* plants by induction of abscisic acid accumulation. *Front Plant Sci* 2017;8:1–17. <https://doi.org/10.3389/fpls.2017.01143>.
- [94] Sukweenadhi J, Balusamy SR, Kim YJ, Lee CH, Kim YJ, Koh SC, et al. A growth-promoting bacteria, *Paenibacillus yonginensis* DCY84T enhanced salt stress tolerance by activating defense-related systems in panax ginseng. *Front Plant Sci* 2018;9:1–17. <https://doi.org/10.3389/fpls.2018.00813>.
- [95] Yoo SJ, Weon HY, Song J, Sang MK. Induced tolerance to salinity stress by halotolerant bacteria *Bacillus aryabhattai* H19–1 and *B. mesonae* H20–5 in tomato plants. *J Microbiol Biotechnol* 2019;29:1124–36. <https://doi.org/10.4014/jmb.1904.04026>.
- [96] Lastochkina O, Aliniaieifard S, Garshina D, Garipova S, Pusenkova L, Allagulova C, et al. Seed priming with endophytic *Bacillus subtilis* strain-specifically improves growth of *Phaseolus vulgaris* plants under normal and salinity conditions and exerts anti-stress effect through induced lignin deposition in roots and decreased oxidative and osmot. *Journal of Plant Physiology* 2021;263:153462–153462. <https://doi.org/10.1016/j.jplph.2021.153462>.
- [97] Gul Jan F, Hamayun M, Hussain A, Jan G, Iqbal A, Khan A, et al. An endophytic isolate of the fungus *Yarrowia lipolytica* produces metabolites that ameliorate the negative impact of salt stress on the physiology of maize. *BMC Microbiol* 2019;19:1–10. <https://doi.org/10.1186/s12866-018-1374-6>.
- [98] Liu CH, Siew W, Hung YT, Jiang YT, Huang CH. 1-aminocyclopropane-1-carboxylate (ACC) deaminase gene in *Pseudomonas azotoformans* is associated with the amelioration of salinity stress in tomato. *J Agric Food Chem* 2021;69:913–21. <https://doi.org/10.1021/acs.jafc.0c05628>.
- [99] Kavi Kishor PB, Sreenivasulu N. Is proline accumulation per se correlated with stress tolerance or is proline homeostasis a more critical issue? *Plant Cell Environ* 2014;37:300–11. <https://doi.org/10.1111/pce.12157>.
- [100] Xu Z, Lei P, Pang X, Li H, Feng X, Xu H. Exogenous application of poly-γ-glutamic acid enhances stress defense in *Brassica napus* L. seedlings by inducing cross-talks between Ca²⁺, H₂O₂, brassinolide, and jasmonic acid in leaves. *Plant Physiol Biochem* 2017;118:460–70. <https://doi.org/10.1016/j.plaphy.2017.07.015>.
- [101] Lei P, Pang X, Feng X, Li S, Chi B, Wang R, et al. The microbe-secreted isopeptide poly-γ-glutamic acid induces stress tolerance in *Brassica napus* L. seedlings by activating crosstalk between H₂O₂ and Ca²⁺. *Sci Rep* 2017;7:1–15. <https://doi.org/10.1038/srep41618>.
- [102] Gad M, Nobuhiro S, Sultan C, Ron M. Reactive oxygen species homeostasis and signalling during drought and salinity stresses. *Plant Cell and Environment* 2010;33:453–467. <https://doi.org/10.1111/j.1365-3040.2009.02041.x>.
- [103] Das K, Roychoudhury A. Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Frontiers in Environmental. Science* 2014;2. <https://doi.org/10.3389/fenvs.2014.00053>.
- [104] Choudhury FK, Rivero RM, Blumwald E, Mittler R. Reactive oxygen species, abiotic stress and stress combination. *Plant J* 2017;90:856–67. <https://doi.org/10.1111/tpj.13299>.
- [105] Apel K, Hirt H. Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol* 2004;55:373–99. <https://doi.org/10.1146/annurev.arplant.55.031903.141701>.
- [106] Estrada B, Aroca R, Barea JM, Ruiz-Lozano JM. Native arbuscular mycorrhizal fungi isolated from a saline habitat improved maize antioxidant systems and plant tolerance to salinity. *Plant Sci* 2013;201–202:42–51. <https://doi.org/10.1016/j.plantsci.2012.11.009>.
- [107] Estrada B, Aroca R, Maathuis FJM, Barea JM, Ruiz-Lozano JM. Arbuscular mycorrhizal fungi native from a Mediterranean saline area enhance maize tolerance to salinity through improved ion homeostasis. *Plant Cell Environ* 2013;36:1771–82. <https://doi.org/10.1111/pce.12082>.
- [108] Andres-Barrao C, Alzubaidy H, Jalal R, Mariappan KG, De Zelicourt A, Bokhari A, et al. Coordinated bacterial and plant sulfur metabolism in *Enterobacter* sp. SA187-induced plant salt stress tolerance. *Proceedings of the National Academy of Sciences of the United States of America* 2021;118:1–12. <https://doi.org/10.1073/pnas.2107417118>.
- [109] Chatterjee P, Samaddar S, Niinemets Ü, Sa TM. *Brevibacterium linens* RS16 confers salt tolerance to *Oryza sativa* genotypes by regulating antioxidant defense and H⁺ ATPase activity. *Microbiol Res* 2018;215:89–101. <https://doi.org/10.1016/j.micres.2018.06.007>.
- [110] Chen J, Zhang H, Zhang X, Tang M. Arbuscular mycorrhizal symbiosis alleviates salt stress in black locust through improved photosynthesis, water status, and K⁺/Na⁺ homeostasis. *Front Plant Sci* 2017;8:1–14. <https://doi.org/10.3389/fpls.2017.01739>.
- [111] Bishop A, Benková E, Helariutta Y. Sending mixed messages: Auxin-cytokinin crosstalk in roots. *Curr Opin Plant Biol* 2011;14:10–6. <https://doi.org/10.1016/j.pbi.2010.08.014>.
- [112] Barnawal D, Bharti N, Pandey SS, Pandey A, Chanotiya CS, Kalra A. Plant growth-promoting rhizobacteria enhance wheat salt and drought stress tolerance by altering endogenous phytohormone levels and TaCTR1/TaDREB2 expression. *Physiol Plant* 2017;161:502–14. <https://doi.org/10.1111/ppi.12614>.
- [113] Li Y, Shao J, Xie Y, Jia L, Fu Y, Xu Z, et al. Volatile compounds from beneficial rhizobacteria *Bacillus* spp. promote periodic lateral root development in *Arabidopsis*. *Plant Cell Environ* 2021;44:1663–78. <https://doi.org/10.1111/pce.14021>.
- [114] Jaroszuk-Ścisł J, Tyśkiewicz R, Nowak A, Ozimek E, Majewska M, Hanaka A, et al. Phytohormones (Auxin, Gibberellin) and ACC deaminase in vitro synthesized by the mycoparasitic *Trichoderma* DENTK23A0 strain and changes in the level of auxin and plant resistance markers in wheat seedlings inoculated with this strain conidia. *Int J Mol Sci* 2019;20:E4923. <https://doi.org/10.3390/ijms20194923>.
- [115] Kazan K. Diverse roles of jasmonates and ethylene in abiotic stress tolerance. *Trends Plant Sci* 2015;20:219–29. <https://doi.org/10.1016/j.tplants.2015.02.001>.
- [116] Gamalero E, Glick BR. Bacterial modulation of plant ethylene levels. *Plant Physiol* 2015;169:13–22. <https://doi.org/10.1104/pp.15.00284>.
- [117] Tavares MJ, Nascimento FX, Glick BR, Rossi MJ. The expression of an exogenous ACC deaminase by the endophyte *Serratia grimesii* BXF1 promotes the early nodulation and growth of common bean. *Lett Appl Microbiol* 2018;66:252–9. <https://doi.org/10.1111/lam.12847>.
- [118] Singh RP, Jha PN. The PGPR *Stenotrophomonas maltophilia* SBP-9 augments resistance against biotic and abiotic stress in wheat plants. *Frontiers in Microbiology* 2017;8:1945–1945. <https://doi.org/10.3389/fmicb.2017.01945>.
- [119] Glick BR. The enhancement of plant growth by free-living bacteria. *Can J Microbiol* 1995;41:109–17. <https://doi.org/10.1139/m95-015>.
- [120] Heydarian Z, Gruber M, Glick BR, Hegedus DD. Gene expression patterns in roots of *Camelina sativa* with enhanced salinity tolerance arising from inoculation of soil with plant growth promoting bacteria producing 1-aminocyclopropane-1-carboxylate deaminase or expression the corresponding acs gene. *Front Microbiol* 2018;9:1–15. <https://doi.org/10.3389/fmicb.2018.01297>.
- [121] Yoolong S, Kruasuan W, Thanh Pham HT, Jaemsang R, Jantasuriyarat C, Thachapenat A. Modulation of salt tolerance in Thai jasmine rice (*Oryza sativa* L. cv. KDML105) by *Streptomyces venezuelae* ATCC 10712 expressing ACC deaminase. *Sci Rep* 2019;9:1–10. <https://doi.org/10.1038/s41598-018-37987-5>.
- [122] Loo E-P-I, Tajima Y, Yamada K, Kido S, Hirase T, Ariga H, et al. Recognition of microbe- and damage-associated molecular patterns by leucine-rich repeat pattern recognition receptor kinases confers salt tolerance in plants. *Molecular Plant-Microbe Interactions* 2022;35(7):554–66. <https://doi.org/10.1094/mpmi-07-21-0185-fj>.
- [123] Rolli E, de Zelicourt A, Alzubaidy H, Karampelias M, Parween S, Rayapuram N, et al. The Lys-motif receptor LYK4 mediates *Enterobacter* sp. SA187 triggered salt tolerance in *Arabidopsis thaliana*. *Environ Microbiol* 2022;24:223–39. <https://doi.org/10.1111/1462-2920.15839>.