Planar universal optimality, tiered: kernel-checked consequences, a non-uniqueness theorem, and a CVODE N-body consistency check for the Yukawa interaction

Xavier Callens (SocrateAI Lab)

Preprint v2, 2026-10-11 (v1: doi:10.5281/zenodo.23293112)




Figure 1: Cover. Three configurations from the $\kappa = 2$ CVODE study (Tier X), with Voronoi cells coloured by local $|\psi_6|$. Dots mark sites that are not 6-fold under the $1.35\,d_0$ rule (5 blue, 7 red, other white). Left: the OC20-derived initial condition `oc20_00`, in which every site is flagged. Middle: its $T = 0$ CVODE quench, a defected local minimum. Right: a *different* initial condition, `oc20_17`, annealed and then quenched, which reaches the triangular lattice. That outcome is uncommon: only 2 of 40 annealed runs at $\kappa = 2$ reach it (Table 2). Each panel is regenerated from a published seed by `yukawa_crystal snapshot`, and its energy equals the published value.

Abstract

The openai/math release contains a Lean 4 proof that the triangular lattice minimises the lower g-energy among all locally finite planar configurations of density one, for every smooth completely monotone g [1]. We file every claim we make about it in a claim ledger with the tiers of the SocrateAI-Scientific-Elenchus method [4]: **A** kernel-checked, **L** cited literature, **C** argument, **X** floating-point exploration, which may steer but may never support a claim. At Tier A, conditional only on the fidelity of upstream's definitions and of our transcription, we prove in Lean that the rescaled triangular lattice is universally optimal at every density in the sense of Cohn–Kumar–Miller–Radchenko–Viazovska's (CKMRV) Definition 1.3, stated with upstream's lower energy. That is weaker than their notion, which also requires the energy to exist

as a limit. We also formalise the remark that follows CKMRV's Theorem 1.4, which upstream also notes: universal optima are not unique among all configurations, because the lattice with one vacancy is another and is not an isometric image of the lattice. In this formulation only the minimum value is determined, not the minimiser. At Tier X we run an overdamped N-body relaxation for the 2D Yukawa interaction, which is in the theorem's class (Tier C), with the CVODE solver of rusty-SUNDIALS, from initial conditions derived from the Hugging Face OC20 dataset. The preregistered falsifiable prediction, that no relaxed periodic configuration falls below the triangular-lattice energy, is reported with its controls. It is consistency evidence, not proof. The Elenchus ledger gate reports 0 blocking findings and 6 flags (exit 1). Each flag marks a Tier A row whose statement adequacy no human has audited, so by the method's own publish gate the ledger does not yet pass.

1 Claims and tiers

Table 1 lists every claim made below, with its tier and its ledger identifier.

Ledger id	Tier	Claim (abbreviated; full statements in `ledger.json`)
UO-A-0001	A	upstream `universal_energy_minimum` (our compile and footprint; definitions AI-audited only)
UO-A-0002–4	A	Riesz admissibility and optimality; finiteness for $s > 2$; every density with attainment [3]
UO-A-0005	A	`triangular_universally_optimal`: $\rho^{-1/2}A$ is `UniversallyOptimal` for every $\rho > 0$
UO-A-0006	A	`universal_optimum_not_unique`: $A \setminus \{0\}$ is universally optimal and not an isometric image of A
UO-L-0001–6	L	CKMRV Definition 1.3, uniqueness remark, history, composition footnote, lattice energy [2]; upstream scope remark [1]
UO-C-0001	C	the 2D Yukawa interaction is completely monotone in r^2
UO-C-0002	C	the interpretation of UO-A-0006 (what it does and does not refute)
UO-X-000n	X	CVODE N-body study, controls, disk study (`claim_licence: false`)

Table 1: The claim ledger. A row may not be cited above its tier, and no row is filed above anything it rests on; both are checked mechanically, transitively, by the Elenchus gate. Every row points to a content-addressed evidence blob (compiled axiom output with source hashes, primary-source excerpts with line numbers, argument text, or result files).

2 Tier A: what the kernel checked

Upstream (UO-A-0001). With A the triangular lattice of covolume one, $N_R(C)$ the number of points of C in the closed disk of radius R, density $\lim N_R/(\pi R^2)$, and lower energy $E_g(C) = \liminf_R N_R^{-1} \sum_{x \neq y} g(|x-y|^2)$ in $[0, \infty]$, upstream proves $\sum_{a \in A \setminus 0} g(|a|^2) \leq E_g(C)$, with equality for $C = A$, for every smooth completely monotone g. We recompiled it with the pinned toolchain (Lean 4.34.1, Mathlib `d13f23b7`, openai/math `adc7f124`, unchanged at `fd4aeeb2`); its footprint is {`propext`, `Classical.choice`, `Quot.sound`}. The proof is checked. What is not established by any gate is that these definitions are the intended ones; our item-by-item comparison with [2, Defs. 1.1–1.3] was done by AI agents and is filed as such.

Earlier corollaries (UO-A-0002–4). Riesz admissibility, finiteness of the lattice Riesz energy for $s > 2$, and optimality of $\rho^{-1/2}A$ at every density with attainment are in [3].

Universal optimality in the sense of CKMRV (UO-A-0005). Define, in upstream's notions,

$$\mathrm{UO}(\rho, C) \;:\Longleftrightarrow\; C \text{ locally finite}, \quad \mathrm{dens}(C) = \rho, \quad \text{and} \\ \forall g \text{ admissible}, \ \forall C' \text{ locally finite of density } \rho: \ E_g(C) \le E_g(C').$$

This transcribes [2, Def. 1.3] with one difference: CKMRV also require the energy of C to exist as a limit, and UO uses only the lim inf, so it is weaker (we do not claim it is strictly weaker; no witness is filed). For a lattice whose sum $\sum_{x \neq 0} g(|x|^2)$ converges absolutely, CKMRV take the energy to be that sum (UO-L-0005). Given the unformalised fact that the disk averages then converge to it, Theorem 1 yields CKMRV's notion for $\rho^{-1/2}A$. That step is Tier C and is not claimed here.

Theorem 1 (`triangular_universally_optimal`). *For every $\rho > 0$,* $\mathrm{UO}(\rho, \rho^{-1/2}A)$.

Non-uniqueness (UO-A-0006).

Theorem 2 (`vacancy_energy`; `vacancy_not_isometric_image`; `universal_optimum_not_unique`). *For every admissible g, $E_g(A \setminus \{0\}) = \sum_{a \in A \setminus 0} g(|a|^2)$. Hence* $\mathrm{UO}(1, A \setminus \{0\})$*, and there is no isometry f of the plane with $f(A) = A \setminus \{0\}$.*

Proof idea. Removing one point removes a sub-sum from each disk energy, and the normalisation changes by $N_R/(N_R - 1) \to 1$; the lim inf comparison closes with `ENNReal.liminf_mul_le`, and upstream's theorem gives the reverse inequality. For the isometry statement, isometries of the plane are affine (Mathlib's `Isometry.affineIsometryOfStrictConvexSpace`), so $f(2a-b) = 2f(a) - f(b)$. Write p for `triangularPoint` and pick $a, b \in A$ with $f(a) = p(1,0)$ and $f(b) = p(2,0)$. Then $2a - b \in A$ and $f(2a - b) = 2p(1,0) - p(2,0) = p(0,0) = 0$, contradicting $0 \notin A \setminus \{0\}$ (Figure 2).




Figure 2: The non-uniqueness theorem (UO-A-0006). Removing the origin leaves the lower energy unchanged for every admissible g. No isometry carries A onto $A \setminus \{0\}$: an isometry of the plane is affine, so it would send $2a - b$ to $2p(1,0) - p(2,0) = 0$.

This is the remark after [2, Thm. 1.4] (UO-L-0002), also noted by upstream ("it identifies the minimum value, not all minimizers" [1, §1], UO-L-0006), now as a theorem in this setting. Its interpretation (UO-C-0002) is Tier C: in the lower-energy formulation over *all* locally finite

configurations, "the minimiser is the triangular lattice" is false, and only the minimum value is pinned down. Uniqueness among *periodic* configurations, which CKMRV prove in dimensions 8 and 24, is not claimed by upstream for dimension 2 and is not addressed here.

Gates. A clean sequential rebuild of the whole chain exits 0 (recorded in the AutoevolveAI commit message `d50dfc4`; no build log is committed). The audit file prints 24 footprints: the 22 audited declarations are within the allowlist, and the 2 planted controls (`sorry`, a custom axiom) are flagged by the same audit. Each layer was re-verified by a separate instance of the same AI model that recompiled it and wrote its own controls. That is producer≠verifier, not human review.

3 Tier L: what is known

Per CKMRV's own text (UO-L-0003): Ventevogel and Nijboer proved universal optimality of $\mathbb{Z}$ in dimension 1 (Cohn and Kumar gave another proof); before CKMRV no universal optimum was proved in any dimension > 1; CKMRV proved it for E_8 and the Leech lattice; and universal optimality of the hexagonal lattice "in full generality is an important open problem". So, per CKMRV, dimension 1 was settled before dimensions 8 and 24, and dimension 2 was then the lowest open case.

4 Tier C: why the Yukawa interaction is in the class

$V(r) = e^{-\kappa r}/r$ is $g(t) = e^{-\kappa\sqrt{t}}/\sqrt{t}$ with $t = r^2$. Here $t^{-1/2}$ is completely monotone; $e^{-\kappa\sqrt{t}} = p \circ q$ with $p(u) = e^{-\kappa u}$ completely monotone and $q'(t) = \frac{1}{2}t^{-1/2}$ completely monotone, hence completely monotone by [2, footnote 2] (UO-L-0004); and products of completely monotone functions are completely monotone (Leibniz). So upstream's theorem, if its definitions are faithful, predicts that no density-one configuration has Yukawa energy below the triangular lattice's. This argument is not kernel-checked (UO-C-0001), and so the numerics below rest on Tier C, not on Tier A.

5 Tier X: a CVODE N-body consistency check

Instrument. rusty-SUNDIALS [5], crate `qf-yukawa-crystal`. $N = 288$ particles in a periodic rectangle that holds the triangular lattice 16×18 exactly at density 1; shifted-force Yukawa with cutoff $r_c = 8$ (minimum image), $\kappa = 2$ ($V(r_c) = 1.4 \times 10^{-8}$) and $\kappa = 1$ ($V(r_c) = 4.2 \times 10^{-5}$). The truncated potential is not exactly completely monotone beyond r_c; that is a disclosed approximation. Overdamped relaxation $\dot{x} = -\nabla E$ is integrated by CVODE (BDF) with the analytic Jacobian $-\nabla^2 E$ until $\max|F| < 10^{-9}$, and minima are checked by the Hessian spectrum (cyclic Jacobi). Two packages named in common descriptions of this pipeline were not used: rusty-SUNDIALS has no KINSOL port, and its IDA crate documents itself as a fixed-step prototype. Annealing is overdamped Langevin (Euler–Maruyama) outside SUNDIALS, because a stochastic equation is not an ODE, followed by a CVODE quench.

Initial conditions. Twenty disordered configurations derived from the Hugging Face dataset `colabfit/OC20_IS2RES_val_id` [13] (CC-BY-4.0): random OC20 structures, rotated uniformly in 3D, projected, rescaled and accepted with a minimum separation. Twenty uniform random configurations serve as a second family. We do not claim that either is more realistic; the dataset supplies disorder, not physics. Machine-learned interatomic potentials were deliberately not used: they model atomic chemistry, not a completely monotone pair potential, and cannot test this theorem.

Preregistration. The predictions and controls were committed (`636c514`) before the solver code (`968b0f2`) and before the results (`1f46e70`) were committed. Git cannot show that no local run preceded the preregistration commit. P1: no relaxed configuration has energy per particle below e_{tri} by more than 10^{-9} relative. P2: at least one annealed run per family reaches the perfect lattice. P3: $T = 0$ quenches mostly end in defected local minima, which is a property of gradient flow and not evidence against the theorem. *Deviations from the preregistration.* (a) The coordination cutoff is $1.35\, d_0$, with d_0 the median nearest-neighbour distance of the configuration, not the registered $1.4a$. (b) At $\kappa = 1$ ten initial conditions per family were run, and five incommensurate controls, instead of the registered twenty per family; the number of incommensurate runs was not registered. (c) The incommensurate control's automatic pass test checks only that every state is above e_{tri}; that the states are defected, also registered, is read from the table (all are).




Figure 3: Reference energies per particle at density 1 (the vertical axes do not start at 0). Left and middle: Yukawa, triangular below square at both κ. Right: the jellium extension (Section 6). This code's Ewald energies match the ten-digit Madelung values of Alves et al. [7], which agree with Bonsall and Maradudin [6], to 2.6×10^{-10} (triangular) and 4.3×10^{-10} (square) relative. The square lattice is a saddle point.

Controls. References: $e_{\mathrm{tri}} = 0.3935333668$ and $e_{\mathrm{sq}} = 0.4009647699$ at $\kappa = 2$ ($e_{\mathrm{sq}} - e_{\mathrm{tri}} = 7.43 \times 10^{-3}$); the periodic box reproduces the direct lattice sum to 2.1×10^{-14}. A perfect lattice stays put (relative gap 5.2×10^{-14}); a lattice with $0.1a$ noise relaxes back (gap 2.5×10^{-14}); the Hessian at the lattice has exactly two zero translation modes and smallest other eigenvalue 0.0236. Negative controls: the square lattice is a stationary point with 72 negative Hessian eigenvalues (a saddle) and relaxes below its energy; in a box that cannot hold the triangular lattice the relaxed states keep defects and stay above e_{tri} (pass).

Results. Table 2. **P1** (the falsifiable prediction): pass at $\kappa = 2$, with 0 undercuts in 100 relaxations (20 of them in the incommensurate control box), and pass at $\kappa = 1$, with 0 undercuts in 50 (10 control). The lowest annealed gaps, 7.6×10^{-14} (OC20-derived) and 8.5×10^{-14} (uniform) at $\kappa = 2$, are the lattice itself to round-off. **P2**: pass at $\kappa = 2$, but narrowly: 1 of 20 (OC20-derived) and 1 of 20 (uniform) annealed runs reached the perfect lattice. At $\kappa = 1$ P2 *fails as preregistered*: 0 of 10 and 2 of 10. The other annealed runs end in defected local minima or in defect-free states above e_{tri}. The annealing schedule is a weak instrument, and this failure says nothing about the theorem in either direction. **P3**, as expected: 15 of 20 and 14 of 20 $T = 0$ quenches at $\kappa = 2$ keep defects. The remainder are defect-free but above e_{tri}; their structure was not analysed. Every run converged to $\max |F| < 10^{-9}$. The final Hessian was computed for the 80 ($\kappa = 2$) and 40 ($\kappa = 1$) non-control

runs. 0 and 0 of them have an eigenvalue below $-10^{-9}\lambda_{\max}$. All have exactly two translation modes ($|\lambda| \leq 1.5 \times 10^{-15}$ at $\kappa = 2$, $\leq 1.8 \times 10^{-15}$ at $\kappa = 1$), and every other eigenvalue is at least 5.4×10^{-5} ($\kappa = 2$) and 5.0×10^{-5} ($\kappa = 1$). These end states are therefore strict local minima modulo translation. The incommensurate-box and disk runs were not Hessian-checked. The negative control behaves as it must: in the incommensurate box every relaxed state is defected and at least 9.6×10^{-4} above e_{tri}. At $\kappa = 2$ the two initial-condition families behave alike. At $\kappa = 1$ they differ (Table 2), with small counts. Nothing in these runs depends on the dataset's chemistry: it supplied disorder only.

	family / stage	runs	perfect	defected	min rel. gap	max rel. gap	not conv.	Hess.< 0
$\kappa = 2$	incomm. box (ctl), anneal + quench	10	0	10	9.60×10^{-4}	1.73×10^{-3}	0	n/a
$\kappa = 2$	incomm. box (ctl), $T = 0$ quench	10	0	10	9.60×10^{-4}	5.67×10^{-3}	0	n/a
$\kappa = 2$	OC20-derived, anneal + quench	20	1	11	7.60×10^{-14}	2.53×10^{-3}	0	0/20
$\kappa = 2$	OC20-derived, $T = 0$ quench	20	0	15	1.84×10^{-4}	4.89×10^{-3}	0	0/20
$\kappa = 2$	uniform, anneal + quench	20	1	11	8.48×10^{-14}	2.39×10^{-3}	0	0/20
$\kappa = 2$	uniform, $T = 0$ quench	20	0	14	1.84×10^{-4}	6.04×10^{-3}	0	0/20
$\kappa = 1$	incomm. box (ctl), anneal + quench	5	0	5	3.09×10^{-4}	6.25×10^{-4}	0	n/a
$\kappa = 1$	incomm. box (ctl), $T = 0$ quench	5	0	5	3.10×10^{-4}	1.07×10^{-3}	0	n/a
$\kappa = 1$	OC20-derived, anneal + quench	10	0	5	5.77×10^{-5}	1.22×10^{-3}	0	0/10
$\kappa = 1$	OC20-derived, $T = 0$ quench	10	0	8	5.77×10^{-5}	1.70×10^{-3}	0	0/10
$\kappa = 1$	uniform, anneal + quench	10	2	3	1.93×10^{-14}	8.51×10^{-4}	0	0/10
$\kappa = 1$	uniform, $T = 0$ quench	10	1	8	3.96×10^{-14}	1.22×10^{-3}	0	0/10

Table 2: CVODE relaxations. "perfect": zero defects and $|e - e_{\text{tri}}| < 10^{-8}|e_{\text{tri}}|$. Relative gap $(e - e_{\text{tri}})/e_{\text{tri}}$. "Hess.$< 0$": runs whose final Hessian has an eigenvalue below $-10^{-9}\lambda_{\max}$, over runs whose Hessian was computed (n/a: not computed). The incommensurate family uses a square box and is a negative control. Tier X throughout.




Figure 4: Every relaxation's relative energy gap to the triangular lattice, one marker per run, on a logarithmic axis. P1 concerns the sign of the gap, which the summary script checks for every run (no gap is negative). Markers at the far left are runs that reached the lattice to round-off. No value is clipped: the smallest gaps are 7.6×10^{-14} at $\kappa = 2$ and 1.9×10^{-14} at $\kappa = 1$. Generated from `main_kappa{2,1}.jsonl`.

Open versus closed disk. $N = 300$, $\kappa = 2$, annealed and quenched: harmonic bowl (open) versus soft wall (closed). Interior defect fraction 0.086 (open) versus 0.033 (closed); edge 0.965 versus 0.602; interior mean $|\psi_6|$ 0.913 versus 0.931. The comparison is confounded: the bowl produces a non-uniform density, the closed disk compresses particles against the wall, and the two "edge" layers (within two spacings of the outermost particle) contain different numbers of particles. It describes finite-size edge effects in these two set-ups and is not a controlled measurement of either. It is not a test of whether upstream's closed-disk counting differs from an open-disk one, which is a question about measure-zero boundary terms in a limit. Figure 5 shows one run of each.




Figure 5: Open (harmonic bowl) versus closed (soft wall) disk, $N = 300$: Voronoi cells coloured by coordination. Defects concentrate at the edge in both cases, with a near-triangular interior. One run each (`disk_00`, regenerated from its published seed); the averages are in the text. The counts in the panel titles are recomputed from the snapshot, with d_0 from its own median, and can differ by one from `disk.jsonl`. Tier X, and confounded as stated.

6 Extension: long-range interactions and jellium (Tier X, in progress)

For $0 < s \leq 2$ and for the logarithm, the raw lattice sum diverges. The comparison of Section 2 then reads $E(C) = \infty$ for every competitor and says nothing. Upstream's manuscript treats this range through renormalized energies against a neutralizing background (its Section 8, after Petrache and Serfaty [8]): for $0 \leq s < 2$ the triangular lattice minimizes the renormalized energy, and the jellium minimum equals it up to normalisation [1]. That corollary is *not* in upstream's Lean. As a first, numerical step we treat $s = 1$, the classical 2D electron gas (Wigner crystal), by Ewald summation in the same periodic box, with CVODE relaxation and a preregistration (rusty-SUNDIALS `e706f85`) committed before the code (`3bc55fa`; snapshot command `fab1e6e`; branch `physics/jellium-wigner-2026-10-11`). Its controls pass:

- the triangular and square Madelung energies equal the ten-digit values of Alves et al. [7], which agree with Bonsall and Maradudin [6], to 2.6×10^{-10} (triangular) and 4.3×10^{-10}

(square) relative (Figure 3). That is the rounding level of the nine-decimal published values; the preregistered tolerance was 10^{-8};

- the energy changes by less than 10^{-10} relative over a $4\times$ range of the Ewald parameter;
- the square lattice is a saddle point with 64 unstable modes, consistent with the imaginary phonon branch reported in [6].

The relaxation study (prediction J1: no relaxed configuration below the triangular Madelung energy) is reported in the companion record when it completes. The $s = 2$ endpoint is outside upstream's statement. The formal targets for this range (a heat-trace comparison, and the Laplace-transform step of upstream's Lemma "Heat comparison across period lattices") are listed with priors in `docs/EXTENSION_RENORMALIZED_JELLIUM_2026-10-11.md`.

7 Laboratory protocols (Tier C proposals)

The theorem is about exact potentials and infinite configurations. A laboratory has finite temperature, finite N, polydispersity and imperfect interaction laws. Experiments therefore cannot test the Lean statement. What they can do is test whether real two-dimensional repulsive systems behave as the theorem and our numerics suggest. Each protocol below is a proposal (Tier C); none has been run. All four share one analysis script, `lab/lab_analysis.py`, which takes tracked particle coordinates and computes the same observables as the simulations.

- *Density* comes from the mean Voronoi cell area of the same interior particles over which the energy is averaged.
- *Coordination* uses the $1.35\,d_0$ rule; the script also gives the local and global $|\psi_6|$.
- *Energy:* the interaction energy per interior particle for a stated law, truncated at r_c with a uniform tail correction, compared with a triangular lattice at the same density and truncation. A Yukawa κ is supplied in the experiment's units and converted internally.

Its self-test passes. A perfect lattice gives gap 6×10^{-14} at an arbitrary length scale; thermal noise raises the energy above the lattice; a square lattice is detected and placed above it. Two caveats bear on the kill criteria. The truncated potential is not exactly completely monotone, so the comparison is between equally truncated energies. And at $s = 3$ the energy scales as $\rho^{3/2}$, so a density error $\delta\rho/\rho$ propagates to about $1.5\,\delta\rho/\rho$ in energy and must be carried into the error bar. Every protocol should be preregistered with that script, a predicted bracket, and the kill criterion below.

L1. Superparamagnetic colloids at a flat air–water interface (Riesz $s = 3$). Following Zahn, Wille, Maret, Sengupta and Nielaba [9], a perpendicular field B induces parallel dipoles. The pair energy is $\propto \chi^2 B^2/r^3$, a Riesz potential with $s = 3$: in the theorem's class, with finite lattice energy (D1b). It realises an interaction in D1's class, up to finite-particle-size and mutual-induction corrections to the point-dipole law. *Procedure.* Ramp B slowly from the fluid to the crystal (the laboratory analogue of annealing) at fixed area fraction. Track positions by video microscopy, and record χ and B so that the law's prefactor is known. *Predictions.*

- (a) At large coupling, $|\psi_6| \to 1$ and the defect fraction $\to 0$ in the interior.
- (b) The measured dipolar energy per interior particle exceeds the triangular value at the measured density, by an amount that shrinks as B grows (the thermal excess).

- (c) A particle removed with optical tweezers leaves a vacancy that relaxes locally, without global restructuring.
- (d) A square arrangement imposed with holographic tweezers and then released relaxes away from square. This is the analogue of the square-saddle control computed here for Yukawa and jellium. The $s = 3$ square lattice was not computed.

Kill criterion. A slowly ramped, defect-free interior whose dipolar energy lies below the triangular value beyond the measurement error, with the law's prefactor independently confirmed.

L2. Dusty-plasma monolayer (Yukawa). Thomas et al. [10] observed Coulomb crystallization of charged microparticles levitated in a plasma sheath. The interaction is screened Coulomb (Yukawa), and the radial confinement is close to parabolic, the analogue of our open bowl. *Procedure.* Measure κ and the coupling from the phonon spectrum or the pair correlation. Image the monolayer, run the shared script, and simulate our bowl model at the measured κ and N. *Predictions.* A near-triangular interior, with defects concentrated at the rim. The interior defect fraction is compared with the simulation at matched κ and N (Figure 5, left). *Confounds.* Ion-wake attraction, out-of-plane motion and departures from a pure Yukawa law at large distance move the system outside the theorem's class. The wake attraction is non-reciprocal, acts chiefly between vertically adjacent grains, and can drive a mode-coupling instability in a monolayer. Neutral-gas drag damps the dynamics but does not change the interaction law.

L3. Charged colloids in circular, hexagonal and square cavities (closed disk). Bubeck, Neser, Bechinger and Leiderer [11] studied small two-dimensional colloidal crystals confined in hexagonal, circular and square regions. Per the publisher's abstract (we did not read the full text), these had about 10 to 100 particles, and their structure was dominated by the shape of the region. *Procedure.* Repeat the circular-cavity case at $N \approx 300$ and image the interior; compare with our closed-wall runs at matched N and wall stiffness. *Prediction.* The edge defect fraction exceeds the interior one, as in Figure 5, right. Compared with L2 at matched N and density, this is the experimental open-versus-closed comparison.

L4. Electrons on liquid helium (jellium, $s = 1$). Grimes and Adams [12] observed the liquid-to-crystal transition of a classical 2D sheet of electrons held above liquid helium. The neutralizing charge sits on electrodes a distance d below the sheet, not in its plane. This is therefore the jellium case of Section 6 only when d is much larger than the interparticle spacing; otherwise image charges make the interaction dipolar at distances large compared with d. *Prediction.* The crystal is triangular. The evidence in [12] is spectroscopic, not a direct image, so a test here compares spectroscopic signatures with the triangular Madelung and phonon structure [6]. Direct imaging of the lattice would need a different detection method, which we do not specify.

8 Limitations

1. No human has audited whether upstream's definitions, or ours, express the intended mathematics; the ledger flags every Tier A row for this.
2. UO is weaker than CKMRV's universal optimality (it does not require the energy limit to exist).

3. Uniqueness among periodic configurations in dimension 2 is not claimed by upstream and was not attempted.

4. The Yukawa step is Tier C; the numerics are Tier X and support no claim.

5. Finite N, truncated potential, heuristic annealing: P2 can fail for reasons that have nothing to do with the theorem.

Reproducibility. Lean: AutoevolveAI `formal_cert/Triangular/`; LeanMaster release `v3.48.1`. Numerics: rusty-SUNDIALS results commit `1f46e70`; summary regenerated by `scripts/summarize.py` at `e1a6083`. Ledger and evidence: `results/discovery/ledger/` (gate: `tools/ledger.py` of [4], exit 1: 22 claims, 0 blocks, 6 flags).

Version history. v1 (doi:10.5281/zenodo.23293112): the text, tables and ledger. v2 adds the figures, all generated by `make_figures.py` from the result files and from snapshots that re-run published seeds. It also adds the jellium extension's preregistration and controls (Section 6) and the laboratory protocols (Section 7). No v1 result changed. Claims added in v2 have their own ledger rows (UO-L-0007, UO-X-0005, UO-X-0006, UO-C-0003); the ledger counts above include them.

Publication decision. The Elenchus publish gate requires the ledger gate to exit 0, and it exits 1 because of the 6 unaudited-statement flags. The author reviewed the draft and decided to publish with those flags open, as a preprint, and to seek the statement-adequacy review from the community. Until that review exists, every Tier A row should be read as "the kernel checked the proof; no person has certified that the statement means what its gloss says".

AI use. The proofs, numerics, ledger and text were produced by AI agents (Claude) under the author's direction; verification runs and the manuscript audit were separate instances of the same model. A DeepSeek-Prover-V2 model (on Cloud Run) was also queried during the Lean work; none of its output was used. The author is responsible for the content.

References

[1] OpenAI, Universal optimality of the triangular lattice, OpenAI Math Release preprint, 2026-09-23, `https://github.com/openai/math`.

[2] H. Cohn, A. Kumar, S. D. Miller, D. Radchenko, M. Viazovska, Universal optimality of the E_8 and Leech lattices and interpolation formulas, Ann. of Math. 196 (2022) 983–1082, doi:10.4007/annals.2022.196.3.3.

[3] X. Callens, Riesz-energy, finiteness and every-density corollaries of the openai/math planar universal-optimality theorem, kernel-checked in Lean 4, Zenodo (2026), doi:10.5281/zenodo.23285631.

[4] X. Callens, SocrateAI-Scientific-Elenchus, `https://github.com/xaviercallens/SocrateAI-Scientific-Elenchus` (commit `c9ae8cd`).

[5] X. Callens, rusty-SUNDIALS, `https://github.com/xaviercallens/rusty-SUNDIALS`.

[6] L. Bonsall, A. A. Maradudin, Some static and dynamical properties of a two-dimensional Wigner crystal, Phys. Rev. B 15 (1977) 1959, doi:10.1103/PhysRevB.15.1959.

[7] E. Alves, G. L. Bendazzoli, S. Evangelisti, J. A. Berger, Accurate ground-state energies of Wigner crystals from a simple real-space approach, Phys. Rev. B 103 (2021) 245125, doi:10.1103/PhysRevB.103.245125.

[8] M. Petrache, S. Serfaty, Crystallization for Coulomb and Riesz interactions as a consequence of the Cohn–Kumar conjecture, Proc. AMS 148 (2020) 3047–3057, doi:10.1090/proc/15003.

[9] K. Zahn, A. Wille, G. Maret, S. Sengupta, P. Nielaba, Elastic properties of 2D colloidal crystals from video microscopy, Phys. Rev. Lett. 90 (2003) 155506, doi:10.1103/PhysRevLett.90.155506.

[10] H. Thomas, G. E. Morfill, V. Demmel, J. Goree, B. Feuerbacher, D. Möhlmann, Plasma crystal: Coulomb crystallization in a dusty plasma, Phys. Rev. Lett. 73 (1994) 652, doi:10.1103/PhysRevLett.73.652.

[11] R. Bubeck, S. Neser, C. Bechinger, P. Leiderer, Structure and dynamics of two-dimensional colloidal crystals in confined geometry, Prog. Colloid Polym. Sci. (1998) 41–45, doi:10.1007/BFb0118046.

[12] C. C. Grimes, G. Adams, Evidence for a liquid-to-crystal phase transition in a classical, two-dimensional sheet of electrons, Phys. Rev. Lett. 42 (1979) 795, doi:10.1103/PhysRevLett.42.795.

[13] L. Chanussot et al., OC20 IS2RES val id, ColabFit (2024), doi:10.60732/b4005525; Open Catalyst 2020, ACS Catal. 11 (2021) 6059, doi:10.1021/acscatal.0c04525.