

Thermodynamic Censorship of Navier-Stokes Singularities: Why Mathematical Blow-ups Are Physically Unrealizable

Xavier Callens

Socrate AI Lab, MechanicaFluidorum Program

September 12, 2026

WITHDRAWN — superseded, retained for the historical record only. Do not cite for its title, abstract or conclusions. This September 12 draft argued that the OpenAI blow-up is “physically unrealizable” through a “thermodynamic censorship” principle; that framing is withdrawn. The OpenAI proofs are correct theorems about a mathematical model, and this project’s position is a physical *reading* of them, not a refutation. Cite instead `01_Verification_Paper/OpenAI_NSE_Verification.tex` (“A Physical Reading, Not a Physical Refutation”, v5.5.0, DOI 10.5281/zenodo.22806767; concept DOI 10.5281/zenodo.22696717). The intermediate “Version 2” records (10.5281/zenodo.22725347, 22727801) also carry withdrawn claims. Nothing in this project shows that something physical stops the construction’s collapse: on its route the incompressible description ends at $\ell_* = \nu/c$ (paper §§5, 9). The specific corrections made there, and applied inline below where a sentence would otherwise be false, are:

- “*Plasma temperatures.*” Because the core keeps $\ell_r^2 = \nu t$, viscous heating over the remaining time is $\Delta T = \varepsilon t/c_p = u^2/c_p$: about 50 K at $\text{Ma} = 0.3$ and 540 K at $\text{Ma} = 1$ in water. Boiling is reached near $\text{Ma} \approx 0.37$; 10^4 K would need $\text{Ma} \approx 4$, far outside the model. No plasma is implied.
- “*The constant $\Omega_{\max} = 1.13 \times 10^{13} \text{ s}^{-2}$ has no derivation.*” A global integrated-enstrophy bound is also the wrong object (it scales with the fluid volume). The physically motivated admissibility condition is local: $|\omega| \lesssim c^2/\nu$ ($2.2 \times 10^{12} \text{ s}^{-1}$ in water), equivalent to $\text{Ma} \lesssim 1$ and $\text{Kn} \lesssim 1$ together; with it, “admissible on $[0, T] \Rightarrow$ regular on $[0, T]$ ” is the Beale–Kato–Majda theorem, not an axiom (§6).
- “*The physics is vacuous*” / “*censorship.*” Withdrawn. The result is a statement about the continuum model; the physical content is a map of *where* the model’s constitutive assumptions end, namely at the single scale $\ell_* = \nu/c$ (water: 0.7 nm).
- “*Lemma 8.7.*” The five-moment matching is Proposition B.8 / Eqs. (5.10)–(5.11) of the OpenAI paper; Lemma 8.7 is a different lemma.
- “*Timeline.*” With $\ell_r \simeq \sqrt{\nu t}$ and $u \simeq \sqrt{\nu/t}$, the breakdown times are essentially independent of the reference length ℓ_0 (which enters only as $(\ell_0^2/\nu t)^h \approx 1.3$); the “6.7 ps” figure is 4.9 ps in unit-free form.

Abstract

In September 2026, an OpenAI multi-agent system produced a Lean 4 formalized proof of finite-time blow-up for the forced 3D incompressible Navier-Stokes equations, establishing Alternatives C and D of the Clay Millennium Prize Problem. We independently verify the logical soundness of this formalization: zero **sorry** axioms, zero custom axioms, correct C^∞

force type, correct uniform L^2 energy bound. The Lean 4 kernel accepts the proof, and the mathematics is irrefutable.

We then ask how far the constructed singularity travels outside the model’s domain of validity. Three lines of evidence: (i) intensive local kinetic energy density diverges as $\tau^{-1.010}$ and integrated enstrophy as $\tau^{-0.515}$; the corresponding viscous heating is $\Delta T = u^2/c_p$, which passes the boiling point of water near $\text{Ma} \approx 0.37$ (not, as an earlier version of this abstract said, a plasma threshold); (ii) the Mach number exceeds 0.3 at physical time $t = T\tau \sim 6.7 \times 10^{-12}$ seconds before singularity (≈ 6.7 picoseconds for $\ell_0 = 1$ cm, $T = \ell_0^2/\nu = 100$ s; 4.9 ps in the unit-free form $t \simeq \nu/(\text{Ma } c)^2$), at which point NSE cease to describe the physical system; (iii) non-dimensionalization analysis reveals the moment-matching system is well-conditioned ($\kappa(B) \approx 4.11 \times 10^5$), confirming the raw 10^{28} condition number in prior literature was an unscaled numerical artifact, while the velocity profile itself relies on non-analytic Gevrey cutoffs ($s = 1.5$).

We state a physical admissibility condition in local form, $|\omega| \lesssim c^2/\nu$, and note that under it the absence of blow-up is a consequence of the Beale–Kato–Majda criterion rather than a postulate.

1 Introduction

The three-dimensional incompressible Navier-Stokes equations (NSE) have stood as a cornerstone of fluid mechanics for nearly two centuries [1]. In 2000, Charles Fefferman formalized the Millennium Prize problem, asking whether solutions to the unforced or forced NSE either exist globally in time and remain smooth, or whether they can develop finite-time singularities (blow-ups) [2].

In September 2026, an OpenAI multi-agent system claimed a formalized proof in Lean 4 establishing a finite-time blow-up for the forced 3D incompressible NSE. This result formally resolves Alternatives C and D of the Millennium Prize problem. Our Physical Verification separates the mathematical validity of the proof from its physical significance. While the mathematics is robust, our analysis shows that the constructed blow-up leaves the regime in which the incompressible continuum model describes a fluid, and it identifies the scale at which this happens. This prompts a distinction between the mathematical and physical validity of NSE solutions. We state a physical admissibility condition (§6) and locate where the AI-generated singularity exits it prior to the blow-up time.

The paper is structured as follows. Section 2 details the formal verification of the OpenAI Lean 4 proof. Section 3 analyzes the paradoxical energy density divergence. Section 4 explores the breakdown of the incompressibility assumption via Mach number self-invalidation. Section 5 discusses the non-dimensionalization of the moment-matching system. Section 6 states the physical admissibility condition. Section 7 discusses the broader implications, and Section 8 concludes.

2 Formal Verification of the OpenAI Lean 4 Proof

2.1 Methodology: AST Crawling and Type Verification

We performed a rigorous Physical Verification of the provided Lean 4 repository. Using Abstract Syntax Tree (AST) crawling, we verified that the final theorem statements strictly adhere to the Millennium Prize definitions.

2.2 Results: Zero Axiomatic Leakage

The formalization is pristine. We confirm:

- Zero sorry keywords in the critical path.

- Zero custom axioms bypassing standard real analysis or Sobolev spaces.
- The external force field $f(x, t)$ is rigorously shown to be in $C^\infty(\mathbb{R}^3 \times [0, \infty))$.
- The total kinetic energy satisfies $\sup_t \int |u(x, t)|^2 dx \leq E < \infty$.

2.3 The Gevrey Bypass

A key innovation in the AI's approach is the use of Gevrey-class functions, specifically of the form $\exp(-1/q^2)$. Gevrey functions are C^∞ but not real-analytic. Their Taylor series diverge, which allows the AI to construct compactly supported localized structures that undergo self-similar collapse. Exact theoretical analysis proves that $\exp(-1/q^2)$ has an asymptotic Gevrey index of $s = 1.5$ (symbolic fits on small $N \leq 15$ yield a pre-asymptotic slope of ≈ 2.22).

2.4 The Method of Manufactured Solutions

The AI constructed the blow-up by specifying a collapsing vortex solution and then calculating the necessary forcing function to satisfy the NSE. In `CandidateFromLimits.lean`, the identity `force_eq_activated_residual` establishes $f(x, t)$ as exactly the residual of the NSE evaluated on the Gevrey candidate. Since the candidate is C^∞ , the force is C^∞ .

3 Thermodynamic Paradox: Energy Density Divergence

Despite the bounded global kinetic energy, the local structure of the singularity exhibits extreme behavior as time t approaches the blow-up time T . Let $\tau = T - t$.

3.1 Scaling Analysis

The AI's collapsing vortex scales radially as $l_r \sim \tau^{1/2}$ and azimuthally as $u_\theta \sim \tau^{-1/2-h}$, where $h = 1/200 > 0$ is an anomalous exponent.

3.2 Energy and Enstrophy Divergence

The global kinetic energy remains bounded because the support volume shrinks as $V \sim \tau^{3/2-h}$. Thus, $E(\tau) \sim V u^2 \sim \tau^{3/2-h} \tau^{-1-2h} = \tau^{1/2-3h} = \tau^{+0.485} \rightarrow 0$, satisfying the finite-energy condition.

However, the intensive local energy density scales purely as $e_{\text{local}} = \frac{1}{2}\rho|u|^2 \sim \tau^{-1-2h} = \tau^{-1.010} \rightarrow \infty$. The enstrophy $\Omega = \int |\nabla \times u|^2 dx$ diverges as $\tau^{-0.515} \rightarrow \infty$. Higher norms such as the L^3 norm $\|u\|_3 \sim \tau^{-0.0067}$ and the Sobolev norm $\|u\|_{H^{3/2}} \sim \tau^{-0.5075}$ also diverge.

Table 1: Scaling exponents of critical quantities as $\tau \rightarrow 0$ ($h = 1/200$)

Quantity	Scaling $\propto \tau^\alpha$
Length scale l_r	+0.5000
Velocity $ u $	-0.5050
Vorticity $ \omega $	-1.0050
Global Energy E	+0.4850
Intensive Energy Density e_{local}	-1.0100
Enstrophy Ω	-0.5150
L^3 Norm $\ u\ _3$	-0.0067
$H^{3/2}$ Sobolev Norm $\ u\ _{H^{3/2}}$	-0.5075

3.3 Viscous Heating and the Isothermal Assumption

The local dissipation generated by the diverging vorticity conflicts with the isothermal assumption of the NSE. Because the core keeps a radial Reynolds number of order one, $\ell_r^2 = \nu t$ where t is the physical time remaining, so the dissipation per unit mass $\varepsilon = \nu(u/\ell_r)^2$ acting over that time gives

$$\Delta T = \frac{\varepsilon t}{c_p} = \frac{u^2}{c_p} \sim \tau^{-1.010}, \quad (1)$$

i.e. an Eckert number of order one. For water this is ≈ 48 K at $Ma = 0.3$ and ≈ 540 K at $Ma = 1$; boiling ($\Delta T = 73$ K) is reached at $u \approx 553$ m/s, $Ma \approx 0.37$, about 3–4.5 ps before the singularity. The formal divergence $\Delta T \rightarrow \infty$ is a property of the model continued past its validity; physically it says only that heating becomes an order-one effect at the same scale as compressibility. It does *not* imply plasma temperatures: 10^4 K would require $Ma \approx 4$, where the incompressible description has long ceased to apply. The estimate assumes the heat stays in the core over the collapse time, which holds when the Prandtl number is of order one or larger (water, $Pr \approx 7$); for gases ($Pr \approx 0.7$) it is an upper estimate.

4 Mach Number Self-Invalidation

The incompressible NSE assume the Mach number $Ma = |u|/c_s \ll 0.3$.

4.1 Physical Reference Scales

Consider water at 300K: speed of sound $c_s = 1500$ m/s, kinematic viscosity $\nu = 10^{-6}$ m²/s.

4.2 Mach Number Trajectory

The peak velocity diverges as $|u|_{max} \sim \tau^{-0.5050}$.

The self-similar collapse profile is parameterized by a dimensionless time-to-singularity τ ; the physical elapsed time is $t = T\tau$ with $T \equiv \ell_0^2/\nu = 100$ s for the water example ($\ell_0 = 1$ cm), *not* τ read directly in seconds. The table below reports t .

Table 2: Physical invalidation timeline prior to singularity (time column corrected to physical seconds $t = T\tau$, $T = 100$ s)

Time to singularity $t = T\tau$ (s)	Mach Number Ma	Physical State
1.0×10^{-1}	10^{-10}	Valid incompressible fluid
6.7×10^{-12}	0.3	Compressibility onset (Invalidation)
6.2×10^{-13}	1.0	Supersonic flow
9.0×10^{-14}	~ 10	Continuum breakdown (Knudsen ~ 1)

At $t = 6.7 \times 10^{-12}$ s, Ma exceeds 0.3. The mathematical model ceases to describe the physical fluid at this exact moment.

5 Matrix Scaling & Non-Dimensionalization of the Moment System

To ensure the Gevrey profile satisfies the NSE, the AI enforces a 5-moment matching condition on the radial profile, a 5×5 linear system in the five unknown coefficients (Proposition B.8 and Eqs. (5.10)–(5.11) of the OpenAI paper; an earlier version of this paper cited “Lemma 8.7”, which is a different lemma).

5.1 Non-Dimensionalization Analysis

Let X_R be the truncation radius. A raw SVD of A yields a condition number $\kappa(A) \sim X_R^{7.75}$ reaching 10^{28} at $X_R = 1000$. However, matrix factorization $A = DBD$ with diagonal scaling $D = \text{diag}(1, X_R^2, X_R^4)$ isolates the physical dimensional powers.

Table 3: Condition number of raw versus non-dimensionalized moment matrix

Truncation Radius X_R	Raw $\kappa(A)$	Non-Dimensionalized $\kappa(B)$
10	2.36×10^{12}	4.11×10^5
100	2.36×10^{20}	4.11×10^5
500	7.22×10^{25}	4.11×10^5
1000	1.78×10^{28}	4.11×10^5

5.2 Structural Stability

Preconditioning with D^{-1} yields a non-dimensionalized condition number $\kappa(B) \approx 4.11 \times 10^5$ that is uniform and bounded for all X_R . The raw 10^{28} condition number is a classic numerical artifact of unscaled dimensional matrices. Proper non-dimensionalization confirms the moment-matching system is mathematically well-conditioned.

6 A Physical Admissibility Condition

To bridge the gap between mathematics and physics, we state the condition under which the incompressible continuum model is being used within its domain of validity.

Definition 1 (Local admissibility). A solution of the Navier-Stokes equations is *physically admissible* on $[0, T)$ if its vorticity satisfies

$$\sup_{t < T} \|\omega(\cdot, t)\|_{L^\infty} \lesssim \frac{c^2}{\nu}, \quad (2)$$

where c is the sound speed and ν the kinematic viscosity of the fluid being modelled.

6.1 Physical Motivation

A vortex of velocity U and size L has $|\omega| \sim U/L$. Compressibility requires $U \lesssim c$ and the continuum hypothesis requires $L \gtrsim \lambda \sim \nu/c$ (the mean free path in a gas, the intermolecular spacing in a liquid, both of order ν/c). Hence $|\omega| \lesssim c^2/\nu$ is the vorticity form of “ $\text{Ma} \lesssim 1$ and $\text{Kn} \lesssim 1$ ”, and $\varepsilon = \nu|\omega|^2 \lesssim c^4/\nu$ is the corresponding bound on dissipation per unit mass. For water at 300 K, $c^2/\nu \approx 2.2 \times 10^{12} \text{ s}^{-1}$ (and $c^4/\nu \approx 5 \times 10^{18} \text{ W/kg}$); for air, $c^2/\nu \approx 7.5 \times 10^9 \text{ s}^{-1}$. The condition is local: a bound on the global integrated enstrophy $\int |\omega|^2 dx$, as proposed in an earlier version of this section with an unjustified constant “ $\Omega_{\max} = 1.13 \times 10^{13} \text{ s}^{-2}$ ”, scales with the volume of fluid and does not express any material limit.

6.2 Consequence via Beale–Kato–Majda

If a smooth solution is admissible on $[0, T)$ then $\int_0^T \|\omega(t)\|_{L^\infty} dt \leq Tc^2/\nu < \infty$, and the Beale–Kato–Majda criterion [4], which holds for Navier–Stokes as well as Euler, gives regularity up to and including T . So “admissible on $[0, T)$ implies no blow-up at T ” is a theorem whose only input is the definition above; conversely, any finite-time singularity must first leave the admissible set. The OpenAI core does so when $\ell_r \simeq \sqrt{\nu t}$ reaches $\ell_* = \nu/c$, i.e. at $t \simeq \nu/c^2 \approx 0.4 \text{ ps}$ in water.

6.3 Lean 4 Formalization

The file `ThermodynamicCensorship.lean` in this repository encodes an earlier, global-entropy version of this condition. *Caveat.* Any Lean “rejection” obtained from it is a direct consequence of the definition of admissibility: the type checker verifies that a solution with unbounded vorticity is not admissible, which is what the definition says, and nothing about real fluids. The content is in the physical motivation of the threshold c^2/ν and in the BKM step, neither of which is formalized against OpenAI’s actual objects at the time of writing; the honest formal target is the BKM-based statement above, applied to OpenAI’s `CandidateProperties`.

7 Discussion

The AI’s achievement is a monumental milestone in automated reasoning. It successfully exploited the analytical blindspot between C^∞ and real-analytic functions using Gevrey-class functions.

However, it solved the Clay Institute’s purely mathematical formulation, not the physicist’s problem. This situation strongly parallels Cosmic Censorship in General Relativity [9], where naked singularities are mathematically possible but hypothesized to be shielded by event horizons in physical reality.

The forced nature of the solution (Alternatives C and D) leaves open whether unforced, compactly supported initial data (Alternatives A and B) can blow up. Furthermore, the AI’s success perfectly encapsulates the adage: “the map is not the territory.”

8 Conclusion

The OpenAI Lean 4 proof of Navier-Stokes blow-up is formally correct. Its physical content is a map of where the continuum model ends: the core reaches $\text{Ma} = 0.3$ about 5–7 ps before the singularity, at which point viscous heating ($\Delta T = u^2/c_p \approx 50\text{ K}$) and rarefaction (ℓ_r a few nanometres) are already order-one effects, all at the single scale $\ell_* = \nu/c$. We state a local admissibility condition, $|\omega| \lesssim c^2/\nu$, under which the absence of blow-up follows from the Beale–Kato–Majda criterion.

The AI has not solved the physicist’s problem; it has solved the mathematician’s problem and, in doing so, exposed the gap between them.

A Appendix: Derivation of the Thermal Energy Equation from Enstrophy

To formally answer how enstrophy divergence implies thermal shock without an explicit temperature equation in the standard incompressible NSE, we must explicitly couple the momentum equations with the thermal energy equation. The classical incompressible NSE assume constant density (ρ) and constant temperature (T). To rigorously demonstrate thermal vaporization, we track the temperature field $T(x, t)$ driven by the viscous dissipation function Φ :

$$\rho c_p \left(\frac{\partial T}{\partial t} + u \cdot \nabla T \right) = k \nabla^2 T + 2\mu \mathbf{D}(u) : \mathbf{D}(u) \quad (3)$$

where $\mathbf{D}(u) = \frac{1}{2}(\nabla u + (\nabla u)^T)$ is the rate-of-strain tensor. For a fluid with suitable boundary conditions, the spatial integral of the dissipation tensor contraction equals the enstrophy:

$$\int_{\mathbb{R}^3} 2\mu \mathbf{D}(u) : \mathbf{D}(u) dx = \mu \int_{\mathbb{R}^3} |\nabla \times u|^2 dx = \mu \Omega(t) \quad (4)$$

For the AI’s collapsing vortex, the local enstrophy density scales as $|\omega|^2 \sim \tau^{-2.010}$ (the integrated enstrophy Ω as $\tau^{-0.515}$). The thermal diffusion length over the remaining time, $\sqrt{\alpha t}$, compares with the core radius $\ell_r = \sqrt{\nu t}$ as $\text{Pr}^{-1/2}$: for water ($\text{Pr} \approx 7$) conduction removes only a fraction of the heat on the collapse time, so the source term dominates and $\Delta T \simeq \varepsilon t / c_p = u^2 / c_p \sim \tau^{-1.010}$; for gases ($\text{Pr} \approx 0.7$) the same expression is an upper estimate. This mechanically driven heating passes the boiling threshold of water ($\Delta T = 73 \text{ K}$) at $\text{Ma} \approx 0.37$, a few picoseconds before the singularity, at essentially the same moment as compressibility becomes order one. The earlier claim in this appendix that it reaches “the plasma threshold ($T > 10^4 \text{ K}$)” is withdrawn: that would require $\text{Ma} \approx 4$, outside the model’s validity.

References

- [1] Leray, J. (1934). Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Mathematica*, 63, 193-248.
- [2] Fefferman, C. L. (2000). Existence and smoothness of the Navier-Stokes equation. *Clay Mathematics Institute*.
- [3] OpenAI. (2026). *Formal Resolution of the Navier-Stokes Problem in Lean 4*.
- [4] Beale, J. T., Kato, T., & Majda, A. (1984). Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Communications in Mathematical Physics*, 94(1), 61-66.
- [5] Escauriaza, L., Seregin, G., & Šverák, V. (2003). $L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness. *Russian Mathematical Surveys*, 58(2), 211.
- [6] Constantin, P., & Foias, C. (1988). *Navier-Stokes Equations*. University of Chicago Press.
- [7] Kolmogorov, A. N. (1941). The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Doklady Akademii Nauk SSSR*, 30, 299-303.
- [8] Caffarelli, L., Kohn, R., & Nirenberg, L. (1982). Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Communications on Pure and Applied Mathematics*, 35(6), 771-831.
- [9] Penrose, R. (1969). Gravitational collapse: The role of general relativity. *Rivista del Nuovo Cimento*, 1, 252-276.
- [10] Tao, T. (2016). Finite time blowup for an averaged three-dimensional Navier-Stokes equation. *Journal of the American Mathematical Society*, 29(3), 601-674.