

Thermodynamic Censorship of Navier-Stokes Singularities: Why Mathematical Blow-ups Are Physically Unrealizable

Xavier Callens

Socrate AI Lab, MechanicaFluidorum Program

September 12, 2026

Abstract

In September 2026, an OpenAI multi-agent system produced a Lean 4 formalized proof of finite-time blow-up for the forced 3D incompressible Navier-Stokes equations, establishing Alternatives C and D of the Clay Millennium Prize Problem. We independently verify the logical soundness of this formalization: zero `sorry` axioms, zero custom axioms, correct C^∞ force type, correct uniform L^2 energy bound. The Lean 4 kernel accepts the proof, and the mathematics is irrefutable.

However, we demonstrate through computational analysis that the constructed singularity is thermodynamically unrealizable. Three independent lines of evidence: (i) intensive local kinetic energy density diverges as $\tau^{-1.010}$ and global enstrophy diverges as $\tau^{-0.515}$, generating temperatures past the plasma threshold, violating Boussinesq incompressibility; (ii) the Mach number exceeds 0.3 at $\tau \sim 6.7 \times 10^{-14}$ seconds before singularity (67 femtoseconds), at which point NSE cease to describe the physical system; (iii) non-dimensionalization analysis reveals the moment-matching system is well-conditioned ($\kappa(B) \approx 4.11 \times 10^5$), confirming the raw 10^{28} condition number in prior literature was an unscaled numerical artifact, while the velocity profile itself relies on non-analytic Gevrey cutoffs ($s = 1.5$).

We formalize a Thermodynamic Censorship principle: physically admissible solutions must satisfy uniform bounded enstrophy. Under this axiom, Gevrey-class collapsing-vortex singularities are provably censored.

1 Introduction

The three-dimensional incompressible Navier-Stokes equations (NSE) have stood as a cornerstone of fluid mechanics for nearly two centuries [?]. In 2000, Charles Fefferman formalized the Millennium Prize problem, asking whether solutions to the unforced or forced NSE either exist globally in time and remain smooth, or whether they can develop finite-time singularities (blow-ups) [?].

In September 2026, an OpenAI multi-agent system claimed a formalized proof in Lean 4 establishing a finite-time blow-up for the forced 3D incompressible NSE. This result formally resolves Alternatives C and D of the Millennium Prize problem. Our Physical Verification separates the mathematical validity of the proof from its physical significance. While the mathematics is robust, our analysis reveals that the constructed blow-up requires unphysical thermodynamic and kinematic extremes. This prompts a fundamental distinction between the mathematical and physical validity of NSE solutions. We introduce the concept of Thermodynamic Censorship, demonstrating that the AI-generated singularity violates fundamental physical limits prior to the blow-up time.

The paper is structured as follows. Section 2 details the formal verification of the OpenAI Lean 4 proof. Section 3 analyzes the paradoxical energy density divergence. Section 4 explores the breakdown of the incompressibility assumption via Mach number self-invalidation. Section 5 discusses the non-dimensionalization of the moment-matching system. Section 6 introduces

the Thermodynamic Censorship Principle. Section 7 discusses the broader implications, and Section 8 concludes.

2 Formal Verification of the OpenAI Lean 4 Proof

2.1 Methodology: AST Crawling and Type Verification

We performed a rigorous Physical Verification of the provided Lean 4 repository. Using Abstract Syntax Tree (AST) crawling, we verified that the final theorem statements strictly adhere to the Millennium Prize definitions.

2.2 Results: Zero Axiomatic Leakage

The formalization is pristine. We confirm:

- Zero `sorry` keywords in the critical path.
- Zero custom axioms bypassing standard real analysis or Sobolev spaces.
- The external force field $f(x, t)$ is rigorously shown to be in $C^\infty(\mathbb{R}^3 \times [0, \infty))$.
- The total kinetic energy satisfies $\sup_t \int |u(x, t)|^2 dx \leq E < \infty$.

2.3 The Gevrey Bypass

A key innovation in the AI’s approach is the use of Gevrey-class functions, specifically of the form $\exp(-1/q^2)$. Gevrey functions are C^∞ but not real-analytic. Their Taylor series diverge, which allows the AI to construct compactly supported localized structures that undergo self-similar collapse. Exact theoretical analysis proves that $\exp(-1/q^2)$ has an asymptotic Gevrey index of $s = 1.5$ (symbolic fits on small $N \leq 15$ yield a pre-asymptotic slope of ≈ 2.22).

2.4 The Method of Manufactured Solutions

The AI constructed the blow-up by specifying a collapsing vortex solution and then calculating the necessary forcing function to satisfy the NSE. In `CandidateFromLimits.lean`, the identity `force_eq_activated_residual` establishes $f(x, t)$ as exactly the residual of the NSE evaluated on the Gevrey candidate. Since the candidate is C^∞ , the force is C^∞ .

3 Thermodynamic Paradox: Energy Density Divergence

Despite the bounded global kinetic energy, the local structure of the singularity exhibits extreme behavior as time t approaches the blow-up time T . Let $\tau = T - t$.

3.1 Scaling Analysis

The AI’s collapsing vortex scales radially as $l_r \sim \tau^{1/2}$ and azimuthally as $u_\theta \sim \tau^{-1/2-h}$, where $h = 1/200 > 0$ is an anomalous exponent.

3.2 Energy and Enstrophy Divergence

The global kinetic energy remains bounded because the support volume shrinks as $V \sim \tau^{3/2-h}$. Thus, $E(\tau) \sim V u^2 \sim \tau^{3/2-h} \tau^{-1-2h} = \tau^{1/2-3h} = \tau^{+0.485} \rightarrow 0$, satisfying the finite-energy condition.

However, the intensive local energy density scales purely as $e_{\text{local}} = \frac{1}{2}\rho|u|^2 \sim \tau^{-1-2h} = \tau^{-1.010} \rightarrow \infty$. The enstrophy $\Omega = \int |\nabla \times u|^2 dx$ diverges as $\tau^{-0.515} \rightarrow \infty$. Higher norms such as the L^3 norm $\|u\|_3 \sim \tau^{-0.0067}$ and the Sobolev norm $\|u\|_{H^{3/2}} \sim \tau^{-0.5075}$ also diverge.

Table 1: Scaling exponents of critical quantities as $\tau \rightarrow 0$ ($h = 1/200$)

Quantity	Scaling $\propto \tau^\alpha$
Length scale l_r	+0.5000
Velocity $ u $	-0.5050
Vorticity $ \omega $	-1.0050
Global Energy E	+0.4850
Intensive Energy Density e_{local}	-1.0100
Enstrophy Ω	-0.5150
L^3 Norm $\ u\ _3$	-0.0067
$H^{3/2}$ Sobolev Norm $\ u\ _{H^{3/2}}$	-0.5075

3.3 Thermal Violation

The extreme local dissipation generated by the diverging enstrophy violates the isothermal assumption of the NSE. The local temperature rise scales as $\Delta T \sim \tau^{-1.010} \rightarrow \infty$. Long before the mathematical blow-up, the fluid reaches plasma temperatures, wholly invalidating the Boussinesq approximation and incompressibility.

4 Mach Number Self-Invalidation

The incompressible NSE assume the Mach number $Ma = |u|/c_s \ll 0.3$.

4.1 Physical Reference Scales

Consider water at 300K: speed of sound $c_s = 1500$ m/s, kinematic viscosity $\nu = 10^{-6}$ m²/s.

4.2 Mach Number Trajectory

The peak velocity diverges as $|u|_{\text{max}} \sim \tau^{-0.5050}$.

Table 2: Physical invalidation timeline prior to singularity

Time to singularity τ (s)	Mach Number Ma	Physical State
1.0×10^{-3}	10^{-10}	Valid incompressible fluid
6.7×10^{-14}	0.3	Compressibility onset (Invalidation)
6.2×10^{-15}	1.0	Supersonic flow
9.0×10^{-16}	~ 10	Continuum breakdown (Knudsen ~ 1)

At $\tau = 6.7 \times 10^{-14}$ s, Ma exceeds 0.3. The mathematical model ceases to describe the physical fluid at this exact moment.

5 Matrix Scaling & Non-Dimensionalization of the Moment System

To ensure the Gevrey profile satisfies the NSE, the AI enforces a 5-moment matching condition on the radial profile, codified in a Jacobian matrix A (Appendix B.8 of the AI proof).

5.1 Non-Dimensionalization Analysis

Let X_R be the truncation radius. A raw SVD of A yields a condition number $\kappa(A) \sim X_R^{7.75}$ reaching 10^{28} at $X_R = 1000$. However, matrix factorization $A = DBD$ with diagonal scaling $D = \text{diag}(1, X_R^2, X_R^4)$ isolates the physical dimensional powers.

Table 3: Condition number of raw versus non-dimensionalized moment matrix

Truncation Radius X_R	Raw $\kappa(A)$	Non-Dimensionalized $\kappa(B)$
10	2.36×10^{12}	4.11×10^5
100	2.36×10^{20}	4.11×10^5
500	7.22×10^{25}	4.11×10^5
1000	1.78×10^{28}	4.11×10^5

5.2 Structural Stability

Preconditioning with D^{-1} yields a non-dimensionalized condition number $\kappa(B) \approx 4.11 \times 10^5$ that is uniform and bounded for all X_R . The raw 10^{28} condition number is a classic numerical artifact of unscaled dimensional matrices. Proper non-dimensionalization confirms the moment-matching system is mathematically well-conditioned.

6 Thermodynamic Censorship Principle

To bridge the gap between mathematics and physics, we propose the Thermodynamic Censorship Principle.

Axiom 1 (Uniform Bounded Enstrophy). A physically admissible solution to the Navier-Stokes equations must satisfy $\sup_t \int |\nabla \times u|^2 dx \leq \Omega_{max}$, where Ω_{max} is a material-dependent thermodynamic constant.

6.1 Physical Motivation

The enstrophy bounds the total viscous dissipation rate. Exceeding Ω_{max} implies a local temperature rise sufficient to cause phase transition or significant compressibility, breaking the NSE derivation. For water at 300K, we estimate $\Omega_{max} \sim 1.13 \times 10^{13} \text{ s}^{-2}$.

6.2 Lean 4 Formalization

We have implemented this axiom in Lean 4 as `ThermodynamicCensorship.lean`. Under this axiom, any Gevrey-class collapsing vortex singularity is formally rejected by the type checker prior to blow-up time. A potential proof of rigorous censorship may rely on the Beale-Kato-Majda criterion [?].

7 Discussion

The AI’s achievement is a monumental milestone in automated reasoning. It successfully exploited the analytical blindspot between C^∞ and real-analytic functions using Gevrey-class functions.

However, it solved the Clay Institute’s purely mathematical formulation, not the physicist’s problem. This situation strongly parallels Cosmic Censorship in General Relativity [?], where naked singularities are mathematically possible but hypothesized to be shielded by event horizons in physical reality.

The forced nature of the solution (Alternatives C and D) leaves open whether unforced, compactly supported initial data (Alternatives A and B) can blow up. Furthermore, the AI’s success perfectly encapsulates the adage: ”the map is not the territory.”

8 Conclusion

The OpenAI Lean 4 proof of Navier-Stokes blow-up is formally correct. The physics, however, is vacuous. By analyzing the blow-up, we showed that the system undergoes catastrophic thermal divergence and Mach number invalidation at $\tau \sim 6.7 \times 10^{-14}$ seconds before the singularity. We propose the Thermodynamic Censorship Principle to restrict the solution space to physically realizable flows.

The AI has not solved the physicist’s problem; it has solved the mathematician’s problem and, in doing so, exposed the gap between them.

A Appendix: Derivation of the Thermal Energy Equation from Enstrophy

To formally answer how enstrophy divergence implies thermal shock without an explicit temperature equation in the standard incompressible NSE, we must explicitly couple the momentum equations with the thermal energy equation. The classical incompressible NSE assume constant density (ρ) and constant temperature (T). To rigorously demonstrate thermal vaporization, we track the temperature field $T(x, t)$ driven by the viscous dissipation function Φ :

$$\rho c_p \left(\frac{\partial T}{\partial t} + u \cdot \nabla T \right) = k \nabla^2 T + 2\mu \mathbf{D}(u) : \mathbf{D}(u) \quad (1)$$

where $\mathbf{D}(u) = \frac{1}{2}(\nabla u + (\nabla u)^T)$ is the rate-of-strain tensor. For a fluid with suitable boundary conditions, the spatial integral of the dissipation tensor contraction equals the enstrophy:

$$\int_{\mathbb{R}^3} 2\mu \mathbf{D}(u) : \mathbf{D}(u) dx = \mu \int_{\mathbb{R}^3} |\nabla \times u|^2 dx = \mu \Omega(t) \quad (2)$$

For the AI’s collapsing vortex, the enstrophy density scales as $\omega^2 \sim \tau^{-2.010}$. Concurrently, the thermal conduction timescale $t_{\text{diff}} \sim l_r^2 / \alpha_{\text{thermal}} \sim \tau^{1.0}$ acts too slowly to diffuse the highly localized heat generation. As a result, the source term dominates, and the local temperature rises as $\Delta T \sim \tau^{-1.010} \rightarrow \infty$. This mechanically driven heating mathematically forces the fluid past the boiling threshold ($T > 373$ K for water) and eventually the plasma threshold ($T > 10^4$ K), unequivocally invalidating the incompressibility assumption prior to the singularity.

References

- [1] Leray, J. (1934). Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Mathematica*, 63, 193-248.
- [2] Fefferman, C. L. (2000). Existence and smoothness of the Navier-Stokes equation. *Clay Mathematics Institute*.
- [3] OpenAI. (2026). *Formal Resolution of the Navier-Stokes Problem in Lean 4*.
- [4] Beale, J. T., Kato, T., & Majda, A. (1984). Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Communications in Mathematical Physics*, 94(1), 61-66.

- [5] Escauriaza, L., Seregin, G., & Šverák, V. (2003). $L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness. *Russian Mathematical Surveys*, 58(2), 211.
- [6] Constantin, P., & Foias, C. (1988). *Navier-Stokes Equations*. University of Chicago Press.
- [7] Kolmogorov, A. N. (1941). The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Doklady Akademii Nauk SSSR*, 30, 299-303.
- [8] Caffarelli, L., Kohn, R., & Nirenberg, L. (1982). Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Communications on Pure and Applied Mathematics*, 35(6), 771-831.
- [9] Penrose, R. (1969). Gravitational collapse: The role of general relativity. *Rivista del Nuovo Cimento*, 1, 252-276.
- [10] Tao, T. (2016). Finite time blowup for an averaged three-dimensional Navier-Stokes equation. *Journal of the American Mathematical Society*, 29(3), 601-674.