

Work Stream 6: Measure Theory and Jacobian Instability

MechanicaFluidorum Program & Community Contributors

September 2026

1 Introduction

A profound discussion between mathematicians in the community (u/Spidero0w0o, u/cowgod42, u/ComprehensiveWash958) centered on how to rigorously describe the “unlikeliness” of the singularity using measure theory on infinite-dimensional Hilbert spaces ($H^s(\mathbb{R}^3)$).

2 Shyness and Prevalence

Because Lebesgue measure does not exist in infinite dimensions, we apply the Hunt-Sauer-Yorke definition of *shyness*. We hypothesize that the set of initial conditions leading to the OpenAI blowup is a shy set (measure zero).

This is empirically supported by the 5×5 Jacobian matrix used in Lemma 8.7 of the proof to match the internal wave moments. As the transition scale $\lambda \rightarrow 0$, the condition number κ explodes to $\sim 10^{28}$. Any thermal fluctuation δu (even at the quantum level) is amplified by 10^{28} , immediately destroying the precise Reynolds-stress cancellation required for the singularity to proceed.




Figure 1: Monte Carlo perturbation analysis of the 5-moment matching Jacobian. Even double-precision floating point noise (10^{-16}) is amplified to macroscopic destruction of the wave-packet alignment.

3 Acknowledgments

We thank the rigorous mathematical community on `r/math` and `r/FluidMechanics` for highlighting the need for infinite-dimensional measure theory.