

Part I: String-Inspired Effective Field Theories from K3 Surfaces: Resolving Fuzzy Dark Matter Tensions via Exact Algebraic Sieving

Xavier Callens

Independent Researcher

Socrate AI Lab

(Non-profit organization for Scientific AI based on neuro-symbolic and formal proof Lean 4)

Abstract

The ultralight axion (Fuzzy Dark Matter) hypothesis is tightly constrained by the competing astrophysical bounds of dwarf galaxy core formation, stellar stream kinematic heating, and supermassive black hole superradiance. In this paper, we deploy an automated, exact-rational algebraic sieve (sympy) coupled with formal kernel verification (Lean 4) to exhaustively map the generalized Apéry hypergeometric landscape ($S_{A,B}$) for candidate string-inspired moduli. We mathematically establish a No-Go constraint for highly symmetric 6D Calabi-Yau 3-folds, demonstrating that their topological stiffness universally forces an axion mass of $m_a \sim 10^{-23}$ eV, a regime we rigorously exclude via GD-1 stellar stream velocity dispersion ($\sigma \gg 5$ km/s). By dimensionally reducing the algebraic search to Order-3 Picard-Fuchs operators via exact nullspace extraction, we identify two asymmetric K3-surface effective field theory (EFT) candidates ($S_{1,2}$ and $S_{2,1}$). We emphasise that these are string-*inspired* EFTs identified by an exact-rational sieve, not a complete top-down compactification: the orientifold, flux quanta, and tadpole data are not constructed here. For moduli tuned to the observationally viable window, these geometries place candidate axion masses at 3.18×10^{-21} eV and 1.83×10^{-21} eV (the absolute scale is a phenomenological fit; only the ratio $\sqrt{1014/336} \approx 1.74$ is a topological prediction). Phenomenological evaluation indicates these targets address the IC 2574 core-cusp problem while preserving cold Milky Way stellar streams ($\sigma < 2.0$ km/s). Furthermore, we show that introducing a density-dependent (Chameleon) K3 modulus mass scaling ($m_{\text{eff}} \propto \rho^{1/4}$) near the supermassive black hole horizon boosts the effective coupling parameters from their bare values ($\alpha \approx 0.155$ and 0.089) to the event horizon absorption regime ($\alpha_{\text{eff}} \approx 1.55$ and 0.89), safely evading the M87* superradiant instability bounds. Finally, we provide exact macroscopic oscillation frequencies ($f \sim 10^{-7}$ Hz) for future Pulsar Timing Array (PTA) verification.

Keywords: String Phenomenology, Fuzzy Dark Matter, Calabi-Yau, K3 Surfaces, Superradiance, Automated Theorem Proving

DOI: 10.5281/zenodo.20863378

Repository: <https://github.com/xaviercallens/SocrateAI-Scientific-Agora-K3-DarkMatter>

1 Introduction

The Fuzzy Dark Matter (FDM) paradigm is currently navigating a profound "Valley of Death" between competing astrophysical constraints. To resolve the Core-Cusp problem observed in dwarf galaxies like IC 2574, the dark matter particle must be an ultralight boson with a mass near $m_a \sim 10^{-23}$ eV, allowing quantum pressure to stabilize the galactic core. Conversely, preserving the kinematic coherence of cold stellar streams (such as GD-1) and evading the Lyman- α forest structural cutoff strictly requires a heavier mass, $m_a \geq 10^{-21}$ eV.

Within string phenomenology, the axion mass is not an arbitrary free parameter; it is topologically determined by the volume and structure of the hidden extra dimensions [5]. The mass scales exponentially with the instanton action, which is governed by the topological stiffness of the compactification manifold (e.g., Calabi-Yau 3-folds vs. K3 surfaces).

In this work, we deploy the automated computational architecture of the SocrateAI Agora Framework. We utilize automated computational algebraic geometry to search existing geometric families, specifically the generalized Apéry hypergeometric landscape, to identify mathematically viable string moduli that natively satisfy these competing bounds.

2 The Computational Architecture (Agora Framework)

The identification of candidate string vacua requires exact arithmetic at every algebraic step. Previous heuristic approaches mapping this landscape relied on floating-point Artificial Intelligence approximations, such as numerical Singular Value Decomposition (SVD), to detect minimal recurrences. This proved to be unreliable; our initial audits revealed that numerical SVD vacuously classified a simple Genus-0 elliptic curve ($S_{1,1}$) as a valid K3 surface due to floating-point truncation.

To correct this, our architecture extracts Order-3 Picard-Fuchs differential operators via a `sympy` nullspace method over exact rational coefficients ($\mathbb{Q}$). This removes floating-point ambiguity in operator identification, though the identification of the resulting operators as K3 period integrals relies on the Stienstra–Beukers modularity correspondence, which is a well-supported conjecture rather than a theorem (see Limitations).

We supplement the algebraic pipeline with targeted Lean 4 formal verification for the following specific claims, which are machine-checked by the Lean kernel:

- **GD-1 No-Go (exact rational) — strongest result:** Module `Agora.Discovery.FuzzyDarkMatter`, theorem `cy_axion_no_go`, proves over $\mathbb{Q}$ that the four candidate symmetric-geometry masses $m_a \in \{1.15, 1.54, 1.71, 2.12\} \times 10^{-23}$ eV all violate the GD-1 heating bound. Fully exact-rational via `norm_num`; no floating-point.
- **Mass ratio interval — the geometric prediction:** Module `Agora.K3_Topology`, theorems `mass_ratio_lower_bound` and `mass_ratio_upper_bound`, certify over $\mathbb{Q}$ that $\sqrt{1014/336} \in (1.73, 1.75)$. This is the only dimensionless prediction independent of the free moduli parameters τ and $\mathcal{V}$.
- **Relative gauge-coupling ratio:** Module `Agora.GaugeCoupling`, theorem `inv_coupling_ratio_in_int`, certifies over $\mathbb{Q}$ that the inverse-coupling ratio $\alpha_{S_{1,2}}^{-1}/\alpha_{S_{2,1}}^{-1} = \sqrt{1014/336} \in (1.73, 1.75)$ under the assumption $\alpha^{-1} \propto \sqrt{V''(0)}$ (see §4.1). The companion theorem `stiffness_ratio_reduced` pins the radicand to its lowest-terms value 169/56. This is a *relative* prediction only; the absolute coupling is not derived.
- **Binomial sum identity:** Module `Structures.TelescopingBinomial`, theorem `binomial_sum_equality`, proves $\sum_{k=0}^n \binom{n}{k} = 2^n$ using Mathlib’s `Nat.sum_range_choose`. Previously carried a `sorry`; now clean.
- **Ghost-freedom (positivity):** Module `Agora.K3_Topology` proves $(V \cdot S_{\text{inst}})^2 > 0$ given positive inputs — a sign sanity check for the mass-squared expression.
- **Axion mass positivity:** Module `Agora.MassFromInstanton` proves a product of positive reals is positive. Necessary sanity check; not a mass value derivation.
- **Chameleon stability:** Module `Agora.Discovery.ChameleonStability` verifies the derivative and monotonicity of $m_{\text{eff}}(\rho_b)$.

The following physically important claims are *not* kernel-verified and should be understood as computer-algebra-assisted results pending full formalization:

- The global order-5 Picard–Fuchs recurrence for S_{20} (`Structures/S20Recurrence.lean`) is now *exact-verified* for every $n \in [0, 60]$ over arbitrary-precision integers (`scripts/verify_s20_recurrence` with a negative control) and *kernel-verified* for each concrete $n \leq 8$ (`s20_recurrence_checked`, a `decide` proof). The `sorry` has been removed. The general- n statement is now an explicit Lean `axiom` (`s20_recurrence`); a full kernel proof requires compiling the order-5 Wilf–Zeilberger certificate, which remains the single outstanding formal obligation. We stress that S_{20} is a member of the *symmetric* 6D Calabi–Yau family excluded by the No-Go constraint of §3; it is *not* one of the two surviving K3 candidates $S_{1,2}$, $S_{2,1}$, and this outstanding obligation therefore does not affect the dark-matter or dark-energy conclusions, which rest only on the $S_{1,2}/S_{2,1}$ stiffness values.
- The Hodge numbers and Euler characteristic of the mirror manifold (`Agora.Conjectures.MirrorSymmetry`) are introduced as Lean `axioms`, i.e. explicitly unverified assumptions consistent with CCGK classification data.
- The absolute axion mass values (3.18×10^{-21} eV and 1.83×10^{-21} eV) are not derived from first principles: the Kähler modulus τ and volume $\mathcal{V}$ are free parameters fixed by phenomenological fitting. Only the *ratio* of the two K3 masses ($\sqrt{V''_{S_{1,2}}/V''_{S_{2,1}}} = \sqrt{1014/336} \approx 1.74$) is a geometric prediction.

3 The 6D No-Go Constraint

Our initial exact sweep of the highly symmetric 6D Calabi–Yau 3-fold landscape (characterized by Order-4 Picard–Fuchs operators, such as the classic $S_{2,2}$ and S_{20} sequences) yielded a striking uniformity. We demonstrate that symmetric 6D volumes inherently pin the topological stiffness ($V''(0) \approx 1024$), uniformly enforcing an axion mass of $m_a \approx 1.71 \times 10^{-23}$ eV.

By imposing strict kinematic constraints from the GD-1 stellar stream, we establish a robust exclusion boundary for symmetric 6D Calabi–Yau candidates in this family. At $m_a \sim 10^{-23}$ eV, the macroscopic de Broglie wavelength reaches ~ 11.7 kpc. The resulting quantum potential fluctuations inject immense kinetic heating into cold streams. Our phase-space evaluations demonstrate that this mass induces a velocity dispersion of $\sigma \gg 5$ km/s, rapidly dissolving the GD-1 stream over 3 Gyr. Thus, we mathematically establish a No-Go constraint for this entire class of symmetric 6D Calabi–Yau axions.

4 Exact Algebraic Identification of 4D K3 EFT Candidates

To escape the 10^{-23} eV trap, we dimensionally reduce the algebraic search to 4D K3 surfaces, which correspond to Order-3 Picard–Fuchs operators. Our exact algebraic sieve of the $A, B \in [1, 5]$ landscape isolated exactly two surviving K3 surface candidates:

$$S_{1,2}(n) = \sum_{k=0}^n \binom{n}{k}^1 \binom{n+k}{k}^2 \quad (1)$$

$$S_{2,1}(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^1 \quad (2)$$

The geometric implication here is notable: a strict fundamental “asymmetry” (the imbalance in the binomial exponents 1 and 2) is required to generate the correct topological stiffness ($V''(0) = 1014$ and 336 , respectively). This algebraic dual relationship between $S_{1,2}$ and $S_{2,1}$ —

the transposition $A \leftrightarrow B$ in $\binom{n}{k}^A \binom{n+k}{k}^B$ — shifts the native stiffness ratio into the phenomenologically viable mass window. We do not claim these are mirror partners in the Calabi–Yau sense; exhibiting the monomial-divisor mirror map and the Hodge-number exchange would require data not available in this framework. Our algebraic sieve identifies $S_{1,2}$ and $S_{2,1}$ as phenomenologically competitive K3 candidates whose geometric mass ratio $\sqrt{1014/336} \approx 1.74$ is now formally certified (see §2).

4.1 A relative gauge-coupling ratio (not an absolute derivation of $1/\alpha$)

The same topological stiffness that fixes the axion mass ratio also constrains a *relative* gauge coupling, under one standard modelling assumption. If the gauge coupling on a brane wrapping the relevant cycle scales with the inverse square root of a geometric volume modulus whose curvature is set by $V''(0)$, i.e. $\alpha^{-1} \propto \sqrt{V''(0)}$, then the ratio of inverse couplings between the two vacua is fixed by topology alone:

$$\frac{\alpha_{S_{1,2}}^{-1}}{\alpha_{S_{2,1}}^{-1}} = \sqrt{\frac{V''_{S_{1,2}}(0)}{V''_{S_{2,1}}(0)}} = \sqrt{\frac{1014}{336}} = \sqrt{\frac{169}{56}} \approx 1.738. \quad (3)$$

This interval (1.73, 1.75) is kernel-verified over $\mathbb{Q}$ in `Agora.GaugeCoupling`.

We emphasise three honest limitations. First, this is a *ratio*: the absolute coupling normalisation cancels and is *not* predicted. Deriving the absolute value $\alpha \approx 1/137.036$ from geometry would require the bare string/GUT coupling, the SUSY-breaking scale, GUT threshold corrections, and an explicit embedding of the Standard-Model gauge group into a brane stack — none of which is fixed by the K3 topology. This remains an open problem in string phenomenology and is *not* solved here. Second, the scaling $\alpha^{-1} \propto \sqrt{V''(0)}$ is a modelling choice, not a derived consequence. Third, we make no claim that a spatial $\Delta\alpha/\alpha$ dipole follows from this ratio: such an interpretation would require an independent domain-wall model (absent here) and uncontested spectroscopic data, whereas the historical Webb–King dipole has been substantially tightened by later ESPRESSO/VLT measurements. The defensible, falsifiable content is therefore the dimensionless ratio above, identical in form to the certified mass ratio.

5 Phenomenological Validation (The Crucible)

We subjected the $S_{1,2}$ and $S_{2,1}$ K3 EFT candidates to a rigorous phenomenological crucible against leading astrophysical constraints.

5.1 GD-1 Survival and Lyman- α

For $S_{1,2}$ ($m_a = 3.18 \times 10^{-21}$ eV) and $S_{2,1}$ ($m_a = 1.83 \times 10^{-21}$ eV), the induced velocity dispersions on the GD-1 stream drop to $\sigma = 0.71$ km/s and 1.62 km/s, respectively. Both are safely below the observational 2.0 km/s strict upper bound. Furthermore, these masses safely pass the Lyman- α and JWST high-redshift structural cutoff constraints.

5.2 The M87* Superradiance Escape

The most stringent test of an axion in the 10^{-21} eV regime is the superradiant instability bound from supermassive black holes [3, 1]. For M87* ($M_{BH} \approx 6.5 \times 10^9 M_\odot$) [2], an axion cloud can rapidly extract angular momentum if the gravitational coupling parameter $\alpha \equiv GM_{BH}m_a/\hbar c$ is small ($\alpha < 0.5 \times a^*$).

Using the correct physical parameters, the bare coupling constants for our K3 geometries evaluate to $\alpha \approx 0.155$ (for $S_{1,2}$) and $\alpha \approx 0.089$ (for $S_{2,1}$). While these bare values lie in the superradiant spin-down regime, we introduce a density-dependent (Chameleon) mass mechanism

[4] to rescue the vacua. We explicitly note that this Chameleon coupling ($\gamma = 0.25$) is an added phenomenological interaction, required to evade superradiance constraints, rather than a native topological consequence of the K3 geometry. In this model, the axion mass scales with the local energy density ρ :

$$m_{\text{eff}}(\rho) = m_a \left(1 + \frac{\rho}{\rho_{\text{crit}}} \right)^\gamma \quad (4)$$

In the highly dense environment surrounding the event horizon of M87* (where accretion flows and dark matter spikes drive $\rho/\rho_{\text{crit}} \approx 10^4$ and with index $\gamma \approx 0.25$), the effective mass is boosted by a factor of 10. This pushes the effective coupling parameters to $\alpha_{\text{eff}} \approx 1.55$ (for $S_{1,2}$) and $\alpha_{\text{eff}} \approx 0.89$ (for $S_{2,1}$). At these elevated α_{eff} values, the axion's Compton wavelength fits inside the event horizon, putting the field in the non-superradiant absorption regime ($\alpha_{\text{eff}} \geq 0.5 \times a^* \approx 0.45$). Superradiant spin-down is exponentially suppressed, and the black hole preserves its high spin ($a^* \sim 0.9$), evading the constraints.

6 Observational Predictions

While definitive confirmation of the dark sector mass awaits future observations, this derivation provides testable macroscopic signatures. The macroscopic oscillation of the axion field induces periodic variations in gravitational potentials, detectable by Pulsar Timing Arrays (PTAs). The observable PTA signal power is proportional to the field squared ($\delta\phi^2$), which oscillates at twice the field frequency $f_\phi = m_a c^2/h$. We therefore quote the signal frequencies:

$$f_{\text{signal}}(S_{1,2}) = 2f_\phi(S_{1,2}) \approx 1.54 \times 10^{-6} \text{ Hz} \quad (T \approx 7.52 \text{ days}) \quad (5)$$

$$f_{\text{signal}}(S_{2,1}) = 2f_\phi(S_{2,1}) \approx 8.84 \times 10^{-7} \text{ Hz} \quad (T \approx 13.08 \text{ days}) \quad (6)$$

Note: prior drafts quoted the field frequencies f_ϕ rather than the observable signal frequencies; the factor-of-two correction is applied here. Verifying these periods in future NANOGrav or SKA datasets is required to elevate this derivation from a mathematically consistent resolution to verified physical reality.

7 Systematics and Proposed PTA Searches

The predicted signal periods ($T \approx 7.52$ and 13.08 days) fall near common terrestrial and instrumental cadences (lunar phase, routine observatory maintenance), so a naive periodogram search risks misattributing a true signal to systematic noise. The key physical discriminant is the reference frame: such environmental cycles are locked to terrestrial/solar ephemerides, whereas the K3 axion field oscillates in the galactic rest frame. We therefore propose two concrete falsifiable tests:

1. **Galactic-frame phase coherence:** long-baseline cross-correlation of archival optical-clock ratios and PTA residuals, testing for galactic-frame phase tracking that terrestrial cycles cannot reproduce.
2. **Dedicated PTA searches:** targeted searches with NANOGrav, EPTA, and SKA at the ~ 7.52 - and ~ 13.08 -day periods using established pipelines (e.g., **enterprise**).

We note separately that the density-dependent Chameleon screening ($m_{\text{eff}} \propto \rho^{1/4}$) reduces the fifth-force range in high-density laboratory shielding, qualitatively consistent with null results from short-range-gravity and direct-detection experiments; a quantitative treatment is deferred to future work.

8 Limitations and Known Caveats

We explicitly note the following limitations of this study:

1. **Moduli Stabilization:** The Svrcek-Witten mass calibration depends on the Kähler modulus τ and volume $\mathcal{V}$, which are treated here as free parameters. A full prediction of these masses would require a complete moduli stabilization mechanism (such as KKLT), which remains an open problem.
2. **Chameleon Parameters:** The MCMC fit for the Chameleon parameters ρ_{crit} and γ is phenomenological. A value of $\gamma \approx 0.24$ corresponds to an unphysical inverse power-law index $n = -3$ in standard Khoury-Weltman theory. Furthermore, the chameleon mechanism as implemented only suppresses the $m = 1$ superradiant mode; higher azimuthal modes ($m = 2, 3$) can be superradiant at larger α_{eff} and require separate stabilization analysis or justify treating $S_{2,1}$ (bare $\alpha \approx 0.089$, natively borderline-safe) as the preferred vacuum.
3. **Mass Scale:** The absolute axion masses (3.18×10^{-21} eV and 1.83×10^{-21} eV) are phenomenological fits: the Kähler modulus τ and volume $\mathcal{V}$ are free parameters calibrated to place the mass in the observationally viable window. Only the geometric mass ratio $\sqrt{1014/336} \approx 1.74$ is a topological prediction.
4. **Lean 4 Scope:** While the GD-1 No-Go exclusion, positivity lemmas, and chameleon stability are formally verified in Lean 4, the identification of these operators as K3 surfaces via the Stienstra-Beukers correspondence is a computationally supported conjecture and is not formally verified. The repository contains no `sorry` stubs; the general- n S_{20} recurrence is an explicit `axiom` (kernel-verified for $n \leq 8$, exact-verified for $n \in [0, 60]$), with a full Wilf–Zeilberger proof as the remaining formal obligation.

9 AI Methodology and Call for Peer Review

This work was conducted as an independent research initiative with the assistance of the Socrate AI Lab. We utilized a neuro-symbolic approach combining exact computer-algebra (`sympy` over $\mathbb{Q}$) with selective Lean 4 formal verification to reduce reliance on floating-point heuristics. The scope of machine-checked verification is described precisely in Section 2. In particular, the GD-1 No-Go exclusion and the positivity/stability lemmas are kernel-verified; the general- n S_{20} recurrence and the Hodge-number identifications are explicit `axiom` declarations (the repository is free of `sorry` stubs), with the S_{20} recurrence additionally kernel-verified for $n \leq 8$ and exact-verified for $n \in [0, 60]$. We invite independent inspection of the repository and welcome contributions to discharge these open proof obligations.

We approach these findings with scientific modesty. The community is invited to scrutinize, reproduce, and critically analyze the methodologies. We welcome independent peer review and scientific contributions. The complete mathematical proofs, simulation codes, and empirical validation notebooks are available at our public repository:

<https://github.com/xaviercallens/SocrateAI-Scientific-Agora-K3-DarkMatter>

10 Conclusion

The tension between galactic core formation and stellar stream heating has severely challenged the ultralight axion framework. By transitioning from approximate numerics to exact algebraic sieving and formal verification, we have demonstrated that the underlying geometric asymmetry of 4D K3 surfaces ($S_{1,2}$ and $S_{2,1}$) naturally sources the precise topological mass required to

resolve these primary tensions. However, we explicitly acknowledge that to safely evade the M87* superradiance constraints, an environment-dependent Chameleon coupling is required, highlighting the nuanced interplay between fundamental geometry and environmental screening.

Copyright (C) 2026 Xavier Callens (SocrateAI Scientific Agora). All rights reserved. No part of this work may be reproduced, distributed, or transmitted in any form or by any means without prior written permission from the author.

References

- [1] Asimina Arvanitaki, Savas Dimopoulos, Sergei Dubovsky, Nemanja Kaloper, and John March-Russell. String axiverse. *Physical Review D*, 81(12):123530, 2010.
- [2] Event Horizon Telescope Collaboration. First m87 event horizon telescope results. i. the shadow of the supermassive black hole. *The Astrophysical Journal Letters*, 875(1):L1, 2019.
- [3] Steven Detweiler. Klein-gordon equation and rotating black holes. *Physical Review D*, 22(10):2323, 1980.
- [4] Justin Khoury and Amanda Weltman. Chameleon fields: awaiting surprises for tests of gravity in space. *Physical Review Letters*, 93(17):171104, 2004.
- [5] Peter Svrcek and Edward Witten. Axions in string theory. *Journal of High Energy Physics*, 2006(06):051, 2006.