

Part II: Project Vafa-Continuity: A String-Inspired $K3 \times T^2$ Quintessence Model and the Swampland Obstruction to Stable Dark Energy

Xavier Callens (Independent researcher)

Socrate AI Lab (none profit organization)

callensxavier@gmail.com

June 2026

DOI: 10.5281/zenodo.20863378

GitHub: SocrateAI-Scientific-Agora-K3-DarkMatter

Abstract

We study the T^2 volume modulus of a $K3 \times T^2$ compactification as a rolling quintessence field and examine its cosmological implications. The best-fit coupling $\lambda \approx 1.6724$ yields a thawing trajectory $w_0 \approx -0.5485$, $w_a \approx -0.3968$, lying outside the DESI 2024 1σ contour. The primary result of this work is the quantitative recovery of the Agrawal–Obied–Steinhardt–Vafa quintessence–Swampland tension in a concrete $K3 \times T^2$ context: since $\lambda_{\text{fit}} = 1.6724 > \sqrt{2}$, the asymptotic scaling attractor gives $w_\phi \approx -0.07$ — not dark energy — while the Swampland bound demands $\lambda \gtrsim 1$. The compatible window $1 \lesssim \lambda < \sqrt{2}$ is narrow ($\Delta\lambda \approx 0.41$) and our best-fit lies outside it. Both facts are formally certified by new Lean 4 theorems (`lambda_fit_exceeds_sqrt2`, `attractor_not_dark_energy`) over exact rationals. In other words, dark energy in this concrete geometry is necessarily a *transient phase* rather than a stable cosmological constant — precisely the obstruction the Swampland conjectures predict for any quasi-de Sitter quintessence. We present this not as a deficiency of the model but as its central, falsifiable physics result. Three resolution programmes for an accelerating epoch are identified: multi-field potentials, hilltop/plateau modifications, and transient dark energy embeddings. We also show that a coupled axion mass variation of 19% ($\Delta\phi \approx 0.37 M_{\text{pl}}$, sub-Planckian) produces an indicative $H_0 \sim 72$ km/s/Mpc shift; this omits a full Boltzmann treatment and is not a precision prediction. All Lean proof scope, open obligations, and caveats are enumerated precisely.

1 Introduction

The ultralight axion (Fuzzy Dark Matter) hypothesis is one of the most compelling alternatives to standard cold dark matter, but it is tightly constrained by competing astrophysical bounds. Historically, resolving dwarf galaxy core-cusp problems (such as in

IC 2574) requires a light boson mass ($m_a \sim 10^{-23}$ eV) to produce a sufficiently large de Broglie wavelength [?]. However, constraints from stellar stream heating (e.g., the GD-1 stream) [?] and black hole superradiance (both vector and scalar instabilities) [?, ?, ?, ?] force the axion mass into a much heavier regime ($m_a \geq 10^{-21}$ eV).

Within string phenomenology, axion masses are topologically determined by the volume and geometry of the compactified extra dimensions [?]. The topological stiffness of a compactification manifold dictates the instanton action, which exponentially suppresses the axion mass. While rigid 6D Calabi-Yau 3-folds typically force axion masses into the excluded 10^{-23} eV regime, asymmetric 4D $K3$ surfaces naturally yield candidate masses of 1.83×10^{-21} eV and 3.18×10^{-21} eV, resolving these primary dark matter tensions.

However, embedding these $K3$ surfaces into a full 6D string geometry via a T^2 Torus fibration introduces an elastic volume modulus, V_{T^2} . The slow expansion of this extra-dimensional volume over cosmic time acts as a rolling quintessence field [?], which provides the negative pressure required for Dark Energy. This paper details Project Vafa-Continuity: the unified field equations governing this coupled dark sector, its cosmological parameters, and its formal mathematical validation.

2 Related Work

The dynamics of dark energy modeled as a slowly rolling scalar field (quintessence) has been studied extensively, with exponential potentials providing tracker behavior that helps alleviate cosmic coincidence problems [?]. In string theory, such scalar fields are identified with moduli describing the shape and size of the compactification manifold [?]. The Swampland Conjectures (specifically the de Sitter and Distance Conjectures) place tight bounds on the slope of these modulus potentials, requiring $|\nabla V|/V \geq \mathcal{O}(1)$ in Planck units [?, ?].

Fuzzy Dark Matter (FDM) constraints have been rigorously extracted from the Lyman- α forest power spectrum [?, ?] and dark matter subhalo stream heating [?], indicating that $m_a < 10^{-21}$ eV is strongly disfavored unless screening mechanisms are present. The Chameleon screening model [?, ?] provides a density-dependent mass boost that allows axions to evade solar system fifth-force tests and supermassive black hole superradiance limits. Recent observations of the supermassive black hole in M87* by the Event Horizon Telescope (EHT) [?] and investigations into jet precession [?] confirm high spin values ($a^* \sim 0.9$), which would be unstable if a light, unscreened boson existed in the superradiant range.

From an algebraic perspective, Calabi-Yau and $K3$ moduli spaces are governed by Picard-Fuchs differential equations, whose coefficients can be extracted from Apéry-like recurrence relations [?, ?, ?]. This mathematical infrastructure allows for the exact determination of $K3$ topological stiffness and mirror map structures.

3 AI Methodology and Peer Review

This research was conducted utilizing a neuro-symbolic approach combining exact computer-algebra (`sympy` over $\mathbb{Q}$) with selective Lean 4 formal verification. The scope of machine-checked proofs is limited and precisely enumerated in Part I, Section 2: the GD-1 No-Go exclusion and positivity lemmas are kernel-verified; the general- n S_{20} Picard-Fuchs recurrence and the Hodge-number assignments are explicit Lean `axioms` (no `sorry` stubs

remain in the repository), with the S_{20} recurrence additionally kernel-verified for $n \leq 8$ and exact-checked for $n \in [0, 60]$. We present this as independent work and actively call for peer review, reproduction, and criticism. All code and formal proofs are available in the public GitHub repository.

4 Methodology

To model the dynamic dark energy, we treat the T^2 volume modulus r (parameterized as a dimensionless scalar field ϕ) rolling down an exponential potential:

$$V(\phi) = V_0 \exp(-\lambda\phi) \quad (1)$$

We perform an active numerical integration of the coupled Klein-Gordon and Friedmann equations from the deep radiation-matter era ($a_{\text{init}} = 10^{-3}$) to the present ($a = 1$):

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0 \quad (2)$$

$$H^2 = \frac{1}{3} \left[\rho_r(a) + \rho_m(a) + \frac{1}{2}\dot{\phi}^2 + V(\phi) \right] \quad (3)$$

The integration is carried out using SciPy's `solve_ivp` solver with the Radau method. To ensure energy budget closure, we implement a bisection shooting method for the potential normalization V_0 such that the dark energy density parameter at the present epoch matches the target value:

$$\Omega_\phi(a = 1) = 1.0 - \Omega_{m0} - \Omega_{r0} \approx 0.685 \quad (4)$$

We fit the resulting dynamic equation of state:

$$w(a) = \frac{\frac{1}{2}\dot{\phi}^2 - V(\phi)}{\frac{1}{2}\dot{\phi}^2 + V(\phi)} \quad (5)$$

to the Chevallier-Polarski-Linder (CPL) parametrization:

$$w(a) = w_0 + w_a(1 - a) \quad (6)$$

where w_0 and $w_a \equiv -dw/da|_{a=1}$ are evaluated at the present epoch.

To address the Hubble tension, we couple the K3 axion mass to the Torus volume modulus: $m_a \propto (V_{T^2})^{-\alpha}$ with $\alpha = 1/2$ [?]. The expanding volume causes the axion mass to decrease over time. The mass-varying dark matter fluid density scales as:

$$\rho_{DM}(a) \propto m_a(a)a^{-3} \propto a^{-3-\epsilon} \quad (7)$$

where $\epsilon = 0.042/\lambda$ represents the mass-varying coupling parameter. The constant 0.042 is a phenomenological calibration: it is the value of $\epsilon\lambda$ that produces a sufficient shift in the acoustic scale to move H_0 by ~ 3 km/s/Mpc for $\lambda \approx 1.67$, determined by reverse-engineering the target $H_0 \sim 72$ km/s/Mpc from the acoustic constraint equation rather than derived from the $K3 \times T^2$ geometry. We integrate the sound horizon at recombination $r_s(z_*)$ and the angular diameter distance $D_A(z_*)$ using:

$$r_s(z_*) = \int_{z_*}^{\infty} \frac{c_s(z)}{H(z)} dz \quad (8)$$

$$D_A(z_*) = \frac{1}{1 + z_*} \int_0^{z_*} \frac{1}{H(z)} dz \quad (9)$$

and solve for the Hubble parameter H_0 by enforcing the CMB acoustic scale constraint measured by Planck 2018 [?]:

$$\theta_s \equiv \frac{r_s(z_*)}{(1+z_*)D_A(z_*)} = 0.010411 \quad (10)$$

5 Results

Our parameter sweep over the coupling constant λ yields a highly constrained set of quintessence models. Table ?? displays the resulting CPL parameters and current dark energy fraction.

Table 1: Quintessence Parameter Sweep and CPL Fitting Results

λ	V_0	$\Omega_\phi(a=1)$	w_0	w_a
0.5000	2.1245	0.6849	-0.9634	-0.0382
1.0000	2.7482	0.6849	-0.8521	-0.1584
1.5000	4.1012	0.6849	-0.6391	-0.3298
1.6724	4.8005	0.6849	-0.5485	-0.3968
2.0000	6.8291	0.6849	-0.3812	-0.4901

The best-fit coupling parameter $\lambda \approx 1.6724$ yields $w_0 \approx -0.5485$ and $w_a \approx -0.3968$. As visible in Figure ??, this best-fit point lies outside the DESI 2024 1σ contour (centred near $w_0 \approx -0.727$, $w_a \approx -1.05$ [?]), representing a qualitative thawing trajectory in the correct direction but remaining in tension with the observed region.

Crucially, the best-fit $\lambda = 1.6724 > \sqrt{2} \approx 1.414$ places the model above the threshold for accelerated expansion on the scaling attractor. The attractor equation of state for a single-exponential potential is $w_\phi = -1 + \lambda^2/3 \approx -0.07$, which gives *no* accelerated expansion. The observed $w_0 = -0.55$ is a transient thawing value off this attractor. This is the Agrawal–Obied–Steinhardt–Vafa quintessence-Swampland tension [?], instantiated in a concrete $K3 \times T^2$ geometry (see §??).

For this best-fit model, the mass-varying dark matter coupling parameter evaluates to $\epsilon \approx 0.0251$, which yields an early-universe axion mass ratio of:

$$\frac{m_a(z=1100)}{m_a(z=0)} \approx 1.1920 \quad (11)$$

Evaluating the acoustic scale constraint $\theta_s = 0.010411$ under this mass-varying regime shifts the predicted Hubble parameter from the Λ CDM baseline of $H_{0,\text{baseline}} \approx 68.31$ km/s/Mpc to an order-of-magnitude estimate of $H_0 \sim 72$ km/s/Mpc, directionally consistent with local distance ladder measurements. We stress that this estimate uses rest initial conditions ($\dot{\phi} = 0$ at $a = 10^{-3}$) and omits the full Boltzmann perturbation treatment; the quoted value should be interpreted as an indicative shift, not a precision prediction. Closing the Hubble tension rigorously requires injection of the interacting modulus fluid into a full Boltzmann solver (see Limitations).

6 Swampland Formal Verification

The de Sitter Swampland conjecture [?], refined in [?], asserts that any consistent potential arising from string theory must satisfy $|\nabla V|/V \geq c$ for some $c = \mathcal{O}(1)$. For




Figure 1: The w_0 - w_a plane. The orange point is the asymptotic scaling attractor ($w_\phi = -1 + \lambda^2/3 \approx -0.07$) toward which the model evolves; the blue point is the current thawing transient. Neither lies within the DESI 2024 1σ contour (red). The dashed arrow indicates the future trajectory, quantitatively recovering the quintessence-Swampland tension of [?].

a single-exponential potential $V = V_0 e^{-\lambda\phi}$ this bound reduces to the statement $\lambda \geq c$, which is a constraint on the model parameter rather than a derived consequence of the geometry. Agrawal et al. [?] showed that satisfying this bound with $c \geq 1$ while also fitting late-time acceleration creates a fundamental tension with observation, a problem this model does not yet resolve (see Limitations).

We use the Lean 4 Theorem Prover [?] with the Mathlib library [?] to formally verify the following specific, kernel-checked statement about the effective potential:

- **Derivative identity (machine-checked):** The theorem `V_has_deriv_at` in module `Agora.SwamplandK3T2` proves that $\nabla V = -c V_0 e^{-cr}$ is the exact analytic derivative of $V(r) = V_0 e^{-cr}$. This is a verified calculus identity.
- **Gradient-to-value ratio (machine-checked):** The theorem `swampland_bound` proves the algebraic identity $|\nabla V| = c \cdot V$ given $V_0 > 0$, $c > 0$. This establishes that the ratio $|\nabla V|/V$ evaluates to the *parameter* c .

Two additional theorems in `Agora.SwamplandK3T2` kernel-verify the tension itself:

- **lambda_fit_exceeds_sqrt2 (machine-checked):** Proves over $\mathbb{Q}$ that $1.6724 > 1.4143$ and $(1.4143)^2 > 2$, certifying $\lambda_{\text{fit}} > \sqrt{2}$.
- **attractor_not_dark_energy (machine-checked):** Proves over $\mathbb{Q}$ that $(1.6724)^2/3 > 1/3$, certifying $w_{\text{attractor}} + 1 > 0$, i.e. the attractor gives no accelerated expansion.

The following points are *not* established by the Lean proof and must be stated explicitly:

- The inequality $c \geq \mathcal{O}(1)$ is not a theorem; it holds trivially because we *choose* $\lambda = 1.6724 > 1$ as a phenomenological fit. The Lean kernel certifies arithmetic, not the physical bound.
- No `sorry` stubs appear in `SwamplandK3T2.lean` itself; the broader Lean library is now free of `sorry` stubs, but it does contain explicit `axiom` declarations (the general- n `Structures/S20Recurrence` law, `Agora.Conjectures.MirrorSymmetry` Hodge data, and the Fano supercongruences). The claim of zero axioms does not hold for the full library; each axiom is disclosed in `OPEN_PROBLEMS.md`.

- **Distance Conjecture constraint.** The Distance Conjecture [?] implies that a field excursion $\Delta\phi$ of the T^2 modulus lowers a tower of states by $m_{\text{tower}} \sim m_0 e^{-\alpha\Delta\phi}$. The 19% axion mass drop from $z = 1100$ to $z = 0$ implies $\Delta V_{T^2}/V_{T^2} \approx 0.44$, and for a canonically normalised modulus $\phi = M_{\text{pl}} \log V_{T^2}$ this gives $\Delta\phi \approx 0.37 M_{\text{pl}}$. The excursion is therefore sub-Planckian, suppressing the tower by $e^{-\alpha \cdot 0.37} \approx 0.7$ for $\alpha = \mathcal{O}(1)$. The tower states are generically Kaluza–Klein modes of T^2 with mass $m_{\text{KK}} \sim V_{T^2}^{-1/2}$, whose 19% shift is consistent with the axion mass variation by construction. A rigorous analysis must verify that no lighter tower states descend into the EFT validity window during the field evolution.

7 The Quintessence-Swampland Tension as a Quantitative Result

The results above instantiate a known structural tension in the Swampland programme. For a single-exponential potential $V = V_0 e^{-\lambda\phi}$, the Friedmann attractor equation of state is:

$$w_{\phi}^{\text{attractor}} = -1 + \frac{\lambda^2}{3} \quad (12)$$

Accelerated expansion requires $w < -1/3$, i.e. $\lambda < \sqrt{2} \approx 1.414$. The de Sitter Swampland conjecture requires $\lambda \gtrsim \mathcal{O}(1)$. The allowed window is therefore $1 \lesssim \lambda < \sqrt{2}$, a narrow interval of width ≈ 0.41 . The best-fit value $\lambda = 1.6724$ falls *outside* this window:

$$\lambda_{\text{fit}} = 1.6724 > \sqrt{2} \approx 1.414 \quad (\text{no attractor acceleration}) \quad (13)$$

$$w_{\phi}^{\text{attractor}} = -1 + \frac{(1.6724)^2}{3} \approx -0.07 \quad (\text{not dark energy}) \quad (14)$$

This quantitatively recovers the tension first analyzed by Agrawal, Obied, Steinhardt, and Vafa [?] in the concrete context of a $K3 \times T^2$ compactification. The observed $w_0 = -0.55$ is a transient thawing value; the asymptotic attractor is matter-like.

Three paths to resolution exist and remain open:

1. **Multi-field potential.** A second modulus could steepen the effective slope at late times while satisfying the Swampland bound at all points on the trajectory.
2. **Hilltop or plateau potential.** A Swampland-compatible potential with a shallow region near $\phi = 0$ (where $\lambda_{\text{eff}} < \sqrt{2}$) and a steep tail (where $\lambda_{\text{eff}} \gtrsim 1$) can in principle satisfy both constraints.
3. **Transient dark energy.** Accept that the current dark energy epoch is transient, and embed the model in a cosmology that predicts the future transition to matter-domination and the associated observational signatures.

Pursuing any of these three programmes within the $K3 \times T^2$ framework is the primary open physics problem of this project.

8 Limitations and Known Caveats

We explicitly note several theoretical and numerical limitations of the current framework:

1. **Moduli Stabilization:** The K3 volume and the Kähler modulus τ are not uniquely fixed by topology alone, requiring a complete moduli stabilization mechanism (e.g., KKLT or Large Volume Scenario) which is not explicitly modeled here [?].
2. **Initial Conditions (updated 2026-07-11, Task T5.1):** The quintessence solver initializes the scalar field from rest ($\phi = 0, \dot{\phi} = 0$) at $a = 10^{-3}$. We have now computed the Copeland-Liddle-Wands (1998) tracker/attractor trajectory instead (`empirical_crucible/tracker_ics.py`); for this potential the tracker and rest initial conditions converge to nearly the same late-time w ($\Delta w_0 \approx 0.0002$) — a modest, real result rather than an unresolved open question. Full detail: `docs/cosmology/ic_sensitivity.md`.
3. **Perturbation Theory (updated 2026-07-11, Task T5.2):** A real CLASS (`classy`) Boltzmann computation was executed. Two findings: (i) *Architectural blocker, confirmed live*: this model’s own formula $\rho_{DM}(a) \propto a^{-3-\epsilon}$ implies $w_{DM} = \epsilon/3 \approx +0.0084 > 0$, which CLASS’s public dark-energy-fluid API hard-rejects (any fluid with $w(a \rightarrow 0) \geq 0$) — a genuine C-source fork of CLASS (`background.c`, `perturbations.c`) is required and was *not* attempted, since a hand-written perturbation patch would have no independent benchmark to validate against. (ii) *Self-consistent background recomputation*: a custom sound-horizon/comoving-distance integrator, validated to $< 0.002\%$ against CLASS’s own derived parameters at $\epsilon = 0$, applied $\epsilon = 0.0251$ consistently to both integrals and found $H_0 \approx 75.8$ km/s/Mpc, **not** the ~ 72 km/s/Mpc quoted above — a real, unreconciled discrepancy that *overshoots* the SH0ES value rather than bridging Planck and SH0ES. The exact derivation behind the original ~ 72 estimate is not fully reproducible from what is documented. Full detail: `docs/cosmology/class_fork_validation.md`.
4. **Cosmic See-Saw / S_8 (updated 2026-07-11, Task T5.3):** the claim (Discussion, below) that a lighter late-time axion mass suppresses small-scale structure was tested quantitatively for the first time against this model’s own axion masses. Two independent real calculations both fail to produce the needed suppression: the standard fuzzy-dark-matter quantum-pressure mechanism (Hu, Barkana & Gruzinov 2000) gives negligible ($< 0.001\%$) σ_8 suppression at these masses ($1.8\text{--}3.2 \times 10^{-21}$ eV, $\sim 20\text{--}30\times$ heavier than the FDM mass range where this effect matters), and the ϵ -modified background growth-factor channel gives the *wrong sign* (a $+4\%$ *increase*, not a decrease). A rigorous JWST-side likelihood was not executed (would require halo-mass-function modeling with no independent benchmark); a real, independently-published qualitative check (Cox et al., arXiv:2307.10302) shows this model’s masses fall within the broad viable ALP window for JWST via a *different* mechanism, which is a necessary-condition check only. Full detail: `docs/cosmology/joint_ep`
5. **Chameleon Parameters:** To evade M87* superradiance bounds, the chameleon coupling must be set to $\gamma \approx 0.24$, corresponding to an unphysical potential index $n = -3$ in standard Khoury-Weltman theory [?]. **Update (2026-07-11, Task T3.2):** using the exact Dolan (2007) continued-fraction superradiance solver (validated to $< 0.4\%$ against published data), $S_{2,1}$ ’s *bare* coupling ($\alpha = 0.089$) survives M87* without any Chameleon screening at all; this unphysical-index tension now applies only to $S_{1,2}$, whose bare coupling ($\alpha = 0.155$) does not survive unscreened.

6. **Modularity Conjecture:** The connection between Order-3 Picard-Fuchs operators and K3 surfaces relies on the Stienstra-Beukers modularity correspondence [?], which is supported by extensive numerical checks but remains a conjecture in the formal Lean 4 pipeline. **Update (2026-07-11):** $S_{2,1}$ is now confirmed to satisfy an order-2 (elliptic), not order-3 (K3), recurrence — see `docs/gap1/ORDER_VERIFICATION_FINDINGS`. only $S_{1,2}$ remains a K3 candidate.

9 Discussion

The physical mechanism of Project Vafa-Continuity is distinct from standard Early Dark Energy (EDE) models. Rather than adding an ad-hoc scalar field that decays rapidly around recombination, our model uses the physical mass variation of the dark matter axion to reduce the sound horizon.

This mass-varying dark matter model introduces potential tensions with Lyman- α forest spectra [?, ?] and dark matter subhalo heating bounds [?, ?], as a lighter dark matter mass at late times could suppress small-scale structure. Future work must evaluate these constraints in detail to verify if the late-time mass of 1.83×10^{-21} eV is compatible with small-scale structure data.

10 Conclusion

The T^2 volume modulus expansion provides a qualitatively consistent dark energy trajectory and a directional shift toward higher H_0 , both motivated by the $K3 \times T^2$ string geometry. The model’s most significant positive result is the quantitative recovery of the Agrawal–Obied–Steinhardt–Vafa quintessence–Swampland tension [?] in a concrete geometric context: the best-fit $\lambda = 1.6724 > \sqrt{2}$ is formally certified to exclude attractor-phase dark energy (theorems `lambda_fit_exceeds_sqrt2` and `attractor_not_dark_energy`), and the resolution window $1 \lesssim \lambda < \sqrt{2}$ is precisely quantified. Three resolution programmes are outlined in §??. The framework is not yet a precision prediction: the best-fit w_0, w_a remains outside the DESI 2024 1σ contour, the H_0 estimate lacks a full Boltzmann treatment, and the general- n S_{20} recurrence remains an explicit `axiom` (kernel-verified for $n \leq 8$, exact-verified for $n \in [0, 60]$) pending a full Wilf–Zeilberger proof. These are explicitly open problems. The formal Lean 4 results provide a clean, machine-checked algebraic foundation on which more complete physics can be built.

Copyright (C) 2026 Xavier Callens (SocrateAI Scientific Agora). All rights reserved. No part of this work may be reproduced, distributed, or transmitted in any form or by any means without prior written permission from the author.

References

- [1] Prateek Agrawal, Georges Obied, Paul J Steinhardt, and Cumrun Vafa. Cosmological implications of the string swampland. *Physics Letters B*, 784:271–276, 2018.
- [2] Gert Almkvist, Christian van Enckevort, Duco van Straten, and Wadim Zudilin. Tables of Calabi-Yau equations. *arXiv preprint math/0507430*, 2005.

- [3] Asimina Arvanitaki, Savas Dimopoulos, Sergei Dubovsky, Nemanja Kaloper, and John March-Russell. String axiverse. *Physical Review D*, 81(12):123530, 2010.
- [4] Asimina Arvanitaki, Savas Dimopoulos, Marios Galanis, Luis Lehner, Joshua O Nee, and Vasileios Sherbrooke. Black hole superradiance signatures of ultralight vectors. *Physical Review D*, 95(4):043001, 2017.
- [5] Ana Bonaca, David W Hogg, Adrian M Price-Whelan, and Charlie Conroy. High-resolution spectroscopy of the gd-1 stellar stream. *The Astrophysical Journal*, 880(1):38, 2019.
- [6] Event Horizon Telescope Collaboration. First m87 event horizon telescope results. i. the shadow of the supermassive black hole. *The Astrophysical Journal Letters*, 875(1):L1, 2019.
- [7] Joseph P Conlon and Fernando Quevedo. Kähler moduli stabilisation and the gravitino mass problem. *Journal of High Energy Physics*, 2006(01):146, 2006.
- [8] Edmund J Copeland, M Sami, and Shinji Tsujikawa. Dynamics of dark energy. *International Journal of Modern Physics D*, 15(11):1753–1935, 2006.
- [9] Yuzhu Cui, Kazuhiro Hada, Tomohisa Kawashima, Motoki Kino, Kotaro Niinuma, et al. Precessing jet nozzle connecting to a spinning black hole in M87. *Nature*, 621:711–714, 2023.
- [10] DESI Collaboration, A G Adame, J Aguilar, S Ahlen, S Alam, D M Alexander, et al. DESI 2024 VI: Cosmological constraints from the measurements of baryon acoustic oscillations. *arXiv preprint*, 2024.
- [11] Steven Detweiler. Klein-gordon equation and rotating black holes. *Physical Review D*, 22(10):2323, 1980.
- [12] Sam R Dolan. Instability of the massive Klein-Gordon field on the Kerr spacetime. *Physical Review D*, 76(8):084001, 2007.
- [13] Vid Iršič, Matteo Viel, Martin G Haehnelt, James S Bolton, and George D Becker. First constraints on fuzzy dark matter from Lyman- α forest data and hydrodynamical simulations. *Physical Review Letters*, 119(3):031302, 2017.
- [14] Justin Khoury and Amanda Weltman. Chameleon cosmology. *Physical Review D*, 69(4):044026, 2004.
- [15] Justin Khoury and Amanda Weltman. Chameleon fields: awaiting surprises for tests of gravity in space. *Physical Review Letters*, 93(17):171104, 2004.
- [16] Federico Lelli, Stacy S McGaugh, and James M Schombert. Sparc: mass models for 175 disk galaxies with spitzer photometry and accurate rotation curves. *The Astronomical Journal*, 152(6):157, 2016.
- [17] David J E Marsh. Axion cosmology. *Physics Reports*, 643:1–79, 2016.
- [18] The mathlib Community. The lean mathematical library. pages 367–381, 2020.

- [19] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021.
- [20] Georges Obied, Hiroshi Ooguri, Lev Spodyneiko, and Cumrun Vafa. De sitter space and the swampland. *arXiv preprint hep-th/1806.08362*, 2018.
- [21] Hiroshi Ooguri, Eran Palti, Gary Shiu, and Cumrun Vafa. Distance and de sitter conjectures on the swampland. *Physics Letters B*, 788:180–184, 2019.
- [22] Hiroshi Ooguri and Cumrun Vafa. Geometry of the string landscape and the swampland. *Nuclear Physics B*, 766:21–33, 2007.
- [23] Planck Collaboration, N Aghanim, Y Akrami, M Ashdown, J Aumont, C Baccigalupi, M Ballardini, et al. Planck 2018 results. VI. cosmological parameters. *Astronomy & Astrophysics*, 641:A6, 2020.
- [24] Keir K Rogers and Hiranya V Peiris. Fuzzy dark matter and the dark energy survey year 1 data. *Physical Review Letters*, 126(7):071302, 2021.
- [25] Hsi-Yu Schive, Tzihong Chiueh, and Tom Broadhurst. Cosmic structure as the quantum interference of a coherent dark wave. *Nature Physics*, 10(7):496–499, 2014.
- [26] Jan Stienstra and Frits Beukers. Picard-fuchs equations and mirror maps for hypersurfaces. *Essays on Mirror Manifolds*, 1992.
- [27] Peter Svrcek and Edward Witten. Axions in string theory. *Journal of High Energy Physics*, 2006(06):051, 2006.
- [28] Jeremy J Webb and Jo Bovy. The effect of dark matter substructure on the gaps and morphology of stellar streams. *Monthly Notices of the Royal Astronomical Society*, 485(4):5929–5938, 2019.
- [29] Don Zagier. Integral solutions of Apéry-like recurrence equations. *Groups and Symmetries: From Neolithic Scots to John McKay*, 47:349–366, 2009.