

1 The epidemic lambda-coalescent model

2 Xavier Didelot^{1,2,*}, Ian Roberts², ...

3 ¹ School of Life Sciences, University of Warwick, United Kingdom

4

5 ² Department of Statistics, University of Warwick, United Kingdom

6

7 * Corresponding author. Tel: 0044 (0)2476 572827. Email: xavier.didelot@gmail.com

8 Running title: Epidemic lambda-coalescent

9 Keywords: lambda-coalescent model; pathogen phylogenetics; multiple mergers; offspring distribution

1 Introduction

Superspreading in infectious disease epidemiology (Lloyd-Smith et al. 2005). For example SARS-CoV-2 superspreading (Wang et al. 2020; Lemieux et al. 2021; Gómez-Carballa et al. 2021). Coalescent model (Kingman 1982a,b). Work by Li and Fraser (Li et al. 2017; Fraser and Li 2017). Lambda-coalescent models (Pitman 1999; Sagitov 1999; Donnelly and Kurtz 1999). Beta-coalescent (Schweinsberg 2003) is a specific type of Lambda-coalescent. Was used in (Hoscheit and Pybus 2019) and (Menardo et al. 2021). David's paper (Helekal et al. 2024).

2 Coalescence probabilities

2.1 General case

Discrete time t . Non-overlapping generations of infected individuals. At time t there are N_t infected individuals. Each of them creates a number $k_{t,i}$ of secondary infections at time $t + 1$, following the offspring distribution $\alpha_t(k)$. The mean of this distribution is the basic reproduction number R_t and the variance is v_t . We have:

$$N_{t+1} = \sum_{i=1}^{N_t} k_{t,i} \quad (1)$$

Let $p_{n,t}$ be the probability that n individuals at time $t + 1$ have the same infector at time t .

2.2 Poisson case

Here the offspring distribution is $\text{Poisson}(R_t)$.

The probability of coalescence for two lines is:

$$p_{2,t} = \frac{1}{N_t} \quad (2)$$

The probability of coalescence for n lines is:

$$p_{n,t} = \frac{1}{N_t^{n-1}} \quad (3)$$




Figure 1: (A) Offspring distribution. (B) Probability of coalescence.

2.3 NegBin case

Here the offspring distribution is Negative-Binomial with mean R_t and variance v_t . The parameters of this distribution are $r = R_t^2/(v_t - R_t)$ and $p = R_t/v_t$.

The probability of coalescence for two lines is:

$$p_{2,t} = \frac{r + 1}{N_t r + 1} \quad (4)$$

The probability of coalescence for n lines is:

$$p_{n,t} = \frac{\prod_{i=1}^{n-1} r + i}{\prod_{i=1}^{n-1} N_t r + i} \quad (5)$$

2.4 Example

See Figure 1.

35 **3 Implementation**

36 We implemented the analytical methods described in this paper in a new R package entitled *EpiLambda*
37 which is available at <https://github.com/xavierdidelot/EpiLambda> for R version 3.5 or later. All
38 code and data needed to replicate the results are included in the “run” directory of the *EpiLambda*
39 repository.

40 **4 Discussion**

41 **Acknowledgements**

42 We acknowledge funding from the National Institute for Health Research (NIHR) Health Protection
43 Research Unit in Genomics and Enabling Data.

References

- Donnelly, P., Kurtz, T.G., 1999. Particle Representations for Measure-Valued Population Models. *The Annals of Probability* 27. doi:10.1214/aop/1022677258.
- Fraser, C., Li, L.M., 2017. Coalescent models for populations with time-varying population sizes and arbitrary offspring distributions. *bioRxiv* , 10.1101/131730doi:10.1101/131730.
- Gómez-Carballa, A., Pardo-Seco, J., Bello, X., Martínón-Torres, F., Salas, A., 2021. Superspreading in the emergence of COVID-19 variants. *Trends in Genetics* 37, 1069–1080. doi:10.1016/j.tig.2021.09.003.
- Helekal, D., Koskela, J., Didelot, X., 2024. Inference of multiple mergers while dating a pathogen phylogeny. *bioRxiv* , 2023.09.12.557403doi:10.1101/2023.09.12.557403.
- Hoscheit, P., Pybus, O.G., 2019. The multifurcating skyline plot. *Virus Evolution* 5, 1–10. doi:10.1093/ve/vez031.
- Kingman, J., 1982a. The coalescent. *Stochastic Processes and their Applications* 13, 235–248. doi:10.1016/0304-4149(82)90011-4.
- Kingman, J.F.C., 1982b. On the genealogy of large populations. *Journal of Applied Probability* 19, 27–43. doi:10.2307/3213548.
- Lemieux, J.E., Siddle, K.J., Shaw, B.M., Loreth, C., Schaffner, S.F., Gladden-Young, A., Adams, G., Fink, T., Tomkins-Tinch, C.H., Krasilnikova, L.A., DeRuff, K.C., Rudy, M., Bauer, M.R., Lagerborg, K.A., Normandin, E., Chapman, S.B., Reilly, S.K., Anahtar, M.N., Lin, A.E., Carter, A., Myhrvold, C., Kembell, M.E., Chaluvadi, S., Cusick, C., Flowers, K., Neumann, A., Cerrato, F., Farhat, M., Slater, D., Harris, J.B., Branda, J.A., Hooper, D., Gaeta, J.M., Baggett, T.P., O’Connell, J., Gnirke, A., Lieberman, T.D., Philippakis, A., Burns, M., Brown, C.M., Luban, J., Ryan, E.T., Turbett, S.E., LaRocque, R.C., Hanage, W.P., Gallagher, G.R., Madoff, L.C., Smole, S., Pierce, V.M., Rosenberg, E., Sabeti, P.C., Park, D.J., MacInnis, B.L., 2021. Phylogenetic analysis of SARS-CoV-2 in Boston highlights the impact of superspreading events. *Science* 371, eabe3261. doi:10.1126/science.abe3261.
- Li, L.M., Grassly, N.C., Fraser, C., 2017. Quantifying Transmission Heterogeneity Using Both Pathogen Phylogenies and Incidence Time Series. *Molecular Biology and Evolution* 34, 2982–2995. doi:10.1093/molbev/msx195.
- Lloyd-Smith, J., Schreiber, S., Kopp, P., Getz, W., 2005. Superspreading and the effect of individual variation on disease emergence. *Nature* 438, 355–9. doi:10.1038/nature04153.
- Menardo, F., Gagneux, S., Freund, F., 2021. Multiple Merger Genealogies in Outbreaks of *Mycobacterium tuberculosis*. *Molecular Biology and Evolution* 38, 290–306. doi:10.1093/molbev/msaa179.
- Pitman, J., 1999. Coalescents with multiple collisions. *The Annals of Probability* 27, 1870–1902.
- Sagitov, S., 1999. The general coalescent with asynchronous mergers of ancestral lines. *Journal of Applied Probability* 36, 1116–1125. doi:10.1239/jap/1032374759.
- Schweinsberg, J., 2003. Coalescent processes obtained from supercritical Galton–Watson processes. *Stochastic Processes and their Applications* 106, 107–139. doi:10.1016/S0304-4149(03)00028-0.
- Wang, L., Didelot, X., Yang, J., Wong, G., Shi, Y., Liu, W., Gao, G.F., Bi, Y., 2020. Inference of person-to-person transmission of COVID-19 reveals hidden super-spreading events during the early outbreak phase. *Nature Communications* 11, 5006. doi:10.1038/s41467-020-18836-4.