

1 Ancestral process for infectious disease outbreaks with superspreading

2 Xavier Didelot^{1,2,*}, David Helekal³, Ian Roberts²

3 ¹ School of Life Sciences, University of Warwick, Coventry, United Kingdom

4 ² Department of Statistics, University of Warwick, Coventry, United Kingdom

5 ³ Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of Public Health,
6 Boston, Massachusetts, USA

7 * Corresponding author. Tel: 0044 (0)2476 572827. Email: `xavier.didelot@warwick.ac.uk`

8 Running title: Ancestry for outbreaks with superspreading

9 Keywords: infectious disease epidemiology modelling; offspring distribution; superspreading;
10 outbreaks; Lambda-coalescent model; multiple mergers

Abstract

When an infectious disease outbreak is of a relatively small size, describing the ancestry of a sample of infected individuals is difficult because most ancestral models assume large population sizes. Given a set of infected individuals, we show that it is possible to express exactly the probability that they have the same infector, either inclusively (so that other individuals may have the same infector too) or exclusively (so that they may not). To compute these probabilities requires knowledge of the offspring distribution, which determines how many infections each infected individual causes. We consider transmission both without and with superspreading, in the form of a Poisson and a Negative-Binomial offspring distribution, respectively. We show how our results can be incorporated into a new Lambda-coalescent model which allows multiple lineages to coalesce together. We call this new model the Omega-coalescent, we compare it with previously proposed alternatives, and advocate its use in future studies of infectious disease outbreaks.

1 Introduction

An outbreak of an infectious disease typically starts when a single or a small number of infected individuals appear within a susceptible population. Each infected individual may come in contact with and transmit the disease to each of the susceptible individuals, who will then become infected in their turn and spread the disease further. Most mathematical models of infectious diseases describe situations where the disease is at an equilibrium, when the number of infected individuals is high and/or with a significant part of the population already infected (Anderson and May 1991; Keeling and Rohani 2008). Here however we focus on the early stages of an epidemic, where the number of infected individuals is small and the number of susceptibles comparatively high and constant. In this situation it is useful to consider the number of new infections that each infected individual is likely to cause, and the probabilistic distribution for this number is often called the offspring distribution (Grassly and Fraser 2008). The mean of the offspring distribution is called the basic reproduction number R_0 and has been given much attention especially since it determines how likely the outbreak is to spread, and how much effort would be needed to bring it under control (Fraser et al. 2004; Ferguson et al. 2006).

If we consider that all individuals are infectious for the same duration and with the same transmission rate, the offspring distribution is Poisson distributed with mean R_0 , in which case the variance of the offspring distribution is also R_0 . We would then say that there is no transmission heterogeneity. However, in practice there are many reasons why this may not be the case, with some individuals being infectious for longer than others, or being more infectious than others, or having more frequent contacts with susceptibles, or being less symptomatic and therefore less likely to reduce contact numbers, etc. All these factors cause the offspring distribution to be more dispersed than it would otherwise be, that is to have a variance greater than its mean R_0 . A frequent choice to capture this overdispersion is to model the offspring distribution using a Negative-Binomial distribution with mean R_0 and dispersion parameter r (Lloyd-Smith et al. 2005; Grassly and Fraser 2008). When r is close to zero the variance is high compared to the mean, whereas when r is high the variance becomes close to the mean. This transmission heterogeneity is often called superspreading, although this is perhaps misleading as it is the rule rather than the exception of how infectious diseases spread. Superspreading has indeed been described in many diseases (Woolhouse et al. 1997; Stein 2011; Kucharski and Althaus 2015; Wang et al. 2021), and most recently for SARS-CoV-2 (Wang et al. 2020; Lemieux et al. 2021; Gómez-Carballa et al. 2021; Du et al. 2022).

As an outbreak unfolds forward-in-time, a transmission tree is generated representing who-infected-

whom, in which each node is an infected individual and points towards a number of nodes distributed according to the offspring distribution. Here we consider the reverse problem of the transmission ancestry, going backward-in-time, from a sample of infected individuals, until reaching the last common transmission ancestor of the whole sample. Given a set of n sampled individuals, we show how to calculate the probability that a given subset of size k have the same infector, either inclusively (so that the remaining $n - k$ may also have the same infector or not) or exclusively (so that none of the remaining $n - k$ have the same infector). We start by considering the general case of an offspring distribution with arbitrary form, and then the specific cases of offspring distributions that follow a Poisson and a Negative-Binomial distribution. The main novelty of our approach is that we consider that the overall population size is small, but we show that in the limit where the population size is large, our results agree with several previous studies (Volz 2012; Koelle and Rasmussen 2012; Fraser and Li 2017). Finally, we show how our results can be incorporated into a new Lambda-coalescent model (Pitman 1999; Sagitov 1999; Donnelly and Kurtz 1999) and compare it with previously proposed models.

2 General offspring distribution case

Let time be measured in discrete units and denoted t . Each discrete value of t corresponds to a unique non-overlapping generation of infected individuals, so that individuals infected at t have offspring at $t + 1$, etc. Let N_t denote the number of infectious individuals at time t . Each of them creates a number $s_{t,i}$ of secondary infections at time $t + 1$, following the offspring distribution $\alpha_t(s)$. The mean of this distribution is the basic reproduction number R_t and the variance is V_t . The total number of infected individuals at time $t + 1$ is given by:

$$N_{t+1} = \sum_{i=1}^{N_t} s_{t,i} \quad (1)$$

2.1 Inclusive coalescence probability

We define the inclusive coalescence probability $p_{k,t}(N_t, N_{t+1})$ as the probability that a specific set of k individuals from generation $t + 1$ have the same infector in generation t , conditional on population sizes N_t and N_{t+1} . Given full information about offspring counts from individuals in generation t ,

80 $\mathbf{s}_t = (s_{t,1}, \dots, s_{t,N_t})$, we have:

$$\begin{aligned}
 p_{k,t}(\mathbf{s}_t, N_t) &= \sum_{i=1}^{N_t} \frac{\binom{s_{t,i}}{k}}{\binom{N_{t+1}}{k}} \\
 &= \sum_{i=1}^{N_t} \frac{s_{t,i}!}{(s_{t,i} - k)!} \frac{(N_{t+1} - k)!}{N_{t+1}!}
 \end{aligned} \tag{2}$$

81 Full information $\{s_{t,i}\}$ yields the population size N_{t+1} as shown in Equation 1, but this is not available
 82 in practice. We can instead express the inclusive coalescence probability conditioning on the next
 83 population size N_{t+1} by summing over possible offspring counts $\mathbf{s}_t = (s_{t,1}, \dots, s_{t,N_t})$ conditional on the
 84 total generation size. Let $S_t^{-(1)} = (S_{t,2}, \dots, S_{t,N_t})$:

$$\begin{aligned}
 p_{k,t}(N_t, N_{t+1}) &= \sum_{\mathbf{s}_t \in \mathbb{N}_0^{N_t}} \mathbb{P} \left[\mathbf{S}_t = \mathbf{s}_t \middle| \sum_{i=1}^{N_t} S_{t,i} = N_{t+1} \right] p_{k,t}(\mathbf{s}_t, N_t) \\
 &= \sum_{\mathbf{s}_t \in \mathbb{N}_0^{N_t}} \mathbb{P} \left[\mathbf{S}_t = \mathbf{s}_t \middle| \sum_{i=1}^{N_t} S_{t,i} = N_{t+1} \right] \sum_{i=1}^{N_t} \frac{\binom{s_{t,i}}{k}}{\binom{N_{t+1}}{k}} \\
 &= \sum_{i=1}^{N_t} \sum_{\mathbf{s}_t \in \mathbb{N}_0^{N_t}} \frac{\binom{s_{t,i}}{k}}{\binom{N_{t+1}}{k}} \mathbb{P} \left[S_{t,1} = s_{t,1}, \mathbf{S}_t^{-(1)} = \mathbf{s}_t^{-(1)} \middle| \sum_{i=1}^{N_t} S_{t,i} = N_{t+1} \right] \\
 &= \frac{N_t}{\binom{N_{t+1}}{k}} \sum_{\mathbf{s}_t \in \mathbb{N}_0^{N_t}} \binom{s_{t,1}}{k} \mathbb{P} \left[S_{t,1} = s_{t,1} \middle| \sum_{i=1}^{N_t} S_{t,i} = N_{t+1} \right] \\
 &\quad \times \mathbb{P} \left[\mathbf{S}_t^{-(1)} = \mathbf{s}_t^{-(1)} \middle| S_{t,1} = s_{t,1}, \sum_{i=1}^{N_t} S_{t,i} = N_{t+1} \right] \\
 &= \frac{N_t}{\binom{N_{t+1}}{k}} \sum_{s_{t,1}=0}^{N_{t+1}} \binom{s_{t,1}}{k} \mathbb{P} \left[S_{t,1} = s_{t,1} \middle| \sum_{i=1}^{N_t} S_{t,i} = N_{t+1} \right] \\
 &\quad \times \underbrace{\sum_{\mathbf{s}_t^{-(1)} \in \mathbb{N}_0^{N_t-1}} \mathbb{P} \left[\mathbf{S}_t^{-(1)} = \mathbf{s}_t^{-(1)} \middle| \sum_{i=2}^{N_t} S_{t,i} = N_{t+1} - s_{t,1} \right]}_{=1} \\
 &= \frac{N_t}{\binom{N_{t+1}}{k}} \mathbb{E} \left[\binom{S_{t,1}}{k} \middle| \sum_{i=1}^{N_t} S_{t,i} = N_{t+1} \right] \\
 &= N_t \frac{(N_{t+1} - k)!}{N_{t+1}!} \mathbb{E} \left[\frac{S_{t,1}!}{(S_{t,1} - k)!} \middle| \sum_{i=1}^{N_t} S_{t,i} = N_{t+1} \right]
 \end{aligned} \tag{3}$$

85 The k -th falling factorial moments $\mathbb{E}\left[\frac{S_{t,1}!}{(S_{t,1}-k)!} \mid \sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right]$ in Equation 3 can be readily obtained
 86 by differentiating the probability generating function of $S_{t,1} \mid (\sum_{i=1}^{N_t} S_{t,i} = N_{t+1})$.

87 2.2 Exclusive coalescence probability

88 Generally, we observe a sample of individuals from each generation rather than the entire population.
 89 In this case, we are interested in the exclusive coalescence probability $p_{n,k,t}(N_t, N_{t+1})$ that a specific
 90 subset of k individuals amongst n sampled individuals arose from a common infector one generation
 91 in the past given knowledge of the total population sizes N_t and N_{t+1} . Let us first assume full
 92 knowledge about offspring counts of the individuals at time t amongst the sample at time $t+1$,
 93 namely $\mathbf{x}_t = (x_{t,1}, \dots, x_{t,N_t})$ such that $x_{t,1} + \dots + x_{t,N_t} = n$. Note that $X_{t,i}$ does not follow the same
 94 offspring distribution as $S_{t,i}$. We have:

$$\begin{aligned} p_{n,k,t}(\mathbf{X}_t = \mathbf{x}_t, N_t) &= \sum_{i=1}^{N_t} \frac{\binom{x_{t,i}}{k}}{\binom{n}{k}} \mathbb{I}\{x_{t,i} = k\} \\ &= \sum_{i=1}^{N_t} \frac{x_{t,i}!}{(x_{t,i}-k)!} \frac{(n-k)!}{n!} \mathbb{I}\{x_{t,i} = k\} \end{aligned} \quad (4)$$

95 Similarly to the inclusive coalescence probability in Equation 3, we can use this to evaluate the exclusive
 96 probability given N_t and N_{t+1} by summing over possible parent offspring configurations (for $k \leq n$):

$$\begin{aligned} p_{n,k,t}(N_t, N_{t+1}) &= \sum_{\mathbf{x}_t \in \mathbb{N}_0^{N_t}} \mathbb{P}\left[\mathbf{X}_t = \mathbf{x}_t \mid \sum_{i=1}^n X_{t,i} = n\right] p_{n,k,t}(\mathbf{x}_t, N_t) \\ &= \sum_{\mathbf{x}_t \in \mathbb{N}_0^{N_t}} \mathbb{P}\left[\mathbf{X}_t = \mathbf{x}_t \mid \sum_{i=1}^n X_{t,i} = n\right] \sum_{i=1}^{N_t} \frac{\binom{x_{t,i}}{k}}{\binom{n}{k}} \mathbb{I}\{x_{t,i} = k\} \\ &= \frac{N_t}{\binom{n}{k}} \sum_{\mathbf{x}_t \in \mathbb{N}_0^{N_t}} \binom{x_{t,1}}{k} \mathbb{P}\left[\mathbf{X}_t = \mathbf{x}_t \mid \sum_{i=1}^{N_t} X_{t,i} = n\right] \mathbb{I}\{x_{t,1} = k\} \\ &= \frac{N_t}{\binom{n}{k}} \sum_{\mathbf{x}_t^{-(1)} \in \mathbb{N}_0^{N_t-1}} \binom{k}{k} \mathbb{P}\left[X_{t,1} = k, \mathbf{X}_t^{-(1)} = \mathbf{x}_t^{-(1)} \mid \sum_{i=1}^{N_t} X_{t,i} = n\right] \\ &= \frac{N_t}{\binom{n}{k}} \mathbb{P}[X_{t,1} = k \mid \sum_{i=1}^{N_t} X_{t,i} = n] \underbrace{\sum_{\mathbf{x}_t^{-(1)} \in \mathbb{N}_0^{N_t-1}} \mathbb{P}\left[\mathbf{X}_t^{-(1)} = \mathbf{x}_t^{-(1)} \mid \sum_{i=1}^{N_t} X_{t,i} = n, X_{t,1} = k\right]}_{=1} \end{aligned}$$

$$= \frac{N_t}{\binom{n}{k}} \mathbb{P} \left[X_{t,1} = k \middle| \sum_{i=1}^{N_t} X_{t,i} = n \right] \quad (5)$$

97 If we consider one of the lines observed amongst a set of n , it can either remain uncoalesced with
 98 probability $p_{n,1,t}(N_t, N_{t+1})$ or coalesce in an event of size k with probability $p_{n,k,t}(N_t, N_{t+1})$ with any
 99 set of $k - 1$ lines among the $n - 1$ other lines, leading to the following complementarity equation:

$$\sum_{k=1}^n \binom{n-1}{k-1} p_{n,k,t}(N_t, N_{t+1}) = 1 \quad (6)$$

100 We can show that it is indeed satisfied by the formula in Equation 5, thus providing a sanity check on
 101 this formula:

$$\begin{aligned} \sum_{k=1}^n \binom{n-1}{k-1} p_{n,k,t}(N_t, N_{t+1}) &= \sum_{k=1}^n \binom{n-1}{k-1} \frac{N_t}{\binom{n}{k}} \mathbb{P} \left[X_1 = k \middle| \sum_{i=1}^{N_t} X_i = n \right] \\ &= \sum_{k=1}^n N_t \frac{k}{n} \mathbb{P} \left[X_1 = k \middle| \sum_{i=1}^{N_t} X_i = n \right] \\ &= \frac{N_t}{n} \sum_{k=0}^n k \mathbb{P} \left[X_1 = k \middle| \sum_{i=1}^{N_t} X_i = n \right] \\ &= \frac{N_t}{n} \mathbb{E} \left[X_1 \middle| \sum_{i=1}^{N_t} X_i = n \right] \\ &= \frac{1}{n} \sum_{i=1}^{N_t} \mathbb{E} \left[X_i \middle| \sum_{i=1}^{N_t} X_i = n \right] \\ &= \frac{1}{n} \mathbb{E} \left[\sum_{i=1}^{N_t} X_i \middle| \sum_{i=1}^{N_t} X_i = n \right] \\ &= 1 \end{aligned} \quad (7)$$

3 Poisson offspring distribution case

In this section we consider that the offspring distribution is $\alpha_t = \text{Poisson}(R_t)$. In this case, we have:

$$\sum_{i=1}^{N_t} S_{t,i} \sim \text{Poisson}(N_t R_t) \quad (8)$$

and the conditional distribution:

$$\begin{aligned} \mathbb{P}\left[S_{t,1} = s \mid \sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right] &= \frac{\mathbb{P}\left[S_{t,1} = s, \sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right]}{\mathbb{P}\left[\sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right]} \\ &= \frac{\alpha_t(s) \mathbb{P}\left[\sum_{i=2}^{N_t} S_{t,i} = N_{t+1} - s\right]}{\mathbb{P}\left[\sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right]} \\ &= \frac{\frac{R_t^s e^{-R_t}}{s!} \cdot \frac{((N_t - 1)R_t)^{N_{t+1} - s}}{(N_{t+1} - s)!}}{\frac{(N_t R_t)^{N_{t+1}} e^{-N_t R_t}}{N_{t+1}!}} \\ &= \binom{N_{t+1}}{s} \left(\frac{1}{N_t}\right)^s \left(1 - \frac{1}{N_t}\right)^{N_{t+1} - s} \end{aligned} \quad (9)$$

This is the probability mass function of a Binomial distribution and therefore we deduce that:

$$S_{t,1} \mid \left(\sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right) \sim \text{Binomial}\left(N_{t+1}, \frac{1}{N_t}\right) \quad (10)$$

The k -th falling factorial moments of $X \sim \text{Binomial}(n, p)$ are (Potts 1953):

$$\mathbb{E}\left[\frac{X!}{(X - k)!}\right] = \binom{n}{k} p^k k! \quad (11)$$

By applying this formula to the Binomial distribution in Equation 10 and injecting into Equation 3, we deduce that the inclusive probability of coalescence for k lines is:

$$p_{k,t}(N_t, N_{t+1}) = \frac{1}{N_t^{k-1}} \quad (12)$$

109 In addition, following a similar reasoning as for Equation 10 we can show that:

$$X_{t,1} \left| \left(\sum_{i=1}^{N_t} X_{t,i} = n \right) \sim \text{Binomial} \left(n, \frac{1}{N_t} \right) \quad (13)$$

110 By injecting the probability mass function of this Binomial distribution into Equation 5 we deduce
 111 that the exclusive probability of coalescence for k lines from a sample of n ($n \geq k$) is:

$$p_{n,k,t}(N_t, N_{t+1}) = \frac{(N_t - 1)^{n-k}}{N_t^{n-1}} \quad (14)$$

112 The definitions of the inclusive and exclusive coalescence probabilities imply that the former is a special
 113 case of the latter, with specifically $p_{k,t}(N_t, N_{t+1}) = p_{k,k,t}(N_t, N_{t+1})$ and this is clearly verified in
 114 Equations 12 and 14. It is interesting to note that neither the inclusive nor the exclusive coalescence
 115 probability depend on the mean R_t of the Poisson offspring distribution or the size N_{t+1} of the
 116 population at time $t + 1$. Both only depend on the population size N_t at time t . The inclusive
 117 coalescent probability in Equation 12 can also be obtained conceptually by considering that among
 118 the k lines, the first one has an ancestor with probability one, and the remaining $k - 1$ need to have the
 119 same ancestor among a set of N_t from which they choose uniformly at random so that the probability of
 120 picking the same ancestor is $1/N_t$. The exclusive coalescent probability in Equation 14 can be derived
 121 likewise by considering that in addition to the above, each of the $n - k$ other lines need to choose a
 122 different ancestor, which happens with probability $(N_t - 1)/N_t$. Figure 1 illustrates the inclusive and
 123 exclusive coalescence probabilities for a set of size $k = 1$ to $k = 10$ amongst a total of $n = 10$ observed
 124 individuals, in a population of size $N_t = 10$, $N_t = 20$ or $N_t = 30$.

125 4 Negative-Binomial offspring distribution case

126 In this section we consider that the offspring distribution is Negative-Binomial, a distribution often
 127 used to model superspreading individuals (Lloyd-Smith et al. 2005) and which can also be used to model
 128 superspreading events (Craddock et al. 2025). Let $\alpha_t = \text{Negative-Binomial}(r, p)$ with parameters (r, p)
 129 set by moment-matching the mean R_t and variance V_t of the offspring distribution which are assumed




Figure 1: Inclusive and exclusive coalescence probabilities for the Poisson case.

constant over time. The resulting parameters for this distribution are $r = R_t^2/(V_t - R_t)$ and $p = R_t/V_t$.

In this case, we have:

$$\sum_{i=1}^{N_t} S_{t,i} \sim \text{Negative-Binomial}(N_t r, p) \quad (15)$$

and similarly to the Poisson offspring distribution case we identify that the conditional distribution of

$S_{t,1} | \sum_{i=1}^{N_t} S_{t,i}$ is as follows:

$$\begin{aligned} \mathbb{P}\left[S_{t,1} = s \mid \sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right] &= \frac{\alpha_t(s) \cdot \mathbb{P}\left[\sum_{i=2}^{N_t} S_{t,i} = N_{t+1} - s\right]}{\mathbb{P}\left[\sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right]} \\ &= \frac{\frac{\Gamma(r+s)}{s! \Gamma(r)} (1-p)^s p^r \cdot \frac{\Gamma((N_t-1)r + (N_{t+1}-s))}{(N_{t+1}-s)! \Gamma((N_t-1)r)} (1-p)^{N_{t+1}-s} p^{(N_t-1)r}}{\frac{\Gamma(N_t r + N_{t+1})}{N_{t+1}! \Gamma(N_t r)} (1-p)^{N_{t+1}} p^{N_t r}} \\ &= \frac{N_{t+1}!}{s! (N_{t+1}-s)!} \frac{\Gamma(r+s) \Gamma((N_t-1)r + (N_{t+1}-s))}{\Gamma(N_t r + N_{t+1})} \frac{\Gamma(N_t r)}{\Gamma(r) \Gamma((N_t-1)r)} \\ &= \binom{N_{t+1}}{s} \frac{B(s+r, N_{t+1}-s+(N_t-1)r)}{B(r, (N_t-1)r)} \end{aligned} \quad (16)$$

where $B(x, y)$ denotes the Beta function defined as $B(x, y) = \Gamma(x)\Gamma(y)/\Gamma(x+y) = \int t^{x-1}(1-t)^{y-1} dt$.

This is the probability mass function of a Beta-Binomial distribution and therefore we deduce that:

$$S_{t,1} \mid \left(\sum_{i=1}^{N_t} S_{t,i} = N_{t+1}\right) \sim \text{Beta-Binomial}(N_{t+1}, r, (N_t-1)r) \quad (17)$$

The k -th falling factorial moments of $X \sim \text{Beta-Binomial}(n, \alpha, \beta)$ are (Tripathi et al. 1994):

$$\mathbb{E}\left[\frac{X!}{(X-k)!}\right] = \binom{n}{k} \frac{B(\alpha+k, \beta)k!}{B(\alpha, \beta)} \quad (18)$$

By applying this formula to the Beta-Binomial distribution in Equation 17 and injecting into Equation

3, we deduce that the inclusive probability of coalescence for k lines is:

$$p_{k,t}(N_t, N_{t+1}) = \frac{B(N_t r + 1, r + k)}{B(r + 1, N_t r + k)} \quad (19)$$

139 In addition, following a similar reasoning as for Equation 17, we can show that:

$$X_{t,1} \left| \left(\sum_{i=1}^{N_t} X_{t,i} = n \right) \right. \sim \text{Beta-Binomial}(n, r, (N_t - 1)r) \quad (20)$$

140 By injecting the probability mass function of this Beta-Binomial distribution into Equation 5 we deduce
141 that the exclusive probability of coalescence for k lines is:

$$p_{n,k,t}(N_t, N_{t+1}) = \frac{N_t B(k + r, n - k + N_t r - r)}{B(r, N_t r - r)} \quad (21)$$

142 As for the Poisson case we can check that $p_{k,t}(N_t, N_{t+1}) = p_{k,k,t}(N_t, N_{t+1})$ using the formulas in
143 Equations 19 and 21 and the definition of the Beta function. It is interesting to note that as for the
144 Poisson case, the inclusive and exclusive coalescence probabilities do not depend on the size N_{t+1} of
145 the population at time $t + 1$. They both depend on the Negative-Binomial offspring distribution only
146 through the dispersion parameter r . If we consider that r is large in Equations 19 and 21, we can
147 derive that the asymptotic behaviour is the same as in the Poisson case shown in Equations 12 and 14.
148 For example this can be derived by rewriting the Beta functions using Gamma functions, and using
149 the following form of Stirling's approximation:

$$\lim_{a \rightarrow \infty} \frac{\Gamma(a + b)}{\Gamma(a)} = a^b e^{-b} \quad (22)$$

150 Figure 2 illustrates the inclusive and exclusive coalescence probabilities for the Negative-Binomial case
151 for a set of size $k = 1$ to $k = 10$ amongst a total of $n = 10$ observed lines, in a population with
152 size $N_t = 20$. Several Negative-Binomial offspring distributions are compared, all of which have the
153 same mean $R_t = 2$, and with the dispersion parameter equal to $r = 0.1$, $r = 1$, $r = 10$ and $r = 100$
154 (Figure 2A). When $r = 1$ the Negative-Binomial reduces to a Geometric distribution. When r is high
155 the dispersion is low and the Negative-Binomial case behaves almost like the Poisson case for both
156 the inclusive (Figure 2B) and the exclusive coalescence probabilities (Figure 2C). When r is lower the
157 dispersion of the offspring distribution increases, so that both the inclusive and exclusive probabilities




Figure 2: (A) Offspring distributions with mean $R_t = 2$. (B) Inclusive probability of coalescence for $N_t = 20$ and $n = 10$. (C) Exclusive probability of coalescence for $N_t = 20$ and $n = 10$.

of larger multiple merger events are increased compared to the Poisson case. In particular, when $r = 0.1$ we see that the exclusive probability can increase with the size of the event considered (Figure 2C). This happens because the probability is not much lower for the common ancestor having say 10 rather than 9 offspring, while on the other hand if the event is of size 9 only then another individual in the generation of the ancestor needs to have had at least one sampled offspring.

5 Limit when the population size is large

Here we consider that the population size N_t is fixed and large, so that we can show the connections between our results and several previous studies on the ancestral process of infectious diseases. In the Poisson case, from Equations 12 and 14 we can see that both inclusive and exclusive probabilities are of order $\mathcal{O}(N_t^{1-k})$. We can therefore ignore events with $k > 2$ and retain only the events with $k = 2$, which means that there are only binary coalescent events and no multiple merger events. The binary coalescent events occur with the same inclusive and exclusive probabilities:

$$p_{2,t}(N_t, N_{t+1}) = p_{n,2,t}(N_t, N_{t+1}) = \frac{1}{N_t} \quad (23)$$

For the Negative-Binomial case, from Equations 19 and 21 we can rewrite using Gamma functions and apply the form of Stirling's equation given in Equation 22 to show that once again both inclusive and exclusive probabilities are also of order $\mathcal{O}(N_t^{1-k})$. We can therefore once again ignore events with $k > 2$ and retain only the events with $k = 2$ which occur with the same inclusive and exclusive probabilities:

$$p_{2,t}(N_t, N_{t+1}) = p_{n,2,t}(N_t, N_{t+1}) = \frac{r+1}{N_t r + 1} \approx \frac{r+1}{N_t r} \quad (24)$$

Koelle and Rasmussen (2012) derived the rates of coalescence of two lineages for several epidemiological models, assuming a large population at equilibrium. For each model they use the equation $N_e = N/\sigma^2$ to relate the effective population size N_e to the actual population size N and the variance σ^2 in the number of offspring. This relationship was first established by Kingman (1982a) to derive the backward-in-time coalescent model from the forward-in-time Cannings exchangeable models (Cannings 1974). This result implies that the rate of coalescence for two lineages is $1/N_e = \sigma^2/N$. From Equation 24 we can take $R_t = 1$ to achieve equilibrium of the population size and the method of moments estimator $r = R_t^2/(V_t - R_t) = 1/(V_t - 1)$ to deduce the equivalent result $p_{2,t}(N_t, N_{t+1}) = V_t/N_t$.

Volz (2012) showed that the rate of coalescence for two lineages under a continuous-time epidemic coalescent model is $2f(t)/I(t)^2$ where $f(t)$ is the incidence of the disease and $I(t)$ its prevalence. Setting in this formula the prevalence as $I(t) = N_{t+1} = N_t R_t$ and the incidence as $f(t) = R_t I(t) = R_t^2 N_t$ we get a coalescent rate of $2/N_t$. To apply our methodology we need to consider that the offspring distribution is Geometric, since the epidemiological models considered have successes (transmission) happening until the first failure (removal). We therefore set $r = 1$ in Equation 24 to make the Negative-Binomial offspring distribution reduce to a Geometric distribution and the same result follows.

Fraser and Li (2017) calculated the effective population size $N_e(t)$ as a function of the actual population size $N(t)$ and the mean and variance of the offspring distribution R and σ^2 . This formula was used to estimate the dispersion parameter of a Negative-Binomial offspring distribution from genetic data (Li et al. 2017). Using our notations, their formula is equivalent to the inclusive coalescence probability for two lineages:

$$p_{2,t}(N_t, N_{t+1}) = \frac{V_t/R_t + R_t - 1}{N_t R_t} \quad (25)$$

In the Poisson case we have $V_t = R_t$ so that Equation 25 simplifies to $1/N_t$ which agrees with Equation 23. In the Negative-Binomial case we have $V_t/R_t = 1/p = 1 + R_t/r$ so that Equation 25 simplifies to $(r+1)/(N_t r)$ which agrees with our Equation 24. Conversely, if we substitute the method of moments estimator $r = R_t^2/(V_t - R_t)$ in Equation 24 we obtain the Equation 25 originally from Fraser and Li (2017).

6 Definition of a new Lambda-coalescent model

The coalescent model (Kingman 1982a,b) describes the ancestry of a sample from a large population evolving according to many forward-in-time models such as the Wright-Fisher model (Wright 1931; Fisher 1930), the Moran model (Moran 1958) and the Cannings exchangeable model (Cannings 1974). Since the coalescent considers a large population in which each individual only has a number of offspring that is small compared to the population size, coalescent trees are always binary and do not feature multiple mergers, making them unsuitable to represent the ancestry of outbreaks considered in this study. However, if the population size is small in any of the aforementioned forward-in-time models, multiple mergers can occur, where more than two sampled individuals have the same ancestor. The

208 Lambda-coalescent model is an extension of the coalescent model that allows for such multiple merger
 209 events (Pitman 1999; Sagitov 1999; Donnelly and Kurtz 1999).

210 A Lambda-coalescent model is defined by a Lambda measure $\Lambda(dx)$ on the interval $[0, 1]$, from which
 211 we deduce the rate $\lambda_{n,k}$ at which any subset of k lineages within a set of n observed lineages coalesce:

$$\lambda_{n,k} = \int_0^1 x^{k-2} (1-x)^{n-k} \Lambda(dx) \quad (26)$$

212 The Beta-coalescent (Schweinsberg 2003) is a specific type of Lambda-coalescent that has been used
 213 recently in several studies analysing genetic data from infectious disease agents (Hoscheit and Pybus
 214 2019; Menardo et al. 2021; Helekal et al. 2025; Zhang and Palacios 2024). The Beta-coalescent model
 215 has a single parameter $\alpha \in [0, 2]$ and is defined as:

$$\Lambda(dx) = \frac{x^{1-\alpha} (1-x)^{\alpha-1}}{B(2-\alpha, \alpha)} dx \quad (27)$$

216 By combining Equations 26 and 27 we deduce that:

$$\lambda_{n,k} = \frac{B(k-\alpha, n-k+\alpha)}{B(2-\alpha, \alpha)} \quad (28)$$

217 Special cases of the Beta-coalescent include $\alpha = 2$ corresponding to the Kingman coalescent, $\alpha = 1$
 218 which is known as the Bolthausen-Sznitman coalescent and $\alpha = 0$ for which the phylogeny is always
 219 star-shaped.

220 We now define a new Lambda-coalescent based on the Negative-Binomial case described previously.
 221 We call this new Lambda-coalescent model the Omega-coalescent (where Omega stands for outbreak).
 222 For ease of comparison with other coalescent models, we consider that time is continuous and
 223 that the population size remains constant equal to $N_t = N$. The exclusive coalescent probability
 224 $p_{n,k,t}(N_t, N_{t+1})$ in the Negative-Binomial case given by Equation 21 can be used to determine the
 225 corresponding rate of the Omega-coalescent, if we consider that the probability of each event in discrete
 226 time is equal to the constant rate of this event happening in continuous time:

$$\lambda_{n,k} = p_{n,k,t}(N_t = N, N_{t+1} = N) = \frac{NB(k+r, n-k+Nr-r)}{B(r, Nr-r)} \quad (29)$$

227 Note that this equation implies that continuous time is measured approximately in number of
 228 transmission generations. For example to measure time in decimal days instead, the time scale would
 229 need to be multiplied by the mean of the generation time distribution measured in days (Svensson
 230 2007). From Equations 26 and 29 we can deduce the Lambda measure associated with the Omega-
 231 coalescent:

$$\Lambda(dx) = \frac{Nx^{r+1}(1-x)^{Nr-r-1}}{B(r, Nr-r)}dx \quad (30)$$

232 For a Lambda-coalescent model to be consistent, when a multiple merger of size k amongst n lineages
 233 occurs, if an additional lineage is revealed it must either take part in the multiple merger or remain
 234 unaffected (Berestycki 2009; Miró Pina et al. 2023). This implies that the rates must satisfy:

$$\lambda_{n,k} = \lambda_{n+1,k} + \lambda_{n+1,k+1} \quad (31)$$

235 This consistency property is easily verified for the Beta-coalescent in Equation 28 and likewise for the
 236 Omega-coalescent in Equation 29, in both cases using recursive properties of the Beta functions used
 237 in the respective definitions.

238 The Omega-coalescent has two parameters: the constant population size N and the dispersion
 239 parameter r . In order to compare the Omega-coalescent defined in Equation 29 with other models
 240 such as the Beta-coalescent defined in Equation 28, we consider the distribution of the size k of the
 241 next event among a set of n lineages. For any Lambda-coalescent this can be computed as:

$$p(k|n) = \frac{\binom{n}{k}\lambda_{n,k}}{\sum_{i=2}^n \binom{n}{i}\lambda_{n,i}} \quad (32)$$

242 Figure 3 compares this distribution for $n = 10$ in the Beta-coalescent with parameter $\alpha \in \{0.5, 1, 1.5\}$
 243 and for the Omega-coalescent with parameters $N \in \{10, 20, 30\}$ and $r \in \{0.1, 1, 10\}$. In the Beta-
 244 coalescent, the distribution shifts towards more larger multiple merger events as the parameter α
 245 decreases. In the Omega-coalescent a wider range of behaviours is obtained when varying the two
 246 parameters N and r . For a given value of N , decreasing the value of r results in more larger events.
 247 Conversely, for a given value of r we can see that increasing the value of N reduces the probability of
 248 larger events.

249 Genealogies can be simulated from the Omega-coalescent model defined in Equation 29 using the same
 250 algorithm as for other Lambda-coalescent models (Pitman 1999). Given n lineages, the next coalescent

Figure 3: Distribution of the size of the next event among a set of $n = 10$ lineages, compared between the Beta-coalescent and the Omega-coalescent model with various parameters.




Figure 4: Example of trees simulated under the Omega-coalescent with $r = 0.1$ (A), $r = 1$ (B), $r = 10$ (C) and $r = 100$ (D).




Figure 5: Summary statistics for trees simulated under the Omega-coalescent with $r = 0.1$, $r = 1$, $r = 10$ and $r = 100$, namely number of multiple mergers (A) the size of the largest multiple merger (B), the time to the most recent common ancestor (C) and the stemminess (D).

251 event happens after a time that is exponentially distributed with rate $\sum_{i=2}^n \binom{n}{i} \lambda_{n,i}$, the size k of this
 252 event is drawn according to Equation 32, and the k lineages that coalesce are chosen uniformly amongst
 253 the n lineages. This process is repeated iteratively until all lineages have coalesced. Figure 4 shows
 254 examples of trees simulated for a sample of size $n = 20$, constant population size $N = 30$ and dispersion
 255 parameter $r \in \{0.1, 1, 10, 100\}$. It is already clear from these single realisations that the lower values
 256 of r result in trees with more larger multiple merger events and lower time to the most recent common
 257 ancestor, but to quantify these properties we need to consider many trees. Figure 5 shows summary
 258 statistics for 10,000 trees simulated in the same conditions as the individual trees shown in Figure 4.
 259 As the dispersion parameter increases from $r = 0.1$ to $r = 100$ multiple merger events become less
 260 and less likely and less large (Figure 5A and B), and the time to the most recent common ancestor
 261 increases (Figure 5C). Furthermore, the stemminess of the tree increases, which is defined as the sum
 262 of lengths of internal branches divided by the total sum of branch lengths (Figure 5D). Stemminess is
 263 usually taken as a sign of population size dynamics (Fiala and Sokal 1985; Didelot et al. 2009), which
 264 would be misleading here since all simulations assumed a constant population size.

265 7 Parameter inference

266 Let us now consider a genealogy T with n leaves and c coalescent nodes, with $t_0 = 0$ the sampling time,
 267 t_1, \dots, t_c the times of the coalescent nodes in increasing order and k_i the number of lineages coalescing
 268 at time t_i . The number of lineages existing between time t_{i-1} and t_i is then $n_i = n - \sum_{j=1}^{i-1} k_j$. Under
 269 a Lambda-coalescent model, the genealogy T has likelihood:

$$p(T|\Lambda) = \prod_{i=1}^c \lambda_{n_i, k_i} \exp \left(- \sum_{j=2}^{n_i} \binom{n_i}{j} \lambda_{n_i, j} (t_i - t_{i-1}) \right) \quad (33)$$

270 Note that in Equation 33 the term $\binom{n_i}{k_i}$ term from the coalescent rate cancels out with its reciprocal
 271 from the probability of sampling k_i specific lineages to coalesce within a set of n_i . Estimating the
 272 Lambda measure from Equation 26 in general is a difficult problem (Koskela 2018; Miró Pina et al.
 273 2023). Here however we focus on estimation under the Omega-coalescent model, where the $\lambda_{n,k}$
 274 terms are given by Equation 29. There are therefore two parameters to estimate which have direct
 275 and important biological meaning: the effective population size N (which remains constant) and the
 276 dispersion parameter r of the Negative-Binomial offspring distribution. We perform estimation simply
 277 by maximising the likelihood in Equation 33, using the Brent algorithm (Brent 1971) when estimating

278 a single parameter and the L-BFGS-B algorithm (Byrd et al. 1995) when estimating both parameters.

279 We simulated 100 genealogies from the Omega-coalescent model each of which had $n = 100$ leaves,
280 with parameter N drawn uniformly at random between 100 and 500 and parameter r drawn uniformly
281 at random between 0.01 and 2. If we assume knowledge of the dispersion parameter, then estimating
282 the population size works really well (Figure 6A). Conversely we obtain good result when estimating
283 the dispersion parameter given a known population size (Figure 6B). However, attempting to estimate
284 both parameters at the same time performed significantly less well (Figures 6C and D). To illustrate
285 the cause of this, we consider a simulation for which the true parameters were $N = 200$ and $r = 0.5$,
286 and we construct the likelihood surface (Figure 6E). This shows a strong inverse tradeoff between the
287 two parameters, which is why it is harder to infer both parameters jointly. This poor identifiability
288 is analogous to the situation of a large population following the Cannings models (Cannings 1974).
289 In this case the coalescent process is fully determined by the effective population size $N_e = N/\sigma^2$
290 as previously noted (Kingman 1982a), where N is the population size and σ^2 is the variance in the
291 number of offspring. Consequently there is a full tradeoff between N and σ^2 , so that the ratio N_e can
292 be estimated but not the parameters N and σ^2 separately.

293 8 Implementation

294 We implemented the analytical methods described in this paper in a new R package entitled *EpiLambda*
295 which is available at <https://github.com/xavierdidelot/EpiLambda> for R version 3.5 or later. All
296 code and data needed to replicate the results are included in the “run” directory of the *EpiLambda*
297 repository. The R package **ape** was used to store, manipulate and visualise phylogenetic trees (Paradis
298 and Schliep 2019).

299 9 Discussion

300 We have described an ancestral process for infectious diseases which is relevant to the analysis of
301 outbreaks of a relatively small size, and to diseases with transmission heterogeneity. We have shown
302 how this process can be incorporated into a new Lambda-coalescent which we called the Omega-
303 coalescent. We only considered the situation where all samples are taken at the same time, but the
304 Omega-coalescent could be extended to allow temporally offset leaves following similar work on the




Figure 6: Maximum likelihood estimation of parameters. (A) Estimation of the population size given the dispersion parameter. (B) Estimation of the dispersion parameter given the population size. (C and D) Joint estimation of both the population size and dispersion parameters. (E) Example of likelihood surface as a function of both parameters.

305 coalescent (Drummond et al. 2003) and the Beta-coalescent (Hoscheit and Pybus 2019). We also made
306 the simplifying assumption of a constant population size, but this could be relaxed following the same
307 approach as previously described for integrating variable population size into the coalescent (Griffiths
308 and Tavaré 1994; Pybus et al. 2000; Ho and Shapiro 2011) and the Beta-coalescent (Hoscheit and
309 Pybus 2019; Zhang and Palacios 2024). Allowing the population size to vary could be especially useful
310 for the Omega-coalescent for several reasons. Firstly, since it is aimed at relatively small outbreaks,
311 it is likely that their sizes varies significantly. Secondly, the probability of multiple merger events of
312 various sizes depends explicitly on the population size in Equation 21. Changes in population size will
313 therefore have an effect on the distribution of events observed, as can be seen for example in Figure
314 3. Thirdly, joint inference of a varying population size could help break the otherwise difficult joint
315 inference of a fixed population size with the dispersion parameter (Figure 6).

316 We compared the Omega-coalescent only to the Beta-coalescent (Schweinsberg 2003) in Figure 3
317 as it is the model that has been most frequently used for infectious diseases (Hoscheit and Pybus
318 2019; Menardo et al. 2021; Helekal et al. 2025). Several other Lambda-coalescent models have been
319 proposed previously, such as the Dirac coalescent (Eldon and Wakeley 2006), the Durrett-Schweinsberg
320 coalescent (Durrett and Schweinsberg 2005) or the extended Beta-coalescent (Helekal et al. 2025).
321 However, none of these models is equivalent to the Omega-coalescent model. Indeed these previously
322 described Lambda-coalescent models are mostly concerned with situations where an individual can be
323 the father of a significant portion of a population in spite of the population being large, as opposed to
324 the small populations with superspreading we considered here. The xi-coalescent models are extensions
325 to the Lambda-coalescent models that admit multiple simultaneous mergers (Schweinsberg 2000). This
326 is clearly relevant to our basic discrete time model for small outbreaks, since in small populations it
327 is quite likely that separate subsets of individuals have the same infector in the previous generation.
328 However the exact timing of ancestry events is never available so that we must rely on ancestral dating
329 estimation with no notion of event co-occurrence (Volz and Frost 2017; Didelot et al. 2018; Bouckaert
330 et al. 2019; Helekal et al. 2025). We therefore introduced a continuous time approximation in Equation
331 29 so that ancestry events do not co-occur. The exact coalescent process for the discrete time Wright-
332 Fisher process in a small population has been previously described (Fu 2006). It would be difficult
333 however to extend this approach to the more complex forward-in-time model we considered here, with
334 variable population size and specific offspring distribution, which further justifies our continuous time
335 approximation.

336 Finally, it should be noted that our model describes the transmission tree during an outbreak, which is
337 different from a phylogeny (Jombart et al. 2011). This difference is often ignored and in some settings it

338 might be appropriate to do so, but not always. Consequently, some previous studies have used models
339 of within-host evolution to bridge the gap between transmission and phylogenetic trees (Didelot et al.
340 2014; Hall et al. 2015; Didelot et al. 2017). However, these models assume that each transmission
341 event happens independently from one infector to each of its infectees. This is not necessarily true
342 especially when considering superspreading events in which many individuals can become infected
343 simultaneously (Riley et al. 2003; Wallinga and Teunis 2004; Ho et al. 2023; Craddock et al. 2025).
344 In conclusion, we have described a new ancestral model for infectious disease outbreaks, which we
345 hope will be useful especially in settings where the outbreaks are small or in the presence of high
346 transmission heterogeneity.

347 **Acknowledgements**

348 We acknowledge funding from the National Institute for Health Research (NIHR) Health Protection
349 Research Unit in Genomics and Enabling Data.

References

- Anderson, R.M., May, R.M., 1991. *Infectious Diseases of Humans: Dynamics and Control*. Oxford University Press, USA.
- Berestycki, N., 2009. Recent progress in coalescent theory. *arXiv* , 0909.3985.
- Bouckaert, R., Vaughan, T.G., Fourment, M., Gavryushkina, A., Heled, J., Denise, K., Maio, N.D., Matschiner, M., Ogilvie, H., Plessis, L., Poppinga, A., 2019. BEAST 2.5 : An Advanced Software Platform for Bayesian Evolutionary Analysis. *PLoS computational biology* 15, e1006650.
- Brent, R.P., 1971. An algorithm with guaranteed convergence for finding a zero of a function. *The computer journal* 14, 422–425.
- Byrd, R.H., Lu, P., Nocedal, J., Zhu, C., 1995. A limited memory algorithm for bound constrained optimization. *SIAM Journal on scientific computing* 16, 1190–1208.
- Cannings, C., 1974. The latent roots of certain Markov chains arising in genetics: a new approach, I. Haploid models. *Adv. Appl. Probab.* 6, 260–290.
- Craddock, H., Spencer, S.E., Didelot, X., 2025. A bayesian modelling framework with model comparison for epidemics with super-spreading. *arXiv* , 2501.12768.
- Didelot, X., Croucher, N.J., Bentley, S.D., Harris, S.R., Wilson, D.J., 2018. Bayesian inference of ancestral dates on bacterial phylogenetic trees. *Nucleic Acids Research* 46, e134–e134.
- Didelot, X., Fraser, C., Gardy, J., Colijn, C., 2017. Genomic infectious disease epidemiology in partially sampled and ongoing outbreaks. *Molecular Biology and Evolution* 34, 997–1007.
- Didelot, X., Gardy, J., Colijn, C., 2014. Bayesian inference of infectious disease transmission from whole genome sequence data. *Molecular Biology and Evolution* 31, 1869–1879.
- Didelot, X., Urwin, R., Maiden, M.C.J., Falush, D., 2009. Genealogical typing of *Neisseria meningitidis*. *Microbiology* 155, 3176–86.
- Donnelly, P., Kurtz, T.G., 1999. Particle Representations for Measure-Valued Population Models. *The Annals of Probability* 27.
- Drummond, A.J., Pybus, O.G., Rambaut, A., Forsberg, R., Rodrigo, A.G., 2003. Measurably evolving populations. *Trends in Ecology and Evolution* 18, 481–488.

377 Du, Z., Wang, C., Liu, C., Bai, Y., Pei, S., Adam, D.C., Wang, L., Wu, P., Lau, E.H.Y., Cowling,
 378 B.J., 2022. Systematic review and meta-analyses of superspreading of SARS-CoV-2 infections.
 379 *Transboundary and Emerging Diseases* 69.

380 Durrett, R., Schweinsberg, J., 2005. A coalescent model for the effect of advantageous mutations on
 381 the genealogy of a population. *Stochastic Processes and their Applications* 115, 1628–1657.

382 Eldon, B., Wakeley, J., 2006. Coalescent Processes When the Distribution of Offspring Number Among
 383 Individuals Is Highly Skewed. *Genetics* 172, 2621–2633.

384 Ferguson, N.M., Cummings, D.A.T., Fraser, C., Cajka, J.C., Cooley, P.C., Burke, D.S., 2006. Strategies
 385 for mitigating an influenza pandemic. *Nature* 442, 448–452.

386 Fiala, K.L., Sokal, R.R., 1985. Factors determining the accuracy of cladogram estimation: Evolution
 387 using computer simulation. *Evolution* 39, 609–622.

388 Fisher, R.A., 1930. *The genetical theory of natural selection*. Clarendon Press.

389 Fraser, C., Li, L.M., 2017. Coalescent models for populations with time-varying population sizes and
 390 arbitrary offspring distributions. *bioRxiv* , 10.1101/131730.

391 Fraser, C., Riley, S., Anderson, R.M., Ferguson, N.M., 2004. Factors that make an infectious disease
 392 outbreak controllable. *Proceedings of the National Academy of Sciences* 101, 6146–6151.

393 Fu, Y.X., 2006. Exact coalescent for the Wright–Fisher model. *Theoretical Population Biology* 69,
 394 385–394.

395 Gómez-Carballa, A., Pardo-Seco, J., Bello, X., Martínón-Torres, F., Salas, A., 2021. Superspreading
 396 in the emergence of COVID-19 variants. *Trends in Genetics* 37, 1069–1080.

397 Grassly, N.C., Fraser, C., 2008. Mathematical models of infectious disease transmission. *Nature*
 398 *Reviews Microbiology* 6, 477–87.

399 Griffiths, R.C., Tavaré, S., 1994. Sampling theory for neutral alleles in a varying environment.
 400 *Philosophical Transactions of the Royal Society B* 344, 403–410.

401 Hall, M., Woolhouse, M., Rambaut, A., 2015. Epidemic Reconstruction in a Phylogenetics Framework:
 402 Transmission Trees as Partitions of the Node Set. *PLOS Computational Biology* 11, e1004613.

403 Heledal, D., Koskela, J., Didelot, X., 2025. Inference of multiple mergers while dating a pathogen
 404 phylogeny. *Systematic Biology* , syaf003.

405 Ho, F., Parag, K.V., Adam, D.C., Lau, E.H.Y., Cowling, B.J., Tsang, T.K., 2023. Accounting for the
406 Potential of Overdispersion in Estimation of the Time-varying Reproduction Number. *Epidemiology*
407 34, 201–205.

408 Ho, S.Y.W., Shapiro, B., 2011. Skyline-plot methods for estimating demographic history from
409 nucleotide sequences. *Molecular Ecology Resources* 11, 423–434.

410 Hoscheit, P., Pybus, O.G., 2019. The multifurcating skyline plot. *Virus Evolution* 5, 1–10.

411 Jombart, T., Eggo, R.M., Dodd, P.J., Balloux, F., 2011. Reconstructing disease outbreaks from genetic
412 data: A graph approach. *Heredity* 106, 383–90.

413 Keeling, M.J., Rohani, P., 2008. Modeling infectious diseases in humans and animals. Princeton
414 university press.

415 Kingman, J., 1982a. The coalescent. *Stochastic Processes and their Applications* 13, 235–248.

416 Kingman, J.F.C., 1982b. On the genealogy of large populations. *Journal of Applied Probability* 19,
417 27–43.

418 Koelle, K., Rasmussen, D.A., 2012. Rates of coalescence for common epidemiological models at
419 equilibrium. *Journal of The Royal Society Interface* 9, 997–1007.

420 Koskela, J., 2018. Multi-locus data distinguishes between population growth and multiple merger
421 coalescents. *Statistical Applications in Genetics and Molecular Biology* 17, 1–24.

422 Kucharski, A.J., Althaus, C.L., 2015. The role of superspreading in Middle East respiratory syndrome
423 coronavirus (MERS-CoV) transmission. *Eurosurveillance* 20, 14–18.

424 Lemieux, J.E., Siddle, K.J., Shaw, B.M., Loreth, C., Schaffner, S.F., Gladden-Young, A., Adams,
425 G., Fink, T., Tomkins-Tinch, C.H., Krasilnikova, L.A., DeRuff, K.C., Rudy, M., Bauer, M.R.,
426 Lagerborg, K.A., Normandin, E., Chapman, S.B., Reilly, S.K., Anahtar, M.N., Lin, A.E., Carter,
427 A., Myhrvold, C., Kembell, M.E., Chaluvadi, S., Cusick, C., Flowers, K., Neumann, A., Cerrato,
428 F., Farhat, M., Slater, D., Harris, J.B., Branda, J.A., Hooper, D., Gaeta, J.M., Baggett, T.P.,
429 O’Connell, J., Gnirke, A., Lieberman, T.D., Philippakis, A., Burns, M., Brown, C.M., Luban, J.,
430 Ryan, E.T., Turbett, S.E., LaRocque, R.C., Hanage, W.P., Gallagher, G.R., Madoff, L.C., Smole, S.,
431 Pierce, V.M., Rosenberg, E., Sabeti, P.C., Park, D.J., MacInnis, B.L., 2021. Phylogenetic analysis
432 of SARS-CoV-2 in Boston highlights the impact of superspreading events. *Science* 371, eabe3261.

433 Li, L.M., Grassly, N.C., Fraser, C., 2017. Quantifying Transmission Heterogeneity Using Both
434 Pathogen Phylogenies and Incidence Time Series. *Molecular Biology and Evolution* 34, 2982–2995.

435 Lloyd-Smith, J., Schreiber, S., Kopp, P., Getz, W., 2005. Superspreading and the effect of individual
436 variation on disease emergence. *Nature* 438, 355–9.

437 Menardo, F., Gagneux, S., Freund, F., 2021. Multiple Merger Genealogies in Outbreaks of
438 *Mycobacterium tuberculosis*. *Molecular Biology and Evolution* 38, 290–306.

439 Miró Pina, V., Joly, É., Siri-Jégousse, A., 2023. Estimating the Lambda measure in multiple-merger
440 coalescents. *Theoretical Population Biology* 154, 94–101.

441 Moran, P., 1958. Random Processes in Genetics. *Mathematical Proceedings of the Cambridge*
442 *Philosophical Society* 54, 60–71.

443 Paradis, E., Schliep, K., 2019. Ape 5.0: An environment for modern phylogenetics and evolutionary
444 analyses in R. *Bioinformatics* 35, 526–528.

445 Pitman, J., 1999. Coalescents with multiple collisions. *The Annals of Probability* 27, 1870–1902.

446 Potts, R.B., 1953. Note on the Factorial Moments of Standard Distributions. *Australian Journal of*
447 *Physics* 6, 498–499.

448 Pybus, O.G., Rambaut, A., Harvey, P.H., 2000. An integrated framework for the inference of viral
449 population history from reconstructed genealogies. *Genetics* 155, 1429–1437.

450 Riley, S., Fraser, C., a Donnelly, C., Ghani, A.C., Abu-Raddad, L.J., Hedley, A.J., Leung, G.M., Ho,
451 L.M., Lam, T.H., Thach, T.Q., Chau, P., Chan, K.P., Lo, S.V., Leung, P.Y., Tsang, T., Ho, W., Lee,
452 K.H., Lau, E.M.C., Ferguson, N.M., Anderson, R.M., 2003. Transmission dynamics of the etiological
453 agent of SARS in Hong Kong: Impact of public health interventions. *Science* 300, 1961–6.

454 Sagitov, S., 1999. The general coalescent with asynchronous mergers of ancestral lines. *Journal of*
455 *Applied Probability* 36, 1116–1125.

456 Schweinsberg, J., 2000. Coalescents with Simultaneous Multiple Collisions. *Electronic Journal of*
457 *Probability* 5.

458 Schweinsberg, J., 2003. Coalescent processes obtained from supercritical Galton–Watson processes.
459 *Stochastic Processes and their Applications* 106, 107–139.

460 Stein, R.A., 2011. Super-spreaders in infectious diseases. *International Journal of Infectious Diseases*
461 15, e510–e513.

462 Svensson, A., 2007. A note on generation times in epidemic models. *Mathematical Biosciences* 208(1),
463 300–311.

464 Tripathi, R.C., Gupta, R.C., Gurland, J., 1994. Estimation of parameters in the beta binomial model.
465 *Annals of the Institute of Statistical Mathematics* 46, 317–331.

466 Volz, E.M., 2012. Complex population dynamics and the coalescent under neutrality. *Genetics* 190,
467 187–201.

468 Volz, E.M., Frost, S.D.W., 2017. Scalable relaxed clock phylogenetic dating. *Virus Evolution* 3, vex025.

469 Wallinga, J., Teunis, P., 2004. Different Epidemic Curves for Severe Acute Respiratory Syndrome
470 Reveal Similar Impacts of Control Measures. *American Journal of Epidemiology* 160, 509–516.

471 Wang, J., Chen, X., Guo, Z., Zhao, S., Huang, Z., Zhuang, Z., Wong, E.L.y., Zee, B.C.Y., Chong,
472 M.K.C., Wang, M.H., Yeoh, E.K., 2021. Superspreading and heterogeneity in transmission of SARS,
473 MERS, and COVID-19: A systematic review. *Computational and Structural Biotechnology Journal*
474 19, 5039–5046.

475 Wang, L., Didelot, X., Yang, J., Wong, G., Shi, Y., Liu, W., Gao, G.F., Bi, Y., 2020. Inference of
476 person-to-person transmission of COVID-19 reveals hidden super-spreading events during the early
477 outbreak phase. *Nature Communications* 11, 5006.

478 Woolhouse, M.E.J., Dye, C., Etard, J.F., Smith, T., Charlwood, J.D., Garnett, G.P., Hagan, P., Hii,
479 J.L.K., Ndhlovu, P.D., Quinnell, R.J., Watts, C.H., Chandiwana, S.K., Anderson, R.M., 1997.
480 Heterogeneities in the transmission of infectious agents: Implications for the design of control
481 programs. *Proceedings of the National Academy of Sciences* 94, 338–342.

482 Wright, S., 1931. Evolution in Mendelian populations. *Genetics* 16, 97–159.

483 Zhang, J., Palacios, J.A., 2024. Multiple merger coalescent inference of effective population size. *arXiv*
484 , 2407.14976.