Chapter 2 Analytic Function

Problem 1

f(z)=f(x+iy)=u(x,y)+iv(x,y) 且 $u,v\in C^{(n)}$, 求 f(z)n 阶可导的 Cauchy-Riemann 条件和 $f^{(n)}(z)$ Solution

设 f'(z) = A + iB,则

$$df = f'(z)dz = f'(z)(dx + idy) \Leftrightarrow df = du + idv$$

$$= \frac{\partial u}{\partial x}dx + \frac{\partial u}{\partial y}dy + i(\frac{\partial v}{\partial x}dx + \frac{\partial u}{\partial y}dy)$$

$$= (Adx - Bdy) + i(Bdx + Ady)$$

由上式得

$$\begin{cases} Adx - Bdy = \frac{\partial u}{\partial x}dx + \frac{\partial u}{\partial y}dy \\ Bdx + Ady = \frac{\partial v}{\partial x}dx + \frac{\partial u}{\partial y}dy \end{cases}$$

解得

$$\begin{cases} \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} = A \\ \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} = -B \end{cases}$$

即

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = F'(z)$$

而

$$F'(z) = \frac{\partial U}{\partial x} + i \frac{\partial V}{\partial x} = \frac{\partial^2 u}{\partial x^2} + i \frac{\partial^2 v}{\partial x^2}$$

由归纳法可证明

$$f^{(n)}(z) = \frac{\partial^n u}{\partial x^n} + i \frac{\partial^n v}{\partial x^n}$$

u,v 需要满足 Cauchy-Riemann 条件

$$\begin{cases} \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \\ \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \end{cases}$$

Problem 2

求 cos(x+iy) 的实部和虚部, 其中 $x,y \in \mathbb{R}$

Solution

$$\begin{aligned} \cos(x+iy) &= \frac{1}{2}(e^{-y+ix} + e^{y-ix}) \\ &= \frac{1}{2}(e^{-y}e^{ix} + e^{y}e^{-ix}) \\ &= \frac{1}{2}[e^{-y}(\cos x + i\sin x) + e^{y}(\cos x - i\sin x)] \\ &= \frac{1}{2}(e^{y} + e^{-y})\cos x + i\frac{1}{2}(-e^{y} + e^{-y})\sin x \end{aligned}$$