

A Hotelling-Downs game for strategic candidacy with binary issues

Javier Esteban Maass Martínez, Vincent Mousseau & Anaëlle Wilczynski

Universidad de Chile – Laboratoire MICS

January 11, 2023



CentraleSupélec

1 Motivation

2 The Model

The Setting

Payoff and Stability Notions

3 Some Results

Negative Results in the general case

4 Restricting the Setting

Distribution of Voters

Candidates' Strategies

5 Some Empirical Insight

Bibliography

The Hotelling-Downs Model

Model of an election as a *political spectrum*. Existence of Equilibria in such a game has been widely studied (e.g. ice-cream vendors).

The Hotelling-Downs Model

Model of an election as a *political spectrum*. Existence of Equilibria in such a game has been widely studied (e.g. ice-cream vendors).

Precedents in the literature

- **Facility Location:** Placing facilities in a metric space to attract customers. More recently, on general graphs (see [Chan et al., 2021]).

The Hotelling-Downs Model

Model of an election as a *political spectrum*. Existence of Equilibria in such a game has been widely studied (e.g. ice-cream vendors).

Precedents in the literature

- **Facility Location:** Placing facilities in a metric space to attract customers. More recently, on general graphs (see [Chan et al., 2021]).
- **Voronoi Games:** Attract closest points in space. Studied in general Graphs and (interestingly) on Transitive Graphs [Feldmann et al., 2009]

The Hotelling-Downs Model

Model of an election as a *political spectrum*. Existence of Equilibria in such a game has been widely studied (e.g. ice-cream vendors).

Precedents in the literature

- **Facility Location:** Placing facilities in a metric space to attract customers. More recently, on general graphs (see [Chan et al., 2021]).
- **Voronoi Games:** Attract closest points in space. Studied in general Graphs and (interestingly) on Transitive Graphs [Feldmann et al., 2009]
- **Strategic Candidacy:** candidates strategically abstain from an election (to achieve another winner they may prefer). [Dutta et al., 2001].

The Hotelling-Downs Model

Model of an election as a *political spectrum*. Existence of Equilibria in such a game has been widely studied (e.g. ice-cream vendors).

Precedents in the literature

- **Facility Location:** Placing facilities in a metric space to attract customers. More recently, on general graphs (see [Chan et al., 2021]).
- **Voronoi Games:** Attract closest points in space. Studied in general Graphs and (interestingly) on Transitive Graphs [Feldmann et al., 2009]
- **Strategic Candidacy:** candidates strategically abstain from an election (to achieve another winner they may prefer). [Dutta et al., 2001].
- **Hotelling Downs Model's** connection to COMSOC only recently studied: *Real Candidacy Games* [Sabato et al., 2017] & specially a 2021 paper by [Harrenstein et al., 2021].


The Hotelling-Downs Model

Model of an election as a *political spectrum*. Existence of Equilibria in such a game has been widely studied (e.g. ice-cream vendors).

Precedents in the literature

- **Facility Location:** Placing facilities in a metric space to attract customers. More recently, on general graphs (see [Chan et al., 2021]).
- **Voronoi Games:** Attract closest points in space. Studied in general Graphs and (interestingly) on Transitive Graphs [Feldmann et al., 2009]
- **Strategic Candidacy:** candidates strategically abstain from an election (to achieve another winner they may prefer). [Dutta et al., 2001].
- **Hotelling Downs Model's** connection to COMSOC only recently studied: *Real Candidacy Games* [Sabato et al., 2017] & specially a 2021 paper by [Harrenstein et al., 2021].
- **Judgment Aggregation:** Aggregating *logical propositions*. Inspiration for *binary issue setting* and *single-peakedness* notion (see [Puppe, 2018]).

The Model



The Setting

- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.

The Setting

- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.
- This population is interested by $K \in \mathbb{N}$ relevant binary issues: their *opinions* can be represented as a vector in the *hypercube* $\mathcal{H} := \{0, 1\}^K$. (Equivalently, we have a *distribution of voters* $f_N : \mathcal{H} \rightarrow \mathbb{N}$)

The Setting

- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.
- This population is interested by $K \in \mathbb{N}$ relevant binary issues: their *opinions* can be represented as a vector in the *hypercube* $\mathcal{H} := \{0, 1\}^K$. (Equivalently, we have a *distribution of voters* $f_N : \mathcal{H} \rightarrow \mathbb{N}$)
- Consider the *Hamming Distance*: $\text{dist}(p, p') = |\{j \in [K] : p_j \neq p'_j\}|$.

- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.
- This population is interested by $K \in \mathbb{N}$ relevant binary issues: their *opinions* can be represented as a vector in the *hypercube* $\mathcal{H} := \{0, 1\}^K$. (Equivalently, we have a *distribution of voters* $f_N : \mathcal{H} \rightarrow \mathbb{N}$)
- Consider the *Hamming Distance*: $\text{dist}(p, p') = |\{j \in [K] : p_j \neq p'_j\}|$.

Binary Strategic Candidacy Game (BSC)


- Candidates announce a *position* $s_i \in \mathcal{H}_i$.
- Strategy Profile: $\mathbf{s} = (s_1, \dots, s_m)$
- Voters will prefer candidates whose announced *opinion* is closest to their own ($p_v \in \mathcal{H}$) $\rightarrow \succsim_v^{\mathbf{s}}$ weak order over C .
- Candidates run for an election whose winner is determined by a voting rule $\mathcal{F} : (\mathcal{W}(C))^n \rightarrow C$ (with tie-breaking \triangleleft)

The Setting

- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.
- This population is interested by $K \in \mathbb{N}$ relevant binary issues: their *opinions* can be represented as a vector in the *hypercube* $\mathcal{H} := \{0, 1\}^K$. (Equivalently, we have a *distribution of voters* $f_N : \mathcal{H} \rightarrow \mathbb{N}$)
- Consider the *Hamming Distance*: $\text{dist}(p, p') = |\{j \in [K] : p_j \neq p'_j\}|$.

Binary Strategic Candidacy Game (BSC)


- Candidates announce a *position* $s_i \in \mathcal{H}_i$.
- Strategy Profile: $\mathbf{s} = (s_1, \dots, s_m)$
- Voters will prefer candidates whose announced *opinion* is closest to their own ($p_v \in \mathcal{H}$) $\rightarrow \succsim_v^{\mathbf{s}}$ weak order over C .
- Candidates run for an election whose winner is determined by a voting rule $\mathcal{F} : (\mathcal{W}(C))^n \rightarrow C$ (with tie-breaking \triangleleft)



- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.
- This population is interested by $K \in \mathbb{N}$ relevant binary issues: their *opinions* can be represented as a vector in the *hypercube* $\mathcal{H} := \{0, 1\}^K$. (Equivalently, we have a *distribution of voters* $f_N : \mathcal{H} \rightarrow \mathbb{N}$)
- Consider the *Hamming Distance*: $\text{dist}(p, p') = |\{j \in [K] : p_j \neq p'_j\}|$.

Binary Strategic Candidacy Game (BSC)

- Candidates announce a *position* $s_i \in \mathcal{H}_i$.
- Strategy Profile: $\mathbf{s} = (s_1, \dots, s_m)$
- Voters will prefer candidates whose announced *opinion* is closest to their own ($p_v \in \mathcal{H}$) $\rightarrow \succsim_v^{\mathbf{s}}$ weak order over C .
- Candidates run for an election whose winner is determined by a voting rule $\mathcal{F} : (\mathcal{W}(C))^n \rightarrow C$ (with tie-breaking \triangleleft)




The Setting

- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.
- This population is interested by $K \in \mathbb{N}$ relevant binary issues: their *opinions* can be represented as a vector in the *hypercube* $\mathcal{H} := \{0, 1\}^K$. (Equivalently, we have a *distribution of voters* $f_N : \mathcal{H} \rightarrow \mathbb{N}$)
- Consider the *Hamming Distance*: $\text{dist}(p, p') = |\{j \in [K] : p_j \neq p'_j\}|$.

Binary Strategic Candidacy Game (BSC)

- Candidates announce a *position* $s_i \in \mathcal{H}_i$.
- Strategy Profile: $\mathbf{s} = (s_1, \dots, s_m)$
- Voters will prefer candidates whose announced *opinion* is closest to their own ($p_v \in \mathcal{H}$) $\rightarrow \succsim_v^{\mathbf{s}}$ weak order over C .
- Candidates run for an election whose winner is determined by a voting rule $\mathcal{F} : (\mathcal{W}(C))^n \rightarrow C$ (with tie-breaking \triangleleft)




The Setting

- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.
- This population is interested by $K \in \mathbb{N}$ relevant binary issues: their *opinions* can be represented as a vector in the *hypercube* $\mathcal{H} := \{0, 1\}^K$. (Equivalently, we have a *distribution of voters* $f_N : \mathcal{H} \rightarrow \mathbb{N}$)
- Consider the *Hamming Distance*: $\text{dist}(p, p') = |\{j \in [K] : p_j \neq p'_j\}|$.

Binary Strategic Candidacy Game (BSC)

- Candidates announce a *position* $s_i \in \mathcal{H}_i$.
- Strategy Profile: $\mathbf{s} = (s_1, \dots, s_m)$
- Voters will prefer candidates whose announced *opinion* is closest to their own ($p_v \in \mathcal{H}$) $\rightarrow \succsim_v^{\mathbf{s}}$ weak order over C .
- Candidates run for an election whose winner is determined by a voting rule $\mathcal{F} : (\mathcal{W}(C))^n \rightarrow C$ (with tie-breaking \triangleleft)




The Setting

- Let's consider $N = [n]$ **voters** and $C = \{c_1, c_2, \dots, c_m\}$ **candidates**.
- This population is interested by $K \in \mathbb{N}$ relevant binary issues: their *opinions* can be represented as a vector in the *hypercube* $\mathcal{H} := \{0, 1\}^K$. (Equivalently, we have a *distribution of voters* $f_N : \mathcal{H} \rightarrow \mathbb{N}$)
- Consider the *Hamming Distance*: $\text{dist}(p, p') = |\{j \in [K] : p_j \neq p'_j\}|$.

Binary Strategic Candidacy Game (BSC)

- Candidates announce a *position* $s_i \in \mathcal{H}_i$.
- Strategy Profile: $\mathbf{s} = (s_1, \dots, s_m)$
- Voters will prefer candidates whose announced *opinion* is closest to their own ($p_v \in \mathcal{H}$) $\rightarrow \succsim_v^{\mathbf{s}}$ weak order over C .
- Candidates run for an election whose winner is determined by a voting rule $\mathcal{F} : (\mathcal{W}(C))^n \rightarrow C$ (with tie-breaking \triangleleft)



Payoff and Stability Notions

Candidates don't necessarily want to win !

- Each one has a **weak ranking** over its rivals with themselves at the top.
- A **better response** for c_i from \mathbf{s} is $\mathbf{s}'_i \in \mathcal{H}_i$ s.t. $\mathcal{F}((\mathbf{s}'_i, \mathbf{s}_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.

Candidates don't necessarily want to win !

- Each one has a **weak ranking** over its rivals with themselves at the top.
- A **better response** for c_i from \mathbf{s} is $\mathbf{s}'_i \in \mathcal{H}_i$ s.t. $\mathcal{F}((\mathbf{s}'_i, \mathbf{s}_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.

Nash Equilibrium (NE)

A state $\mathbf{s} \in \prod_{i=1}^m \mathcal{H}_i$ is a Nash equilibrium if there is no strategy $\mathbf{s}'_i \in \mathcal{H}_i$ for a candidate $c_i \in C$ such that $\mathcal{F}((\mathbf{s}'_i, \mathbf{s}_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.

Candidates don't necessarily want to win !

- Each one has a **weak ranking** over its rivals with themselves at the top.
- A **better response** for c_i from \mathbf{s} is $\mathbf{s}'_i \in \mathcal{H}_i$ s.t. $\mathcal{F}((\mathbf{s}'_i, \mathbf{s}_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.

Nash Equilibrium (NE)

A state $\mathbf{s} \in \prod_{i=1}^m \mathcal{H}_i$ is a Nash equilibrium if there is no strategy $\mathbf{s}'_i \in \mathcal{H}_i$ for a candidate $c_i \in C$ such that $\mathcal{F}((\mathbf{s}'_i, \mathbf{s}_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.

t -local Equilibrium

A state $\mathbf{s} \in \prod_{i=1}^m \mathcal{H}_i$ is a t -local equilibrium if there is no strategy $\mathbf{s}'_i \in \mathcal{H}_i$ for a candidate $c_i \in C$ such that $\text{dist}(\mathbf{s}'_i, \mathbf{s}_i) \leq t$ and $\mathcal{F}((\mathbf{s}'_i, \mathbf{s}_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.

Candidates don't necessarily want to win !

- Each one has a **weak ranking** over its rivals with themselves at the top.
- A **better response** for c_i from \mathbf{s} is $s'_i \in \mathcal{H}_i$ s.t. $\mathcal{F}((s'_i, s_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.

Nash Equilibrium (NE)




A state $\mathbf{s} \in \prod_{i=1}^m \mathcal{H}_i$ is a Nash equilibrium if there is no strategy $s'_i \in \mathcal{H}_i$ for a candidate $c_i \in C$ such that $\mathcal{F}((s'_i, s_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.

t -local Equilibrium

A state $\mathbf{s} \in \prod_{i=1}^m \mathcal{H}_i$ is a t -local equilibrium if there is no strategy $s'_i \in \mathcal{H}_i$ for a candidate $c_i \in C$ such that $\text{dist}(s'_i, s_i) \leq t$ and $\mathcal{F}((s'_i, s_{-i})) \succ_{c_i} \mathcal{F}(\mathbf{s})$.


A Nash equilibrium is equivalent to a K -local equilibrium. Also, a t -local equilibrium is a t' -local equilibrium for every $1 \leq t' \leq t \leq K$.

Some Results




Negative Results in the general case

A 1-local equilibrium may fail to exist even with $m = 2$, $K = 3$



		$s_2 \in \mathcal{H}_2$	
		$(1, 0, 1)$	$(1, 1, 1)$
$s_1 \in \mathcal{H}_1$	$(0, 1, 0)$	$(1, \mathbf{2})$	$\xleftarrow{c_2} (\mathbf{1.5}, 1.5)$
	$(0, 1, 1)$	$(\mathbf{1.5}, 1.5)$	$\xrightarrow{c_2} (1, \mathbf{2})$

A 1-local equilibrium may fail to exist even with $m = 2$, $K = 3$



		$s_2 \in \mathcal{H}_2$	
		$(1, 0, 1)$	$(1, 1, 1)$
$s_1 \in \mathcal{H}_1$	$(0, 1, 0)$	$(1, \mathbf{2})$ $c_1 \downarrow$	$\xleftarrow{c_2} (\mathbf{1.5}, 1.5)$ $\uparrow c_1$
	$(0, 1, 1)$	$(\mathbf{1.5}, 1.5)$	$\xrightarrow{c_2} (\mathbf{1}, \mathbf{2})$

NP-Hardness of the decision problem

Deciding whether there exists a t -local equilibrium is NP-hard, for $t \in \{2, \dots, K\}$, even under *narcissistic* preferences.

We make a reduction of *Exact Cover by 3-Sets (X3C)*, which is NP Complete:

Some structure of the $m = 2$ case

Influence Sets (for $\mathbf{s} = (s_1, s_2)$)

- $P_i^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) < \text{dist}(p, s_{-i})\} \rightarrow p_v \in P_i^{\mathbf{s}} \Leftrightarrow c_i \succ_v^{\mathbf{s}} c_{-i}$
- $I^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) = \text{dist}(p, s_{-i})\} \rightarrow p_v \in I^{\mathbf{s}} \Leftrightarrow c_i \sim_v^{\mathbf{s}} c_{-i}$
- $\mathcal{H} = P_1^{\mathbf{s}} \cup P_2^{\mathbf{s}} \cup I^{\mathbf{s}}$.
- Let $r^{\mathbf{s}} := \text{dist}(s_1, s_2)$ and $d_{r^{\mathbf{s}}} := \lceil \frac{r^{\mathbf{s}}}{2} \rceil - 1$ (*critical distance* of influence).
- $p \in P_i^{\mathbf{s}} \Leftrightarrow \text{dist}_{\neq}^{\mathbf{s}}(p, s_i) \leq d_{r^{\mathbf{s}}}$ and $p \in I^{\mathbf{s}} \Leftrightarrow r^{\mathbf{s}}$ is even & $\text{dist}_{\neq}^{\mathbf{s}}(p, s_i) = \frac{r^{\mathbf{s}}}{2}$.

Some structure of the $m = 2$ case

Influence Sets (for $\mathbf{s} = (s_1, s_2)$)


- $P_i^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) < \text{dist}(p, s_{-i})\} \rightarrow p_v \in P_i^{\mathbf{s}} \Leftrightarrow c_i \succ_v^{\mathbf{s}} c_{-i}$
- $I^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) = \text{dist}(p, s_{-i})\} \rightarrow p_v \in I^{\mathbf{s}} \Leftrightarrow c_i \sim_v^{\mathbf{s}} c_{-i}$
- $\mathcal{H} = P_1^{\mathbf{s}} \cup P_2^{\mathbf{s}} \cup I^{\mathbf{s}}$.
- Let $r^{\mathbf{s}} := \text{dist}(s_1, s_2)$ and $d_{r^{\mathbf{s}}} := \lceil \frac{r^{\mathbf{s}}}{2} \rceil - 1$ (*critical distance of influence*).
- $p \in P_i^{\mathbf{s}} \Leftrightarrow \text{dist}_{\neq}^{\mathbf{s}}(p, s_i) \leq d_{r^{\mathbf{s}}}$ and $p \in I^{\mathbf{s}} \Leftrightarrow r^{\mathbf{s}}$ is even & $\text{dist}_{\neq}^{\mathbf{s}}(p, s_i) = \frac{r^{\mathbf{s}}}{2}$.

Dynamic Aspect

When doing a 1 – local deviation, a candidate can only win/lose influence over positions that *agree/disagree* with him in the deviated issue.

Antipodality

Influence sets are *antipodal*: $P_i^{\mathbf{s}} = \check{P}_{-i}^{\mathbf{s}}$



Some structure of the $m = 2$ case

Influence Sets (for $\mathbf{s} = (s_1, s_2)$)


- $P_i^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) < \text{dist}(p, s_{-i})\} \rightarrow p_v \in P_i^{\mathbf{s}} \Leftrightarrow c_i \succ_v^{\mathbf{s}} c_{-i}$
- $I^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) = \text{dist}(p, s_{-i})\} \rightarrow p_v \in I^{\mathbf{s}} \Leftrightarrow c_i \sim_v^{\mathbf{s}} c_{-i}$
- $\mathcal{H} = P_1^{\mathbf{s}} \cup P_2^{\mathbf{s}} \cup I^{\mathbf{s}}$.
- Let $r^{\mathbf{s}} := \text{dist}(s_1, s_2)$ and $d_{r^{\mathbf{s}}} := \lceil \frac{r^{\mathbf{s}}}{2} \rceil - 1$ (*critical distance of influence*).
- $p \in P_i^{\mathbf{s}} \Leftrightarrow \text{dist}_{\neq}^{\mathbf{s}}(p, s_i) \leq d_{r^{\mathbf{s}}}$ and $p \in I^{\mathbf{s}} \Leftrightarrow r^{\mathbf{s}}$ is even & $\text{dist}_{\neq}^{\mathbf{s}}(p, s_i) = \frac{r^{\mathbf{s}}}{2}$.

Dynamic Aspect

When doing a 1 – local deviation, a candidate can only win/lose influence over positions that *agree/disagree* with him in the deviated issue.

Antipodality

Influence sets are *antipodal*: $P_i^{\mathbf{s}} = \check{P}_{-i}^{\mathbf{s}}$



Some structure of the $m = 2$ case

Influence Sets (for $\mathbf{s} = (s_1, s_2)$)


- $P_i^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) < \text{dist}(p, s_{-i})\} \rightarrow p_v \in P_i^{\mathbf{s}} \Leftrightarrow c_i \succ_v^{\mathbf{s}} c_{-i}$
- $I^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) = \text{dist}(p, s_{-i})\} \rightarrow p_v \in I^{\mathbf{s}} \Leftrightarrow c_i \sim_v^{\mathbf{s}} c_{-i}$
- $\mathcal{H} = P_1^{\mathbf{s}} \cup P_2^{\mathbf{s}} \cup I^{\mathbf{s}}$.
- Let $r^{\mathbf{s}} := \text{dist}(s_1, s_2)$ and $d_{r^{\mathbf{s}}} := \lceil \frac{r^{\mathbf{s}}}{2} \rceil - 1$ (critical distance of influence).
- $p \in P_i^{\mathbf{s}} \Leftrightarrow \text{dist}_{\neq}^{\mathbf{s}}(p, s_i) \leq d_{r^{\mathbf{s}}}$ and $p \in I^{\mathbf{s}} \Leftrightarrow r^{\mathbf{s}}$ is even & $\text{dist}_{\neq}^{\mathbf{s}}(p, s_i) = \frac{r^{\mathbf{s}}}{2}$.

Dynamic Aspect

When doing a 1 – local deviation, a candidate can only win/lose influence over positions that *agree/disagree* with him in the deviated issue.

Antipodality

Influence sets are *antipodal*: $P_i^{\mathbf{s}} = \check{P}_{-i}^{\mathbf{s}}$



Some structure of the $m = 2$ case

Influence Sets (for $\mathbf{s} = (s_1, s_2)$)


- $P_i^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) < \text{dist}(p, s_{-i})\} \rightarrow p_v \in P_i^{\mathbf{s}} \Leftrightarrow c_i \succ_v^{\mathbf{s}} c_{-i}$
- $I^{\mathbf{s}} := \{p \in \mathcal{H} : \text{dist}(p, s_i) = \text{dist}(p, s_{-i})\} \rightarrow p_v \in I^{\mathbf{s}} \Leftrightarrow c_i \sim_v^{\mathbf{s}} c_{-i}$
- $\mathcal{H} = P_1^{\mathbf{s}} \cup P_2^{\mathbf{s}} \cup I^{\mathbf{s}}$.
- Let $r^{\mathbf{s}} := \text{dist}(s_1, s_2)$ and $d_{r^{\mathbf{s}}} := \lceil \frac{r^{\mathbf{s}}}{2} \rceil - 1$ (critical distance of influence).
- $p \in P_i^{\mathbf{s}} \Leftrightarrow \text{dist}_{\neq}^{\mathbf{s}}(p, s_i) \leq d_{r^{\mathbf{s}}}$ and $p \in I^{\mathbf{s}} \Leftrightarrow r^{\mathbf{s}}$ is even & $\text{dist}_{\neq}^{\mathbf{s}}(p, s_i) = \frac{r^{\mathbf{s}}}{2}$.

Dynamic Aspect

When doing a 1 – local deviation, a candidate can only win/lose influence over positions that *agree/disagree* with him in the deviated issue.

Antipodality

Influence sets are *antipodal*: $P_i^{\mathbf{s}} = \check{P}_{-i}^{\mathbf{s}}$



Some Positive Results

Let $p^m \in \mathcal{H}$ be defined, $\forall j \in \mathcal{H}$ as: $(p^m)_j = \arg \max_{e \in \{0,1\}} f_N(\mathcal{H} \mid_{j=e})$. i.e.
 p^m captures the *majoritarian view* on each issue.

Some Positive Results

Let $p^m \in \mathcal{H}$ be defined, $\forall j \in \mathcal{H}$ as: $(p^m)_j = \arg \max_{e \in \{0,1\}} f_N(\mathcal{H} \mid_{j=e})$. i.e. p^m captures the *majoritarian view* on each issue.

Existence of 2-local equilibria

There always exists a 2-local equilibrium in the BSC game when $m = 2$, n is odd, and $p^m \in \mathcal{H}_1 \cap \mathcal{H}_2$.

Unfortunately...

A 3-local equilibrium may not exist, even when $m = 2$, $K = 3$, and the sets of candidates' strategies coincide, contain p^m and are connected.

Some Positive Results


Let $p^m \in \mathcal{H}$ be defined, $\forall j \in \mathcal{H}$ as: $(p^m)_j = \arg \max_{e \in \{0,1\}} f_N(\mathcal{H}|_{j=e})$. i.e. p^m captures the *majoritarian view* on each issue.

Existence of 2-local equilibria

There always exists a 2-local equilibrium in the BSC game when $m = 2$, n is odd, and $p^m \in \mathcal{H}_1 \cap \mathcal{H}_2$.


Unfortunately...

A 3-local equilibrium may not exist, even when $m = 2$, $K = 3$, and the sets of candidates' strategies coincide, contain p^m and are connected.



		$s_2 \in \mathcal{H}_2$			
		(0,0,0)	(1,0,0)	(1,1,0)	(1,1,1)
$s_1 \in \mathcal{H}_1$	(0,0,0)	(30.5 , 30.5)	(30, 31)	(30, 31)	(31 , 30)
	(1,0,0)	(31 , 30)	(30.5 , 30.5)	(30, 31)	(30, 31)
	(1,1,0)	(31 , 30)	(31 , 30)	(30.5 , 30.5)	(30, 31)
	(1,1,1)	(30, 31)	(31 , 30)	(31 , 30)	(30.5 , 30.5)

Restricting the Setting



SinglePeaked Distribution of voters


A distribution of voters $f : \mathcal{H} \rightarrow \mathbb{N}$ is SP if:

$\exists p^* \in \mathcal{H}; \forall x, y \in \mathcal{H} :$

$$y \in [x, p^*] \implies f(x) \leq f(y)$$

General agreement around the opinion p^ .*

(An example of a SP distribution is the *uniform*)



SinglePeaked Distribution of voters


A distribution of voters $f : \mathcal{H} \rightarrow \mathbb{N}$ is SP if:

$\exists p^* \in \mathcal{H}; \forall x, y \in \mathcal{H} :$

$$y \in [x, p^*] \implies f(x) \leq f(y)$$

General agreement around the opinion p^ .*

(An example of a SP distribution is the *uniform*)



Existence of Nash Equilibrium under **uniform** distribution

When $m = 2$ and the distribution of voters is **uniform**, **every state** is a NE.

SinglePeaked Distribution of voters


A distribution of voters $f : \mathcal{H} \rightarrow \mathbb{N}$ is SP if:

$\exists p^* \in \mathcal{H}; \forall x, y \in \mathcal{H} :$

$$y \in [x, p^*] \implies f(x) \leq f(y)$$

General agreement around the opinion p^ .*

(An example of a SP distribution is the *uniform*)



Existence of Nash Equilibrium under **uniform** distribution

When $m = 2$ and the distribution of voters is **uniform**, **every state** is a NE.

Theorem: NE existence under SP distributions

When $m = 2$, under a single-peaked distribution of voters, if $p^* \in \mathcal{H}_1$, there always exists a Nash equilibrium in the BSC game !

SinglePeaked Distribution of voters


A distribution of voters $f : \mathcal{H} \rightarrow \mathbb{N}$ is SP if:

$\exists p^* \in \mathcal{H}; \forall x, y \in \mathcal{H} :$

$$y \in [x, p^*] \implies f(x) \leq f(y)$$

General agreement around the opinion p^ .*

(An example of a SP distribution is the *uniform*)



Existence of Nash Equilibrium under **uniform** distribution

When $m = 2$ and the distribution of voters is **uniform**, **every state** is a NE.

Theorem: NE existence under SP distributions

When $m = 2$, under a single-peaked distribution of voters, if $p^* \in \mathcal{H}_1$, there always exists a Nash equilibrium in the BSC game !

It's actually stronger: c_1 **always wins** when taking the peak

[Betweenness Matching]

For any $\mathbf{s} = (s_1, s_2) \in H_1 \times H_2$ s.t. $s_{i_0} = p^* \neq s = s_{-i_0}$,

$\exists \phi : \mathcal{H}_s \rightarrow \mathcal{H}_p$ **bijection** , s.t.: $\forall x \in \mathcal{H}_i(\mathbf{s}), \phi(x) \in [x, p^*]$




Every $x \in \mathcal{H}_s$ is mapped to exactly one $\phi(x) \in \mathcal{H}_p$ *between* x and the peak.

[Betweenness Matching]

For any $\mathbf{s} = (s_1, s_2) \in H_1 \times H_2$ s.t. $s_{i_0} = p^* \neq s = s_{-i_0}$,

$\exists \phi : \mathcal{H}_s \rightarrow \mathcal{H}_p$ **bijection** , s.t.: $\forall x \in \mathcal{H}_i(\mathbf{s}), \phi(x) \in [x, p^*]$

Every $x \in \mathcal{H}_s$ is mapped to exactly one $\phi(x) \in \mathcal{H}_p$ *between* x and the peak.






[Betweenness Matching]

For any $\mathbf{s} = (s_1, s_2) \in H_1 \times H_2$ s.t. $s_{i_0} = p^* \neq s = s_{-i_0}$,

$\exists \phi : \mathcal{H}_s \rightarrow \mathcal{H}_p$ **bijection** , s.t.: $\forall x \in \mathcal{H}_i(\mathbf{s}), \phi(x) \in [x, p^*]$

Every $x \in \mathcal{H}_s$ is mapped to exactly one $\phi(x) \in \mathcal{H}_p$ *between* x and the peak.



This is achieved using Hall's Theorem's Corollary for regular graphs. The problem is reduced to the *antipodal* case and decomposed by *layers*.

Unfortunately, NOT scalable !


NOT even a 1-local equilibrium !

A 1-local equilibrium may not exist in the populism game even when $m = 3$, $K = 2$, the candidates' preferences are fixed, and the distribution of voters is uniform.

Unfortunately, NOT scalable !

NOT even a 1-local equilibrium !

A 1-local equilibrium may not exist in the populism game even when $m = 3$, $K = 2$, the candidates' preferences are fixed, and the distribution of voters is uniform.



C_1 :	C_1	\succ	C_3	\succ	C_2
C_2 :	C_2	\succ	C_1	\succ	C_3
C_3 :	C_3	\succ	C_2	\succ	C_1

Seeing the Game Table, we notice that players have a 1-local deviation from every possible state:

		$s_2 \in \mathcal{H}_2$				$s_2 \in \mathcal{H}_2$	
		(0,0)	(1,0)			(0,0)	(1,0)
$s_1 \in \mathcal{H}_1$	(0,0)	$(w, w, 2w)$ $c_1 \downarrow$	$(w, \frac{3}{2}w, \frac{3}{2}w)$ $\downarrow c_1$	$s_1 \in \mathcal{H}_1$	(0,0)	$(\frac{7}{6}w, \frac{7}{6}w, \frac{5}{3}w)$ $c_1 \uparrow$	$(\frac{3}{2}w, w, \frac{3}{2}w)$ $\uparrow c_1$
	(1,0)	$(\frac{3}{2}w, w, \frac{3}{2}w)$	$(\frac{7}{6}w, \frac{7}{6}w, \frac{5}{3}w)$		(1,0)	$(w, \frac{3}{2}w, \frac{3}{2}w)$	$(w, w, 2w)$
		$s_3 = (0,1)$				$s_3 = (1,1)$	

Theorem: 1-local by following your rival

There always exists a 1-local equilibrium in the BSC game when $m = 2$ and $\mathcal{H}_2 \subseteq \mathcal{H}_1$. Such an equilibrium can be found in polynomial time.

Theorem: 1-local by following your rival

There always exists a 1-local equilibrium in the BSC game when $m = 2$ and $\mathcal{H}_2 \subseteq \mathcal{H}_1$. Such an equilibrium can be found in polynomial time.

Theorem: 1-local for balls

There always exists a 1-local equilibrium in the BSC game when $m = 2$ and candidates' strategies are **balls of radius one**. Such an equilibrium can be found in polynomial time.


Theorem: 1-local by following your rival

There always exists a 1-local equilibrium in the BSC game when $m = 2$ and $\mathcal{H}_2 \subseteq \mathcal{H}_1$. Such an equilibrium can be found in polynomial time.

Theorem: 1-local for balls


There always exists a 1-local equilibrium in the BSC game when $m = 2$ and candidates' strategies are **balls of radius one**. Such an equilibrium can be found in polynomial time.

2-local eq. may NOT exist with $m = 2$, $K = 3$, and \mathcal{H}_i balls of radius one




		$s_2 \in \mathcal{H}_2$			
		(1,0,0)	(0,0,0)	(1,0,1)	(1,1,0)
$s_1 \in \mathcal{H}_1$	(0,0,0)	(4,5)	(4.5,4.5)	(4,5)	(4,5)
	(1,0,0)	(4.5,4.5)	(5,4)	(4,5)	(4,5)
	(0,1,0)	(4.5,4.5)	(5,4)	(5,4)	(4,5)
	(0,0,1)	(4.5,4.5)	(5,4)	(4,5)	(5,4)

Some Empirical Insight



Empirical Insight: Local Equilibria




Iterative Games : t -local Dynamic

By turns, candidates change their current strategy for a t -local improving move. It will stop at a t -local equilibrium!


Iterative Games : t -local Dynamic

By turns, candidates change their current strategy for a t -local improving move. It will stop at a t -local equilibrium!




Iterative Games : t -local Dynamic

By turns, candidates change their current strategy for a t -local improving move. It will stop at a t -local equilibrium!




Iterative Games : t -local Dynamic

By turns, candidates change their current strategy for a t -local improving move. It will stop at a t -local equilibrium!





Iterative Games : t -local Dynamic

By turns, candidates change their current strategy for a t -local improving move. It will stop at a t -local equilibrium!





Iterative Games : t -local Dynamic

By turns, candidates change their current strategy for a t -local improving move. It will stop at a t -local equilibrium!





Iterative Games : t -local Dynamic

By turns, candidates change their current strategy for a t -local improving move. It will stop at a t -local equilibrium!




Iterative Games : t -local Dynamic

By turns, candidates change their current strategy for a t -local improving move. It will stop at a t -local equilibrium!



Empirical Insight: Dynamics





Thank you for your attention !

- [Chan et al., 2021] Chan, H., Filos-Ratsikas, A., Li, B., Li, M., and Wang, C. (2021).
Mechanism design for facility location problem: A survey.
In Proceedings of the 30th International Joint Conference on Artificial Intelligence (IJCAI-21), pages 1–17.
- [Dutta et al., 2001] Dutta, B., Jackson, M. O., and Breton, M. L. (2001).
Strategic candidacy and voting procedures.
Econometrica, 69(4):1013–1037.
- [Feldmann et al., 2009] Feldmann, R., Mavronicolas, M., and Monien, B. (2009).
Nash equilibria for voronoi games on transitive graphs.
In Proceedings of the 5th International Workshop on Internet and Network Economics (WINE-09), pages 280–291.
- [Harrenstein et al., 2021] Harrenstein, P., Lisowski, G., Sridharan, R., and Turrini, P. (2021).
A hotelling-downs framework for party nominees.
In Proceedings of the 21st International Conference on Autonomous Agents and MultiAgent Systems, page 593–601.
- [Puppe, 2018] Puppe, C. (2018).
The single-peaked domain revisited: A simple global characterization.
Journal of Economic Theory, 176:55–80.
- [Sabato et al., 2017] Sabato, I., Obratzsova, S., Rabinovich, Z., and Rosenschein, J. S. (2017).
Real candidacy games: A new model for strategic candidacy.
In Proceedings of the 16th Conference on Autonomous Agents and MultiAgent Systems, AAMAS '17, page 867–875, Richland, SC. International Foundation for Autonomous Agents and Multiagent Systems.

Lemma: [d-regular bipartite graph matching] - Uses Hall's Theorem

A graph $G = (V, E)$ is said to be d -regular if $\forall v \in V$, $\deg_G(v) = d$ (i.e. all vertices have the same degree d).

We always have that **any d -regular bipartite graph has an X -perfect matching**.

Lemma

Let $r \in \{1, \dots, K\}$. Let $s = \vec{0}_r$ and $p = \vec{1}_r$, both in the Hypercube $\mathcal{H}^r = \{0, 1\}^r$. (Also, define $\mathcal{H}_s^r := \{x \in \mathcal{H}^r : d(x, s) < d(x, p)\}$, and \mathcal{H}_p^r in a similar way)

There **exists a bijection** $\phi : \mathcal{H}_s^r \rightarrow \mathcal{H}_p^r$ s.t. $\forall x \in \mathcal{H}_s^r$, $\phi(x) \in [x, p]$

Schema of Proof

Let $\ell \in \{0, \dots, c_r\}$, we consider

$$\mathcal{H}_s^r|_\ell := \{x \in \mathcal{H}_s^r : d(x, s) = \ell\} \quad \text{and} \quad \mathcal{H}_p^r|_\ell := \{y \in \mathcal{H}_p^r : d(y, p) = \ell\}.$$

The GOOD bipartite graph to check: $G|_\ell = (X + Y, E)$:

$$X = \mathcal{H}_s^r|_\ell, \quad Y = \mathcal{H}_p^r|_\ell, \quad E = \{(x, y) \in X \times Y : (x, y, p) \in \tau_r\}$$

We can see that $G|_\ell$ is **regular!** The degree of nodes in $G|_\ell$ is:

For $x \in X = \mathcal{H}_s^r|_\ell$,

$$\deg(x) = |\{(a, b) \in E : a = x\}| = |\{b \in Y : (x, b, p) \in \tau_r\}| = |[x, p] \cap \mathcal{H}_p^r|_\ell|.$$

We see that, by definition, this set is:

$$\begin{aligned} [x, p] \cap \mathcal{H}_p^r|_\ell &= \{z \in \mathcal{H}_p^r : d(z, \vec{1}_r) = \ell, \forall i \in \{1, \dots, r\} : [x_i = 1 \implies z_i = 1]\} \\ &= \{y \in \mathcal{H}_p^r : \exists A \subseteq \mathbb{I}_{x \equiv 0}, |A| = r - 2\ell; y = \mathbf{1}_{\mathbb{I}_{x \equiv 1}} + \sum_{i \in A} \mathbf{e}_i\} \end{aligned}$$

$$\text{So, we have: } \deg(x) = |[x, p] \cap \mathcal{H}_p^r|_\ell| = |\{A \subseteq \mathbb{I}_{x \equiv 0} : |A| = r - 2\ell\}| = \binom{r-\ell}{r-2\ell}$$

Similarly (as $(a, y, \vec{1}_r) \in \tau_r \iff (y, a, \vec{0}_r) \in \tau_r$):

$$\deg(y) = |\{(a, b) \in E : b = y\}| = |\{a \in X : (a, y, p) \in \tau_r\}| = |[y, s] \cap \mathcal{H}_s^r|_\ell.$$

And, of course, we notice:

$$\begin{aligned} [y, s] \cap \mathcal{H}_s^r|_\ell &= \{x \in \mathcal{H}_s^r : d(x, \vec{0}_r) = \ell, \forall i \in \{1, \dots, r\} : [y_i = 0 \implies x_i = 0]\} \\ &= \{x \in \mathcal{H}_s^r : \exists A \subseteq \mathbb{I}_{y=1}, |A| = \ell; x = 0_{\mathbb{I}_{y=0}} + \sum_{i \in A} e_i\} \end{aligned}$$

So that: $\deg(y) = |[y, s] \cap \mathcal{H}_s^r|_\ell = |\{A \subseteq \mathbb{I}_{y=1} : |A| = \ell\}| = \binom{r-\ell}{\ell}$

As $G|_\ell$ is a **regular bipartite graph**, by the Lemma, we have a **perfect matching between $\mathcal{H}_s|_\ell$ and $\mathcal{H}_r|_\ell$** (call it ϕ^ℓ)

We can properly define the following to have the bijection we want:

$$\begin{aligned} \phi : \mathcal{H}_s^r &\rightarrow \mathcal{H}_p^r \\ x &\mapsto \phi^\ell(x) \text{ if } d(x, \vec{0}_r) = \ell \end{aligned}$$

(bijective cause surjective and pidgeon principle; and $(x, \phi(x), p) \in \tau_r$ cause every ϕ^ℓ satisfy that).

From General to Particular

Let $s, p \in \mathcal{H} = \{0, 1\}^K$. We consider $\mathbb{I} := \{i : s_i = 1\}$ and define:

$$\psi : x \mapsto \psi(x) := (x_{\mathbb{I}^c}, (1 - x_i)_{\mathbb{I}})$$

This is such that: $\psi(s) = \vec{0}$, and for $\mathbb{J} = \{i : s_i = p_i\}$: $\psi(p) = (\vec{0}_{\mathbb{J}}, \vec{1}_{\mathbb{J}^c})$.

We can then partition the hypercube as:

$$\mathcal{H} = \{0, 1\}^K = \{0, 1\}^{\mathbb{J}} \times \{0, 1\}^{\mathbb{J}^c} = \bigcup_{a \in \{0, 1\}^{\mathbb{J}}} \{a\} \times \{0, 1\}^{\mathbb{J}^c}.$$

This is perfect to establish (as $\mathcal{H}_{\vec{s}}|_{\{a\} \times \{0, 1\}^{\mathbb{J}^c}} = \{a\} \times \mathcal{H}_{\vec{0}_r}^r$; ϕ from the lemma):

$$\begin{aligned} \phi^a : \{a\} \times \mathcal{H}_{\vec{0}_r}^r &\rightarrow \{a\} \times \mathcal{H}_{\vec{1}_r}^r \\ (a, z) &\mapsto \phi^a((a, z)) := (a, \phi(z)) \end{aligned}$$

And ultimately: $f : \mathcal{H}_{\vec{s}} \rightarrow \mathcal{H}_{\vec{p}}$ given by $x \mapsto \phi^{x_{\mathbb{J}}}(x)$. By finally conjugating with our original bijection of the hypercube, we come back to:

$F := \psi^{-1} \circ f \circ \psi : \mathcal{H}_{\vec{s}} \rightarrow \mathcal{H}_{\vec{p}}$ which satisfies all the properties we want
(**bijective** and $\forall x \in \mathcal{H}_{\vec{s}}, F(x) \in [x, p]$)

Another way of reducing to the *antipodal* case:

We can restrict our attention to issues where $s_1 \neq s_2$. Let $r = |s| = \text{dist}(s_1, s_2)$ and focus on a new BSC game BSC^r , where the sets of voters and candidates are the same as in BSC , and taking the hypercube $\mathcal{H}^r := \{0, 1\}^r$.

Assume, w.l.o.g., that s_1 and s_2 differ on the **first** r issues. We transform game BSC into game BSC^r by using the function $\phi : \mathcal{H} \rightarrow \mathcal{H}^r$ where $(\phi(p))_j = p_j$ for every $j \in \mathbf{s}$ and every position $p \in \mathcal{H}$.

It is easy to see that, by definition, $\phi(s_i) = \phi(\hat{s}_{-i})$ (**they are now antipodal!**) and $\text{dist}(\phi(s_i), \phi(s_{-i})) = r$. Also: $\forall p \in P_i^{\mathbf{s}}$, we have $\phi(p) \in P_i^{\phi(\mathbf{s})}$.

We denote by $w_p := f_N(p)$ and $w_{p^r} := f_N(p^r)$. In addition, we define $F(p^r) := \{p \in \mathcal{H} : \phi(p) = p^r\}$ for each $p^r \in \mathcal{H}^r$; and also w^r as follows: $w^r(p^r) := \sum_{p \in F(p^r)} w_p$. Hence, we have $sc(\phi(s_i)) = \sum_{p^r \in i\phi(\mathbf{s})} w_{p^r} = \sum_{p^r \in i\phi(\mathbf{s})} \sum_{p \in F(p^r)} w_p = sc(s_i)$ for every $i \in \{1, 2\}$. The distribution of voters in this new populism game is still single-peaked:

Lema

The distribution of voters in populism game BSC^r is single-peaked w.r.t. peak position $\phi(p^*)$.

X3C

In an instance of X3C, we are given a set $X = \{x_1, x_2, \dots, x_{3q}\}$ and a set $S = \{S_1, S_2, \dots, S_r\}$ of 3-element subsets of X and we ask whether there exists an exact cover, i.e., a subset $S' \subseteq S$ such that every element of X occurs in exactly one member of S' , in other words S' is a partition of X .

We construct our game as follows. We consider $K = 3q + 4$ issues, and we create $(3q + 10)w_p + 23$ voters, given an arbitrary integer w_p such that $w_p > 24$, where the voters are distributed as follows:

- w_p voters on each position $e^i = (0, \dots, 0, 1, 0, \dots, 0)$ such that $e_i^i = 1$ and $e_j^i = 0$ for every $j \in [3q + 4] \setminus \{i\}$, for every $i \in [3q]$;
- $\frac{5}{2}w_p + 11$ voters on position $p_1 := (0, \dots, 0, 1, 1, 0, 0)$;
- 7 voters on position $p_2 := (0, \dots, 0, 0, 0, 1, 1)$;
- $\frac{5}{2}w_p + 3$ voters on position $p_3 := (0, \dots, 0, 0, 0, 1, 0)$;
- 2 voters on position $p_4 := (0, \dots, 0, 0, 0, 0, 1)$.

- We create $q + 2$ candidates and denote the set of candidates by $C := C_S \cup \{c_a, c_b\}$, where the set $C_S := \bigcup_{j=1}^q c_j$ regroups the so-called subset-candidates.
- The sets of strategies are:
 - $\mathcal{H}_c := \mathcal{H}_S := \bigcup_{j=1}^q \{s^j = (s_1, \dots, s_{3q}, 0, 0, 0, 0) \in \{0, 1\}^K : \forall i \in [3q], s_i = 1 \text{ iff } x_i \in S_j\}$ for every $c \in C_S$;
 - $\mathcal{H}_{c_a} := \{s_a^1 := (0, \dots, 0, 1, 0, 0, 1), s_a^2 := (0, \dots, 0, 1, 1, 0, 0)\}$;
 - $\mathcal{H}_{c_b} := \{s_b^1 := (0, \dots, 0, 0, 0, 1, 1), s_b^2 := (0, \dots, 0, 1, 0, 1, 0)\}$.
- The candidates' truthful positions are arbitrary and their preferences are narcissistic.

One can prove that there exists a Nash equilibrium in the populism game iff there exists a subset of S that is a partition of X .

The idea is that only candidates c_a and c_b may have an incentive to deviate and they would do so only if there is a position e^i for $i \in [3q]$ not “covered” by the strategy position of a subset-candidate.

We report in the table the number of votes that candidates c_a and c_b can get from positions p_1, p_2, p_3 , and p_4 .

		\mathcal{H}_{c_b}	
		s_b^1	s_b^2
\mathcal{H}_{c_a}	s_a^1	$(\frac{5}{2}w_p + 12, \frac{5}{2}w_p + 11)$	$(\frac{5}{4}w_p + 11, \frac{15}{4}w_p + 12)$
	s_a^2	$(\frac{5}{2}w_p + 11, \frac{5}{2}w_p + 12)$	$(\frac{5}{2}w_p + 12, \frac{5}{2}w_p + 11)$

Table: Number of votes, from the voters whose truthful position is in $\{p_1, p_2, p_3, p_4\}$, that candidates c_a and c_b get according to all their possible strategies.

- A better response for candidate c_a or c_b would trigger a cycle of local deviations, preventing a Nash equilibrium to exist.
- Moreover, the only deviations that c_a or c_b can make are towards another strategy position at distance 2 from their previous strategy position.

It follows that the complexity result also holds for 2-local equilibria.